

Analysis of the vegetation structure in the savanna - tropical forest transition of northern South America using The Global Ecosystem Dynamics Investigation (GEDI) data

Santiago Valencia^{1,1}, Juan Camilo Villegas^{1,1}, and Juan F. Salazar^{1,1}

¹Universidad de Antioquia

January 20, 2023

Abstract

The savanna - forest transition in the tropics has a large and complex variation in vegetation structure both vertically and horizontally. 3D-imaging technologies provide detailed high-resolution measurements of the vegetation structure. However, the use of these observations globally faces practical challenges due to spatio-temporal gaps and operational restrictions, mainly in tropical regions. NASA's Global Ecosystem Dynamics Investigation (GEDI) is the first quasi-global LiDAR (light detection and ranging) observations of 3D vegetation structure at a footprint resolution of 25 m. Here we use GEDI data (GEDI02_Bv001) to analyze vegetation structure in the savanna - tropical forest transition of northern South America, using canopy height, canopy cover, total Plant Area Index (PAI), maximum Plant Area volume Density (PAVD), and vertical profile of PAI and PAVD as vegetation structure descriptors. Despite contrasts between savanna (open-canopy) and forest (closed-canopy), our results show a gradual variation along the transition in canopy height, canopy cover, total PAI, and maximum PAVD. Our results support that the savanna- forest transition in tropical regions can be described as a grassland - forest continuum. Results also indicate that GEDI data allow a better characterization of vegetation lower than 5 m in height, mainly in savanna, an improvement from other global databases (e.g. MODIS). Further, our study illustrates the potential of GEDI data to advance in the characterization of large-scale patterns of vegetation structure in tropics, key for supporting biogeography, and macroecology studies relevant in the phase of current ecosystem changes.

Analysis of vegetation structure in the savanna - tropical forest transition of northern South America using The Global Ecosystem Dynamics Investigation (GEDI) data

Analysis of vegetation structure in the savanna - tropical forest transition of northern South America using The Global Ecosystem Dynamics Investigation (GEDI) data
Santiago Valencia¹, Juan Camilo Villegas², Juan F. Salazar¹

¹ Grupo GIGA, Escuela Ambiental, Facultad de Ingeniería, Universidad de Antioquia, Medellín-Colombia
² Grupo de Investigación en Ecología Aplicada, Escuela Ambiental, Facultad de Ingeniería, Universidad de Antioquia, Medellín-Colombia

Motivation

The savanna - forest transition is the most widespread transition in tropical areas, which has a large and complex variation in vegetation structure both vertically and laterally (Oliver & Muller, 2016).

3D imaging technologies provide detailed high-resolution measurements of the vegetation structure. However, the use of these observations faces practical challenges due to spatial-temporal gaps (Villegas et al., 2020).

Results

Despite the apparent structural contrasts between savanna and forest and typical representation of transition as a sharp line, our results suggest a gradual transition in canopy cover, and most notably in structural characteristics indicated by PAI and PAVD profiles (Figure 3).




Figure 3. Canopy cover (left panel), PAI (middle panel), and PAVD (right panel) profiles (meters along transect). The vertical axis indicates height in meters. The horizontal axis indicates distance along the transect. The vertical axis is labeled 'm' and ranges from 0 to 100. The horizontal axis is labeled 'm' and ranges from 0 to 100. The canopy cover profile shows a gradual increase from savanna to forest. The PAI and PAVD profiles show a similar gradual transition, with PAI increasing and PAVD decreasing along the transect.

The below figures show canopy cover (Figure 4), PAI (Figure 5), and PAVD (Figure 6) increases along forest and savanna.

Discussion

The development of tools such as postglows (GEDI to R) (Shan et al., 2020) and pyGEDI (GEDI to Python) (Gómez & Torres, 2020) in Python facilitate the use of GEDI data, particularly for non-expert users.


Despite the high resolution of GEDI data only along inter-national space station trajectories for spatial analysis, additionally, low quality data decreases the number of available transects as shown in Figure 1b.

Our results highlight the importance of using structural descriptors such PAI and PAVD that provide more information about structures than simple canopy cover (Gómez et al., 2020).

Study area and data

In this study, we use GEDI data to analyze vegetation structure in the savanna - tropical forest transition of northern South America (Figure 1).

The savanna-forest transition zone differs according to the Latin American (Figure 1a, Clements et al., 2012). For this analysis, we selected 18 transects that cross the savanna (SL, ME), transition (TL, TE), and forest (FL, FE) portions of the transect. We use canopy height, canopy cover, and structure (Plant Area Index (PAI) and Plant Area Volume Density (PAVD)) profiles as vegetation structure descriptors.



Future work

Detailed GEDI data validation has shown high performance. However, it is necessary to combine with the validation, mainly in regions with substantial LIDAR data gaps such as northern South America.

Previous studies have identified differences in GEDI data performance between power vs. coverage losses and day vs. night observations, which require further analysis for our study area.

The fusion of GEDI data with other datasets (e.g. Landsat) and availability of ground GEDI products (SGL) could allow more detailed spatial-temporal analysis of vegetation structure, tracking their changes, and improved biomass estimations.


[CHATting](#) [AUTHOR INFORMATION](#) [ABSTRACT](#) [REFERENCES](#) [CONTACT AUTHOR](#) [PRINT](#) [GET POSTER](#)

Santiago Valencia¹, Juan Camilo Villegas², Juan F. Salazar¹

¹ Grupo GIGA, Escuela Ambiental, Facultad de Ingeniería, Universidad de Antioquia, Medellín-Colombia

² Grupo de Investigación en Ecología Aplicada, Escuela Ambiental, Facultad de Ingeniería, Universidad de Antioquia, Medellín-Colombia

PRESENTED AT:



MOTIVATION

- **The savanna – forest transition** is the most widespread ecotone in **tropical areas**, which has a **large and complex variation in vegetation structure** both vertically and horizontally (Oliveras & Malhi, 2016).
- **3D-imaging technologies** provide detailed high-resolution measurements of the **vegetation structure**. However, the use of these observations **faces practical challenges due to spatio-temporal gaps** (Valbuena et al., 2020).
- NASA's Global Ecosystem Dynamics Investigation (GEDI) is the **first quasi-global** (51.6°N-51.6°S) LiDAR (light detection and ranging) observations of 3D vegetation structure at a **footprint resolution of 25 m** (Dubayah et al., 2020).
- No studies have previously tested the capacity of this product to **detect and characterize ecotones and transitions in tropical regions**.

STUDY AREA AND DATA

In this study, we use GEDI data to analyze vegetation structure in the **savanna - tropical forest transition of northern South America** (Figure 1).

The savanna-forest transition was defined according to the **Llanos ecoregion** (Figure 1b, Dinerstein et al., 2017). For the analysis, we selected **18 transects** that cover the savanna (S1-S6), transition (T1-T6), and forest (F1-F6) portions of the ecotone. We use **canopy height, canopy cover, and cumulative Plant Area Index (PAI) and Plant Area Volume Density (PAVD)** profiles as vegetation structure descriptors.




Figure 1. (a) General location of the study area in the northern South America and (b) total available high-quality GEDI data of canopy cover (GEDI2B) over the study area. The brown line represents the limits of Llanos ecoregion (Dinerstein et al., 2017). The red boxes show the selected transect in savanna (S1 to S6), transition (T1 to T6), and forest (F1 to F6). The Google Earth imagery as the background.

As seen in Figure 2, GEDI data consists of 8 beam ground transects. In this study, we obtained data only for **one full power beam** (beam0101) excluding low-quality data. We download, extract, and process data with the **rGEDI package in R** (Silva et al., 2020).




Figure 2. Example of 8 beam ground transect. Red rows indicate the selected beam in this study and the brown line represents the limits of Llanos ecoregion (Dinerstein et al., 2017). The Google Earth imagery as the background.

RESULTS

Despite the apparent structural contrasts between savanna and forest and typical representation of transition as a sharp line, our results suggest a **gradual transition** in canopy cover, and most notably in structural characteristics indicated by PAI and PAVD profiles (Figure 3).




Figure 3. Canopy cover (left panel), PAI (middle panel), and PAVD (right panel) transects along transition. The vertical red and white dashed lines indicate the savanna – forest transition defined for the Llanos ecoregion. PAI and PAVD values were derived for 5 m vertical bins.

The below figures show canopy cover (Figure 4), PAI (Figure 5), and PAVD (Figure 6) transects along forest and savanna.




Figure 4. Canopy cover transects along forest (left panel, F1 to F6) and savanna (right panel, S1 to S6).

As expected, the savannas have low canopy cover (S1 to S6, Figure 4). However, **high canopy cover values along the savanna transects**, mainly associated riparian forest (Figure 2), fundamental for biodiversity in these systems.




Figure 5. PAI vertical profile as a function of canopy height transect along forest (left panel, F1 to F6) and savanna (right panel, S1 to S6). PAI values were derived for 5 m vertical bins.

The PAI and PAVD profiles show that footprints in savanna with similar canopy cover to the forest (> 80%), have lower canopy height and structural complexity, typical of savannas.




Figure 6. PAVD vertical profile as a function of canopy height transect along forest (left panel, F1 to F6) and savanna (right panel, S1 to S6). PAVD values were derived for 5 m vertical bins.

These differences between canopy cover and vertical profiles are also observed in the forest, where canopy cover shows high homogeneity, while PAVD and PAI highlight variations in canopy height and complexity structural along and between transects.

The PAVD profiles also indicate that savanna and forest have a greater density of vegetation in heights between 0-10 m, and 15-25 m, respectively.

DISCUSSION

The development of tools such as packages **rGEDI** in R (Silva et al., 2020) and **pyGEDI** (Serna & Serna, 2020) in Python **facilitate the use of GEDI data**, particularly for non-expert users.

Despite the high resolution of GEDI, **data only along international space station trajectories** limit spatial analyzes. Additionally, **low-quality data** decreases the number of available footprints as shown in Figure 1b.

Our results **highlight** the importance of using structural descriptors such **PAI and PAVD that provide more information about ecotones that simple canopy cover** (Marselis et al., 2018).

Our results support that **transition shows many shades of green** depending on the local biophysical factors (e.g. climate, soil, fire) and human-related disturbances (e.g. deforestation, degradation) (Oliveras & Malhi, 2016; Stark et al., 2020).

Multiple studies have highlighted that although **canopy cover** is useful in many cases, it can lead to **underestimating the structural complexity** of the vegetation. For example, **open biomes**, as savannas with typically low canopy cover, has been **wrongly classified** as **degraded forest** and priority areas for **restoration** (Silveira et al., 2020) solely based on a canopy cover criteria.

Our results support that the **savanna-forest transition in tropical regions** can be described as a **grassland - forest continuum** (Breshears, 2006), which contrasts with the **typical representation as a simple line in maps** (Marques et al., 2020). Considering a **transition zone instead of a line** is key to understand the **impacts of environmental change** (e.g. deforestation, degradation, fire) and **related ecosystems process** (Stark et al., 2020).

FUTURE WORK

Global GEDI data validation has show high performance. However, it is necessary to **continue with the validation**, mainly in regions with substantial LiDAR data gaps such as northern South America.

Previous studies have identified differences in **GEDI data performance between power vs. coverage beams and day vs. night observations**, which require further analysis for our study area.

The fusion of GEDI data with other databases (e.g. Landsat) and availability of gridded GEDI products (L3 & L4) would allow more detailed spatio-temporal analysis of vegetation structure, including their change, and improved biomass estimations.

Acknowledgment

This works was funded by the **Universidad de Antioquia** through the Estudiante Instructor Program for graduate studies. JCV and JFS were funded by the **Colombian Ministry of Science, Technology and Innovation (Minciencias)** through program "Sostenibilidad de sistemas ecológicos y sociales en la cuenca Magdalena-Cauca bajo escenarios de cambio climático y pérdida de bosques" (code 1115-852-70719) with funds from "Patrimonio Autónomo Fondo Nacional de Financiamiento para la Ciencia, la Tecnología y la Innovación, Fondo Francisco José de Caldas".

AUTHOR INFORMATION

Santiago Valencia

Master's student

Escuela Ambiental, Facultad de Ingeniería

Universidad de Antioquia

santiago.valencia8@udea.edu.co

Twitter: [@Sav_Cardenas](https://twitter.com/Sav_Cardenas)

ABSTRACT

The savanna - forest transition in the tropics has a large and complex variation in vegetation structure both vertically and horizontally. 3D-imaging technologies provide detailed high-resolution measurements of the vegetation structure. However, the use of these observations globally faces practical challenges due to spatio-temporal gaps and operational restrictions, mainly in tropical regions. NASA's Global Ecosystem Dynamics Investigation (GEDI) is the first quasi-global (51.6°N-51.6°S) LiDAR (light detection and ranging) observations of 3D vegetation structure at a footprint resolution of 25 m. Here we use GEDI data (GEDI02_Bv001) to analyze vegetation structure in the savanna - tropical forest transition of northern South America, using canopy height, canopy cover, and vertical profiles of Plant Area Index (PAI) and Plant Area Volume Density (PAVD) as vegetation structure descriptors. Despite contrasts between savanna (open-canopy) and forest (closed-canopy), our results show a gradual variation along the transition in canopy height, canopy cover, total PAI, and maximum PAVD. Our results support that the savanna- forest transition in tropical regions can be described as a grassland - forest continuum. Results also indicate that GEDI data allow a better characterization of vegetation lower than 5 m in height, mainly in savanna, an improvement from other global databases (e.g. MODIS). Further, our study illustrates the potential of GEDI data to advance in the characterization of large-scale patterns of vegetation structure in tropics, key for supporting biogeography, and macroecology studies relevant in the phase of current ecosystem changes.

REFERENCES

- Breshears, D. D. (2006). **The grassland–forest continuum: trends in ecosystem properties for woody plant mosaics?** *Frontiers in Ecology and the Environment*, 4(2), 96-104.
- Dubayah, R., Blair, J. B., Goetz, S., Fatoyinbo, L., Hansen, M., Healey, S., ... & Armston, J. (2020). **The Global Ecosystem Dynamics Investigation: High-resolution laser ranging of the Earth's forests and topography.** *Science of Remote Sensing*, 1, 100002.
- Dinerstein, E., Olson, D., Joshi, A., Vynne, C., Burgess, N. D., Wikramanayake, E., ... & Hansen, M. (2017). **An ecoregion-based approach to protecting half the terrestrial realm.** *BioScience*, 67(6), 534-545.
- Marques, E. Q., Marimon-Junior, B. H., Marimon, B. S., Matricardi, E. A., Mews, H. A., & Colli, G. R. (2020). **Redefining the Cerrado–Amazonia transition: implications for conservation.** *Biodiversity and conservation*, 29(5), 1501-1517.
- Marselis, S. M., Tang, H., Armston, J. D., Calders, K., Labrière, N., & Dubayah, R. (2018). **Distinguishing vegetation types with airborne waveform lidar data in a tropical forest-savanna mosaic: A case study in Lopé National Park, Gabon.** *Remote sensing of environment*, 216, 626-634.
- Oliveras, I., & Malhi, Y. (2016). **Many shades of green: the dynamic tropical forest–savannah transition zones.** *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371(1703), 20150308.
- Serna, E; Hernandez-Serna, A. **pyGEDI: NASA's Global Ecosystem Dynamics Investigation (GEDI) mission data extraction, analysis, processing and visualization.** version 0.2, April. 5th 2020, available at: <https://pypi.org/project/pyGEDI/>
- Silva, C.A; Hamamura,C.; Valbuena, R.; Hancock,S.; Cardil,A.; Broadbent, E. N.; Almeida,D.R.A.; Silva Junior, C.H.L; Klauberg, C. **rGEDI: NASA's Global Ecosystem Dynamics Investigation (GEDI) Data Visualization and Processing.** version 0.1.8, accessed on October. 22 2020, available at: <https://CRAN.R-project.org/package=rGEDI>
- Silveira, F. A., Arruda, A. J., Bond, W., Durigan, G., Fidelis, A., Kirkman, K., ... & Siebert, S. J. (2020). **Myth-busting tropical grassy biome restoration.** *Restoration Ecology*, 28(5), 1067-1073.
- Stark, S. C., Breshears, D. D., Aragón, S., Villegas, J. C., Law, D. J., Smith, M. N., ... & Saleska, S. R. (2020). **Reframing tropical savannization: linking changes in canopy structure to energy balance alterations that impact climate.** *Ecosphere*, 11(9), e03231.
- Valbuena, R., O'Connor, B., Zellweger, F., Simonson, W., Vihervaara, P., Maltamo, M., ... & Chirici, G. (2020). **Standardizing ecosystem morphological traits from 3D information sources.** *Trends in Ecology & Evolution*.