

Telomere erosion as an intrinsic mechanism of species extinction: the sudden and complete disappearance of the passenger pigeon 100 years ago

Reinhard Stindl¹

¹Alpharm GesmbH, apo-med-center, Plättenstr. 7-9, 2380 Perchtoldsdorf, Austria

April 17, 2023

Abstract

Correspondence re: Hung, C. M., et al. 2014. “Drastic population fluctuations explain the rapid extinction of the passenger pigeon.” *Proc Natl Acad Sci U S A* no. 111 (29):10636-41. doi: 10.1073/pnas.1401526111.



Telomere erosion as an intrinsic mechanism of species extinction: the sudden and complete disappearance of the passenger pigeon 100 years ago

REINHARD STINDL¹

1. Alpharm GesmbH, apo-med-center, Plättenstr. 7-9, 2380 Perchtoldsdorf, Austria

ABSTRACT

Correspondence re: Hung, C. M., et al. 2014. "Drastic population fluctuations explain the rapid extinction of the passenger pigeon." *Proc Natl Acad Sci U S A* no. 111 (29):10636-41. doi: 10.1073/pnas.1401526111.

READ REVIEWS

WRITE A REVIEW

CORRESPONDENCE:

rs@telomere.at

DATE RECEIVED:

June 11, 2015

DOI:

10.15200/winn.140984.49964

ARCHIVED:

September 04, 2014

KEYWORDS:

evolution, extinction, telomere erosion

CITATION:

Reinhard Stindl, Telomere erosion as an intrinsic mechanism of species extinction: the sudden and complete disappearance of the passenger pigeon 100 years ago, *The Winnower* 2:e140984.49964, 2014, DOI: 10.15200/winn.140984.49964

© Stindl This article is distributed under the terms of the [Creative Commons Attribution 4.0 International License](#), which permits unrestricted use, distribution, and redistribution in any medium, provided that the original author and source are

In North America, the passenger pigeon population was estimated at 3-5 billion individuals in the mid-1800s. A few decades later, the last individual died at the Cincinnati Zoo in 1914 during an unsuccessful breeding program (Hung et al. 2014). Hung and colleagues concluded from sequencing data that the passenger pigeon frequently experienced dramatic population fluctuations, possibly due to habitat change. According to their hypothesis, the passenger pigeon simply did not recover from the last population low, because of human hunting activities.

However, according to their genetic data (Hung et al. 2014), a population low equals 300,000 to 500,000 individuals living on the North American continent, which makes the hunting-to-extinction scenario rather unlikely. In the same article, the authors cite the mysterious and sudden extinction of the Rocky Mountain grasshopper, a widespread agricultural pest (Hung et al. 2014). Yet, there are no reports of massive grasshopper hunting activities in the late 1800s and there was no widespread use of insecticides until several decades later.

Since more than 99.9% of all species, which have ever lived on this planet, have become extinct (Raup 1991), the reported cases of the sudden disappearance of pigeons and grasshoppers might be the rule, not the exception. The mysterious observations could point to an intrinsic mechanism of species extinction, an old and forgotten European concept of organic evolution (Stindl 2014). There is a reason why intrinsic causes of extinction and saltatory speciation are not listed in contemporary biology textbooks anymore. It is the lack of any known mechanism, which can act over hundreds or thousands of species generations. Even Otto H. Schindewolf, one of the best-known supporters of the saltatory evolution model, could not provide any mechanistic explanation for his paleontological findings (Stindl 2014).

Barbara McClintock described broken chromosome ends (eroded telomeres) as leading to chromosomal instability and activated transposable elements (McClintock 1984) that are capable of rewiring the genome (Kunarso et al. 2010). Whereas it is a scientific fact that telomeres erode in somatic tissues during aging, they are thought to remain stable in the germ line of a species. However, the results of a large multigenerational telomere study on healthy subjects (Eisenberg, Hayes, and

credited.



Kuzawa 2012), especially regarding the loss of the positive grandfather effect on offspring telomere length in the maternal line, deliver indirect proof of telomere erosion in the human lineage (Stindl 2014). If telomeres erode in the female germ line, the long telomeres in old men`s testes would be no age effect, but a birth cohort effect. Accordingly, transgenerational telomere erosion is the ideal candidate for a biological clock that can accomplish sudden extinctions of widespread and successful species (Stindl 2014).

REFERENCES

- Eisenberg, D. T., M. G. Hayes, and C. W. Kuzawa. 2012. "Delayed paternal age of reproduction in humans is associated with longer telomeres across two generations of descendants." *Proc Natl Acad Sci U S A* no. 109 (26):10251-6. doi: 10.1073/pnas.1202092109.
- Hung, C. M., P. J. Shaner, R. M. Zink, W. C. Liu, T. C. Chu, W. S. Huang, and S. H. Li . 2014. "Drastic population fluctuations explain the rapid extinction of the passenger pigeon." *Proc Natl Acad Sci U S A* no. 111 (29):10636-41. doi: 10.1073/pnas.1401526111.
- Kunarsow, G., N. Y. Chia, J. Jeyakani, C. Hwang, X. Lu, Y. S. Chan, H. H. Ng, and G. Bo urque. 2010. "Transposable elements have rewired the core regulatory network of human embryonic stem cells." *Nat Genet* no. 42 (7):631-4. doi: 10.1038/ng.600.
- McClintock, B. 1984. "The significance of responses of the genome to challenge." *Science* no. 226 (4676):792-801. doi: 10.1126/science.15739260.
- Raup, D. M. 1991. *Extinction: bad genes or bad luck?* New York: W.W. Norton.
- Stindl, R. 2014. "The telomeric sync model of speciation: species-wide telomere erosion triggers cycles of transposon-mediated genomic rearrangements, which underlie the saltatory appearance of nonadaptive characters." *Naturwissenschaften* no. 101 (3):163-186. doi: 10.1007/s00114-014-1152-8.

APPENDIX

Submitted to PNAS 08/15/2014, rejected 08/20/2014.