

Hospital hyperglycemia: Status Investigation and Effect of a Real-time Glycemic Alert System

Chenxiang Cao¹, Victor Bernet², Zhaoxiang Liu¹, Caihong Li¹, Chongyang Bi¹, Zhenhua Sang¹, and Jianzhong Xiao¹

¹Tsinghua University

²Mayo Clinic Hospital Jacksonville

January 4, 2021

Abstract

BACKGROUND Hospital hyperglycemia is common and associated with potential adverse outcomes. A Hospital-wide Mobile Phone Alert (HMA) system was built to achieve real time glucose monitoring with warnings for glucose excursions. This study investigated the status of glucose control and evaluated the impact of HMA system on inpatient glycemia management. **METHODS** Inpatients with hyperglycemia hospitalized between 1 January, 2017 and 31 December, 2018 were identified excluding those < 18 years of age. The HMA system was activated on 1 October, 2017. It sent real time cellphone warning messages to the patient's designated team physician whenever glucose levels > 10 mmol/L or < 3 mmol/L were detected. A serum glucose > 7.8 mmol/L was defined as hospital hyperglycemia (HH), and > 10 mmol/L was defined as significant HH (SHH). Glucose excursions before and after the HMA system was instituted were compared. **RESULTS** The incidence of HH, SHH and hypoglycemia was 26.1%, 12.8% and 2.5%, respectively. With the HMA system, the monthly glucose related consultation rate for all inpatients increased 65.9%. The rate of HH glucose amount/ total glucose amount improved with the HMA system, being lower than pre HMA system activation for the surgical wards ($15.8 \pm 4.7\%$ vs $21.1 \pm 6.1\%$, $p < 0.05$). **CONCLUSIONS** In this study, one third of inpatients were noted to experience hyperglycemia. Real time cellphone warning messages to the patient's designated team physician can improve consultation utilization for blood glucose excursions. The alert system was found to reduce the incidence of hyperglycemia on surgical wards.

INTRODUCTION

The prevalence of hyperglycemia in hospitalized patients is common and in our study varied from 12% to 50% of all inpatients at our facility^{1,2 1-3}. Episodes of hyperglycemia were associated with several reasons, to include: history of diabetes mellitus (DM), newly diagnosed diabetes mellitus and stress-related hyperglycemia^{4,5}. Inpatient hyperglycemia has been associated with various adverse outcomes to include a higher mortality, longer hospital stay, increased infections, and complications^{5,6}. In clinic practice, the achievement of blood glucose goal in inpatients with hyperglycemia is not satisfied and a number of barriers can prevent the achievement^{7,8}. The physician's omissions of hyperglycemia and impaired medication timing due to hospital system are important reasons that impact the management of hospitalized patients^{7,8}. While there have been numerous publications on how to improve the management of glucose excursions in hospitalized patients, available systems to date appear to have fallen short of achieving optimal results⁹⁻¹³. Although diabetes management guidelines for inpatient diabetes advise the development and use of inpatient blood glucose monitoring systems, no gold standard models or systems presently exist.

We established a hospital-wide inpatient glucose management model for glucose monitoring and response to clinically relevant glucose excursions including: a glycemic care team (GCT), standard operation procedures (SOP), and glucose care information system (GCI). The GCI was integrated with our electronic medical record (EMR), and was developed to provide an alarm message whenever noting hyperglycemia (>7.8 mmol/L)

or hypoglycemia (< 3 mmol/L) in a patient. This system was named the Hospital-wide Mobile Phone Alert (HMA) with it being developed to achieve real time warnings and resultant timely response for glucose excursions with the intent of improving overall glucose control. The study aimed to evaluate the impact of the system HMA on inpatient hyperglycemia control.

RESEARCH DESIGN AND METHODS

Patients

We searched our facility's EMR database in order to identify all adult patients hospitalized in our facility between January 1, 2017 through December 31, 2018 and found that 47,093 inpatients (99.5% of total adult inpatients) who had been admitted were tested for blood glucose levels. The study did not include transplant patients. There were 12,292 inpatients (26.1%) who experienced hyperglycemia > 7.8 mmol/L and 1,175 inpatients (2.5%) that experienced hypoglycemia < 3 mmol/L. Their demographic data, diagnosis, treatment, and laboratory test were analyzed.

Study design

This study was reviewed and approved by the Ethics Committee of Beijing Tsinghua Changgung Hospital and Beijing Tsinghua Changgung Hospital Fund (Grant No. 12017C1002). The HMA system was activated on October 1, 2017. The HMA system was set to send real time warning messages to the patient's immediate team physician mobile phone when hypoglycemia (< 3 mmol/L) or hyperglycemia (> 10 mmol/L) was detected. HMA alert messages were set not to repeat until 72 hours had passed. The alert would be activated by results from either point of care POC or serum glucose testing. Hospital hyperglycemia (HH) was defined as random blood glucose > 7.8 mmol/L, whereas a blood glucose > 10 mmol/L was defined as significant HH (SHH). The HH and SHH rates were defined as HH amount/ total tested glucose amount % and SHH amount/ total tested glucose amount %, respectively, and used as indicators for glycemic control. The patients hospitalized before the HMA system was activated acted as the baseline group and with those patients monitored with the HMA system being labelled as the HMA group.

Statistical analysis

Continuous variables were analyzed using the Student's t-test and reported as the mean-standard deviation. Categorical variables were analyzed using the chi-squared test and Fisher's exact test and reported as frequency, percentages and standard errors. Statistical analyses were conducted using SPSS 22. All enrolled patients were assigned a unique identifier in the form of a participant number as to ensure anonymity.

RESULTS

The status of diabetes and hospital hyperglycemia

The rate of pre-existing diabetes history in adult inpatients was 17.7%. Excluding the inpatient Endocrinology ward, Cardiac Surgery (42.9%), Nephrology (36.7%) and Neurology (35.5%) were noted to have the highest rates of DM history, while the Plastic Surgery (4.5%), Gynecology (4.4%), and Intensive Care Unit (ICU) (2.6%) wards had the lowest rates. The incidence of HH and SHH based on serum glucose testing was 26.1% and 12.8%, overall. In non-intensive care unit inpatients, the rates of HH and SHH were 20.7% and 10.8%, respectively. Excluding the Endocrinology inpatient ward, the ICU had the highest incidence of hyperglycemia (HH-73.6% and of SHH- 30.6%) being followed by Cardiac surgery (HH- 48.6% and SHH- 34.3%) and Cardiology + CCU (HH- 36.5% and SHH- 22.5%). On the other hand, Plastic surgery (HH and SHH- 0%), Obstetrics (HH- 1.9% and SHH - 0.7%), and Gynecology (HH - 2.9% and SHH - 1.3%) had the lowest incidences (Table 1).

Before HMA activation, Neurology required the largest amount of glycemic related consultations (16.4/month) followed by Cardiology + CCU (12.7/month) and Orthopedics (5.8/month). Thoracic Surgery, Gastrointestinal Surgery, Cardiac Surgery, and General Practice had requested no more than one consultation per month (Table 1).

Effect of HMA on inpatients management

The total amount of HMA messages for adult inpatients was 9539 during the 15 month test period. Within the Internal Medicine wards (excluding Endocrinology) the HMA rate was: Neurology (93/month), Cardiology + CCU (55/month), and Nephrology (52/month), these being the top three departments in receiving HMA warning messages while Hepatobiliary Surgery (85/month), Vascular Surgery (37/month), and Orthopedics (36/month) were noted as the top three surgical wards receiving HMA alerts (Fig. 1).

The glycemia related consultation (number per month) of all adult inpatients increased by 65.9% in the period of HMA monitoring. The consult rates for the Gastrointestinal Surgery (476%) and Gastroenterology (446.7%) wards increased by more than fourfold, while the rates for Cardiac Surgery (200%) and Hepatobiliary Medicine (218%) were doubled. In contradistinction, a decrease in glycemia related consultations were noted for the Cardiology + CCU (20%) and Neurosurgery (16.4%) wards (Fig. 2).

The glycemia excursions between baseline and HMA group

There were no significant differences noted between the two groups in occurrence of either HH or SHH rate amongst all inpatients or patients admitted to internal medicine wards, while the HH rate for surgical wards decreased significantly (HMA group vs baseline = $15.8 \pm 4.7\%$ vs $21.1 \pm 6.1\%$, $p < 0.05$) (Table 2).

DISCUSSION

Hospital hyperglycemia which is defined as random blood glucose > 7.8 mmol/L with or without diabetes history occurs frequently in inpatients¹⁴⁻¹⁶. Inpatient hyperglycemia has been associated with an increase in mortality, complications, hospital stay length, and overall hospital cost^{17,18}. Previous reports have indicated that the inpatient hyperglycemia incidence is around 32.2% of ICU patients and 32% of patients admitted to non-ICU wards in the U.S.¹⁹. The reported incidence of hyperglycemia was 45.7% in China²⁰. In our study, 26.1% of total adult inpatients were noted to experience hyperglycemia (> 7.8 mmol/L) with 12.8% having a significant episode of hyperglycemia (> 10 mmol/L). The HH and SHH prevalence of ICU inpatients are as high as 73.6% and 30.6%, respectively. Meanwhile, there are 20.7% of non-ICU inpatients with HH and 10.8% with SHH. The inpatient wards with higher hyperglycemia rates can be divided into two categories: the first category is those who tend to admit more patients with pre-existing diabetes history who are at risk for developing excursions in blood glucose control, such as the Cardiology ward; the second category consists of patients without a history of diabetes but who experience illness, stress, and/or medication related hyperglycemia. Surgical and ICU wards typically represent patients falling into the latter category^{21,22}. As is known, medications other than glucocorticoids can induce hyperglycemia, such as lipid-modifying agents, antidepressants and estrogen²³⁻²⁶. In our hospital, patients following craniocerebral operations who are admitted to the Neurosurgery ward regularly receive dexamethasone therapy. However, episodes of hyperglycemia were low which appeared to be related to intensive glucose monitoring whenever glucocorticoid therapy was prescribed. This finding seems to support the belief that intensive monitoring of blood glucose is of benefit for inpatient glycemic control^{9,27,28}. Illness associated stress which occurs in 12 -22% of inpatients has been reported to induce hyperglycemia which a resultant increase in both morbidity and mortality^{5,22,29-31}. Stress-induced hyperglycemia is felt to be related to over activation of the hypothalamic-pituitary-adrenal axis and/or the sympathetic autonomic nervous system thereby causing insulin resistance followed by hyperglycemia³². Sleep loss which is common in inpatients is considered as an independent risk factor which can disrupt glycemic homeostasis^{33,34}. Our study also indicates that thoracic surgery is unique in character as the rate of mild hyperglycemia (7.8-10mmol/L) is nearly three times higher than the rate of pre-existing diabetes history. This association may be related to the fact that many thoracic surgery cases were for a malignant tumor, and both lung cancer itself and its treatment have been associated with the development of hyperglycemia³⁵⁻³⁷. Traditional inpatient glycemic care is typically passive in nature and not very efficient^{14,20,38}. Lack of timely reaction to glucose excursions, delayed treatment, and undiagnosed new onset diabetes or pre-diabetes have all been identified as common factors related to poor inpatient glycemic control³⁹. Our facility developed an alert system for glucose excursions which sent real time cellphone message to the hospitalized patient's designated inpatient team physician with the aim

of improving the reaction time and intervention for episodes of hypo- or hyperglycemia. Previously studied glucose alert systems have shown improvement of inpatient glycemic control^{40,41}. With the HMA system, the monthly glycemia related consultation rate for all inpatients increased nearly 66%. While the HMA system was not noted to reduce the rate of total inpatient hyperglycemic episodes, the HH rate for the surgical wards was decreased. These findings indicate that the effect of HMA system on utilization of formal consultations for blood glucose excursions varied by the specialty service of the ward. The surgical wards actively utilized the alert by requesting endocrinology team consultation. It should be noted that two kinds of endocrinologist consultation are performed in our hospital: 1) individual endocrinology consults and 2) glycemic care team (GCT) assessments. The former is the traditional way to deal glucose excursions in Chinese hospitals, and is performed during a single visit by an endocrine physician to the inpatient with recommendations being provided to fulfill the consultation requirement. The GCT is composed of a nurse, a diabetes educator, nutritionist and an endocrinologist. GCT assessment is led by the endocrinologist who takes the initiative to follow-up the patient during the period that glucose excursions are occurring. In order to cover the entire complement of hospitalized patients, the GCT would require a large number of endocrinologists which presently is beyond the capacity of our facility. Therefore, at present, GCT assessments are only available to a select number of inpatients in our hospital. The GCT glucose assessments were noted to correlate with a reduced rate of hyperglycemia and a shorter length of hospital stay, while individual endocrinologist consults were noted to decrease the average length of hospital stay and medical care expenses for inpatients experiencing hyperglycemia⁴². However, the HMA system was found to reduce the HH rate for the surgical wards even when just individual endocrinologist consultations were utilized. Within internal medicine wards, a quite different response to HMA alerts was noted. Nephrology was the fourth highest ward receiving HMA alerts but the third lowest ward in increased utilization of endocrinologist consults. The glycemia related consultation rate for Cardiology + CCU ward even declined by 20%. There is rare previous report describing the relationship between inpatient ward response to glycemic alert, hyperglycemia episodes and glycemia related consultations. We postulate that the less active response to the glucose excursions alerts may lead to suboptimal glycemic control secondary to a delay in endocrinologist consultation, or possibly that individual endocrinologist consults were not felt to be effective enough for internal medicine wards patients with more complicated glucose control issues. Future investigation is warranted as to determine the exact causes for variations in utilization of consultations for HMA alerts.

In conclusion, inpatient hyperglycemia which is associated with inferior outcomes and longer hospital stays is commonly encountered in hospitalized adults, and real time glycemic alert systems for inpatients hold the potential to improve glycemic control thereby reducing complications, morbidity and mortality related to excursions in glucose from either hyper- or hypoglycemia. This study revealed the prevalence of hospital hyperglycemia at 26.1% overall, 20.7% in non-ICU inpatients, and 73.6% in ICU inpatients. It is noted that real time glycemic alert cellphone messages to the team physician who is directly responsible for the inpatient with the glucose excursions can increase the glycemia related consultation rate overall and also appears to decrease hyperglycemia episodes in patients on the surgical wards.

References

1. Rodriguez-Calero MA, Barceló Llodrá E, Cruces Cuberos M, Blanco-Mavillard I, Pérez Axartell MA. Effectiveness of an evidence-based protocol for the control of stress-induced hyperglycaemia in critical care. *Enfermería Intensiva (English ed)*.2019;30(1):4-12.
2. Boord JB, Greevy RA, Braithwaite SS, et al. Evaluation of hospital glycemic control at US academic medical centers. *J Hosp Med*.2009;4(1):35-44.
3. Cook CB, Elias B, Kongable GL, Potter DJ, Shepherd KM, McMahon D. Diabetes and hyperglycemia quality improvement efforts in hospitals in the United States: current status, practice variation, and barriers to implementation. *Endocr Pract*. 2010;16(2):219-230.
4. Clement S, Braithwaite SS, Magee MF, et al. Management of diabetes and hyperglycemia in hospitals. *Diabetes Care*.2004;27(2):553-591.

5. Umpierrez GE, Pasquel FJ. Management of Inpatient Hyperglycemia and Diabetes in Older Adults. *Diabetes Care*. 2017;40(4):509-517.
6. CHARLES KODNER MLA, MD; and KATHERINE POHLGEERS, MD. Glucose Management in Hospitalized Patients. *Am Fam Physician* 2017;96(10):648-654 Copyright © 2017 American Academy of Family Physicians. 2017.
7. Kodner C, Anderson L, Pohlgeers K. Glucose Management in Hospitalized Patients. *Am Fam Physician*. 2017;96(10):648-654.
8. Pichardo-Lowden A, Farbaniec M, Haidet P. Overcoming barriers to diabetes care in the hospital: The power of qualitative observations to promote positive change. *J Eval Clin Pract*. 2019;25(3):448-455.
9. Thabit H, Hovorka R. Bridging technology and clinical practice: innovating inpatient hyperglycaemia management in non-critical care settings. *Diabet Med*. 2018;35(4):460-471.
10. Sheen YJ, Huang CC, Huang SC, et al. Implementation of an Electronic Dashboard with a Remote Management System to Improve Glycemic Management among Hospitalized Adults. *Endocr Pract*. 2019.
11. Momesso DP, Costa Filho RC, Costa JLF, et al. Impact of an inpatient multidisciplinary glucose control management program. *Arch Endocrinol Metab*. 2018;62(5):514-522.
12. Mathioudakis N, Bashura H, Boyer L, et al. Development, Implementation, and Evaluation of a Physician-Targeted Inpatient Glycemic Management Curriculum. *J Med Educ Curric Dev*. 2019;6:2382120519861342.
13. Aloji JA, Mulla C, Ullal J, Lieb DC. Improvement in inpatient glycemic care: pathways to quality. *Curr Diab Rep*. 2015;15(4):18.
14. O'Sullivan EP, Duignan J, O'Shea P, Griffin D, Dinneen SF. Evaluating hyperglycaemia in the hospitalised patient: towards an improved system for classification and treatment. *Ir J Med Sci*. 2014;183(1):65-69.
15. Bar-Dayyan Y, Beer I, Boaz M, Landau Z, Jakubowicz D, Wainstein J. Hyperglycaemia in patients with no history of diabetes presenting to Internal Medicine Departments. *International journal of clinical practice*. 2015;69(11):1289-1295.
16. Feldman-Billard S, Joubert M, Morello R, et al. High prevalence of diabetes mellitus and hospital-related hyperglycaemia in French general wards. *Diabetes Metab*. 2013;39(5):454-458.
17. Leite SA, Locatelli SB, Niece SP, Oliveira AR, Tockus D, Tosin T. Impact of hyperglycemia on morbidity and mortality, length of hospitalization and rates of re-hospitalization in a general hospital setting in Brazil. *Diabetol Metab Syndr*. 2010;2:49.
18. Cakir M, Altunbas H, Karayalcin U. Hyperglycemia: an independent marker of in-hospital mortality in patients with undiagnosed diabetes. *J Clin Endocrinol Metab*. 2003;88(3):1402; author reply 1402.
19. Swanson CM, Potter DJ, Kongable GL, Cook CB. Update on inpatient glycemic control in hospitals in the United States. *Endocr Pract*. 2011;17(6):853-861.
20. Hao S, Zhang N, Fish AF, et al. Inpatient glycemic management in internal medicine: an observational multicenter study in Nanjing, China. *Curr Med Res Opin*. 2017;33(8):1371-1377.
21. Kwon S, Thompson R, Dellinger P, Yanez D, Farrohki E, Flum D. Importance of perioperative glycemic control in general surgery: a report from the Surgical Care and Outcomes Assessment Program. *Ann Surg*. 2013;257(1):8-14.
22. Cheisson G, Jacqueminet S, Cosson E, et al. Perioperative management of adult diabetic patients. Review of hyperglycaemia: definitions and pathophysiology. *Anaesth Crit Care Pain Med*. 2018;37 Suppl 1:S5-S8.
23. Paredes S, Alves M. [Management and Treatment of Glucocorticoid-Induced Hyperglycemia]. *Acta Med Port*. 2016;29(9):556-563.

24. Tamez-Perez HE, Quintanilla-Flores DL, Rodriguez-Gutierrez R, Gonzalez-Gonzalez JG, Tamez-Pena AL. Steroid hyperglycemia: Prevalence, early detection and therapeutic recommendations: A narrative review. *World J Diabetes*. 2015;6(8):1073-1081.
25. Fathallah N, Slim R, Larif S, Hmouda H, Ben Salem C. Drug-Induced Hyperglycaemia and Diabetes. *Drug Saf*. 2015;38(12):1153-1168.
26. Liu MZ, He HY, Luo JQ, et al. Drug-induced hyperglycaemia and diabetes: pharmacogenomics perspectives. *Arch Pharm Res*. 2018;41(7):725-736.
27. Krinsley JS, Chase JG, Gunst J, et al. Continuous glucose monitoring in the ICU: clinical considerations and consensus. *Critical Care*. 2017;21(1).
28. Krinsley JS, Bruns DE, Boyd JC. The impact of measurement frequency on the domains of glycemic control in the critically ill—a Monte Carlo simulation. *J Diabetes Sci Technol*. 2015;9(2):237-245.
29. Davis G, Fayfman M, Reyes-Umpierrez D, et al. Stress hyperglycemia in general surgery: Why should we care? *J Diabetes Complications*. 2018;32(3):305-309.
30. Russo MP, Grande Ratti MF, Giunta DH, Elizondo CM. Hospitalized patients with stress hyperglycemia: Incidence of diabetes and mortality on follow-up. *Endocrinología, Diabetes y Nutrición (English ed)*. 2018;65(10):571-576.
31. Russo MP, Elizondo CM, Giunta DH, Grande Ratti MF. Prevalence of hyperglycemia and incidence of stress hyperglycemia in hospitalized patients: A retrospective cohort. *Eur J Intern Med*. 2017;43:e15-e17.
32. Mifsud S, Schembri EL, Gruppetta M. Stress-induced hyperglycaemia. *Br J Hosp Med (Lond)*. 2018;79(11):634-639.
33. Stewart NH, Arora VM. Sleep in Hospitalized Older Adults. *Sleep Med Clin*. 2018;13(1):127-135.
34. DePietro RH, Knutson KL, Spampinato L, et al. Association Between Inpatient Sleep Loss and Hyperglycemia of Hospitalization. *Diabetes Care*. 2017;40(2):188-193.
35. Kirakli EK, Yilmaz U, Yilmaz H, Komurcuoglu B. Fasting Blood Glucose Level in Locally Advanced Non-Small Cell Lung Cancer: a New Prognostic Factor? *Horm Cancer*. 2018;9(3):188-196.
36. Sakuma I, Nagano H, Yoshino I, Yokote K, Tanaka T. Ceritinib Aggravates Glycemic Control in Insulin-treated Patients with Diabetes and Metastatic ALK-positive Lung Cancer. *Intern Med*. 2019;58(6):817-820.
37. Luo J, Chen YJ, Chang LJ. Fasting blood glucose level and prognosis in non-small cell lung cancer (NSCLC) patients. *Lung Cancer*. 2012;76(2):242-247.
38. Drincic AT, Akkireddy P, Knezevich JT. Common Models Used for Inpatient Diabetes Management. *Current diabetes reports*. 2018;18(3):10.
39. El Khoury G, Mansour H, Kabbara WK, Chamoun N, Atallah N, Salameh P. Prevalence, Correlates and Management of Hyperglycemia in Diabetic Non-critically Ill Patients at a Tertiary Care Center in Lebanon. *Curr Diabetes Rev*. 2019;15(2):133-140.
40. Seheult JN, Pazderska A, Gaffney P, et al. Addressing Inpatient Glycaemic Control with an Inpatient Glucometry Alert System. *Int J Endocrinol*. 2015;2015:807310.
41. Kyi M, Wraight PR, Rowan LM, Marley KA, Colman PG, Furlanos S. Glucose alert system improves health professional responses to adverse glycaemia and reduces the number of hyperglycaemic episodes in non-critical care inpatients. *Diabet Med*. 2018;35(6):816-823.
42. Yu DJ, Ebaid A. To Consult or Not to Consult: The Role of the Endocrinologist in the Management of Diabetes Mellitus in the Hospital Setting. *Curr Emerg Hosp Me R*. 2015;3(1):38-42.

Table page 1:

Table 1 - The baseline of pre-existing diabetes history, hyperglycemia and consultation in different inpatient wards before HMA activation (1/2017 - 9/2017)

Department	Total Inpatients (N)	Pre-existing diabetes inpatients (N)	HH inpatients (N)	SHH inpatients (N)
ICU	1134	29	835	3
Cardiac Surgery	35	15	17	0
Cardiology+CCU	1069	332	497	2
Nephrology	447	164	167	9
Hepatobiliary Surgery	821	129	304	3
Hepatobiliary Medicine	108	28	35	3
General Medicine	356	114	115	5
Respiration	261	62	79	3
Gastrointestinal Surgery	688	48	206	8
Urology	1300	149	336	3
Neurology	828	294	196	3
Thoracic Surgery	140	10	28	3
Vascular Surgery	469	113	93	4
Hematology and Oncology	398	69	78	4
Orthopedics	1217	189	238	3
Hepatobiliary Intervention	92	9	16	8
General Surgery	562	80	94	4
Neurosurgery	722	59	119	4
Radiotherapy	21	6	3	3
Rehabilitation Medicine	415	113	37	3
Gastroenterology	349	51	29	3
Ophthalmology	414	80	23	3
O-HNS	1016	85	43	3
Gynecology	1142	50	33	3
Obstetrics	858	134	16	6
Plastic Surgery	22	1	0	0
Total adult inpatients	15350	2760	4013	2

HH, hospital hyperglycemia (serum random glucose > 7.8 mmol / L); SHH, significant hospital hyperglycemia (blood glucose > 10 mmol / L); CCU, cardiac intensive care unit; ICU, intensive care unit; O-HNS, Otolaryngology-Head and Neck surgery.

Table page 2:

Table 2 - The HH and SHH rates of adult inpatients between baseline and HMA group.

	Baseline	HMA group	p
HH rate (%)	21.9 ± 0.8	22.6 ± 1.6	0.222
Internal Medicine wards	23.8±9.6	26.7±7.0	0.456
Surgical wards	21.1± 6.1	15.8 ± 4.7	<0.05
SHH rate (%)	9.7 ± 0.5	10 ± 1	0.279
Internal Medicine wards	12.6±5.1	14.5±5.2	0.417
Surgical wards	9.3±6.6	9.7±4.8	0.878

HH, hospital hyperglycemia; SHH, significant hospital hyperglycemia; HMA, Hospital-Wide Mobile Phone

Alert System.

Figure 1 - **The average monthly HMA message amount for different inpatient wards (10/2017 - 12/2018)** . HMA- Hospital-wide Mobile Phone Alert System; Otolaryngology-Head and Neck surgery; CCU- cardiac intensive care unit; ICU- intensive care unit; Holistic hospital - all adult inpatients.

Figure 2 - **Percentage of average monthly consultation amount change between baseline and HMA group.** HMA- Hospital-Wide Mobile Phone Alert System; O-HNS, Otolaryngology-Head and Neck surgery; CCU- cardiac intensive care unit; ICU- intensive care unit; Holistic hospital- all adult inpatients.

Hosted file

Figure page 1_V 12.27.2020.pdf available at <https://authorea.com/users/325309/articles/502134-hospital-hyperglycemia-status-investigation-and-effect-of-a-real-time-glycemic-alert-system>

Hosted file

Figure page 2_V 12.27.2020.pdf available at <https://authorea.com/users/325309/articles/502134-hospital-hyperglycemia-status-investigation-and-effect-of-a-real-time-glycemic-alert-system>