

# Extensive keloid scarring in a patient with pemphigus vulgaris

Dorsaf Elinkichari<sup>1</sup>, Anissa Zaouak<sup>1</sup>, Amal Chamli<sup>2</sup>, Houda Hammami Ghorbel<sup>2</sup>, and Samy Fenniche<sup>1</sup>

<sup>1</sup>Habib Thameur Hospital

<sup>2</sup>University of Tunis El Manar Faculty of Medicine of Tunis

August 15, 2022

## Abstract

Since pemphigus blisters are intraepidermal, scarring should induce at most a post-inflammatory hyperpigmentation. We describe a very atypical and unusual course of pemphigus vulgaris with extensive keloid formation despite high systemic steroids. This could be promoted by the severe flare of the disease, the delay of scarring and the superinfection.

## Extensive keloid scarring in a patient with pemphigus vulgaris

**Dorsaf Elinkichari<sup>1,2</sup>, Anissa Zaouak<sup>1,2</sup>, Amal Chamli<sup>1,2</sup>, Houda Hammami<sup>1,2</sup> Samy Fenniche<sup>1,2</sup>**

<sup>1</sup>Department of Dermatology, Habib Thameur Hospital, Tunis, Tunisia

<sup>2</sup>Faculty of Medicine of Tunis, University of Tunis El Manar, Tunis, Tunisia

## Correspondence

Dorsaf Elinkichari, Department of Dermatology, Habib Thameur Hospital, Tunis, Tunisia.

Email: [dorsaf.elinkichari@gmail.com](mailto:dorsaf.elinkichari@gmail.com)

## Key Clinical message

Keloid scarring in PV may be promoted by the severity of the flare, the delay of scarring and the superinfection. Treatment in such cases may be challenging.

## Abstract

Since pemphigus blisters are intraepidermal, scarring should induce at most a post-inflammatory hyperpigmentation. We describe a very atypical and unusual course of pemphigus vulgaris with extensive keloid formation despite high systemic steroids. This could be promoted by the severe flare of the disease, the delay of scarring and the superinfection.

## Key words

Clinical Dermatology, Pemphigus vulgaris, Keloid

The patient has given written informed consent to the publication of his case details.

## Introduction

Keloid scarring in pemphigus vulgaris (PV) is uncommon. Since pemphigus blisters are intraepidermal, scarring should induce at most a post-inflammatory hyperpigmentation. Only four cases of keloid scars

following pemphigus have been reported [1–4]. Herein, we describe an unusual extensive keloid scarring following a severe flare of PV.

## Observation

A 47-year-old Caucasian woman with a 4-month history of PV presented with extensive keloid scarring at areas of prior PV erosions. Four months ago, she had been hospitalized for a severe PV flare with erosions affecting her trunk, upper and lower limbs (figure 1a). Oral, nasal and conjunctival mucosa were also involved. In addition, nail damage was noted with onycholysis, onychomadesis and necrotic paronychia (figure 1b). The Pemphigus Disease Area Index (PDAI) was 55. The diagnosis of PV was confirmed by histological and immunological investigation showing intraepidermal blistering, supra-basal epidermal acantholysis and inter-keratinocyte IgG deposits in direct immunofluorescence. ELISA was positive for anti-desmoglein 3 (120 U/ml) and anti-desmoglein 1 (180 U/ml). The erosions did not heal after 16 days of treatment with 1.5mg/kg/day of prednisone-equivalent and became extensive and hemorrhagic. The patient was afebrile. The complete blood count revealed hyperleukocytosis and hyperneutrophilia. CRP level was 386 mg/l. Two days later, a greenish malodorous discharge covered almost all the erosions (Figure 2a). Cultures grew *Pseudomonas aeruginosa*. Thus, the diagnosis of ecthyma gangrenosum complicating a PV was retained. A dramatic improvement of the patient’s skin erosions was observed two weeks after introducing an antibiogram-adjusted antibiotic therapy based on Tazobactam and Amikacin. After two weeks, the erosions had re-epithelialized with post-inflammatory hyperpigmentation, and the patient was discharged. We started oral steroids tapering without relapse. At the four-month follow-up, extensive keloids were noted on previous superinfected PV sites (figure 2b), for which intralesional corticosteroids were scheduled.

## Discussion

Three main factors of keloids are known: infection, deep injury and mechanical trauma [5]. So far, only four cases of keloid scars following pemphigus have been reported [1–4]. Gupta et al. reported a case of uninfected PV erosions that healed with keloids despite high dose of systemic steroids [2]. Shivaswamy et al. described keloids in a patient whose pemphigus was treated, *inter alia*, with cyclosporin. The onset of keloids in this patient was attributed to cyclosporin [3]. In the latest two cases, such as in our patient, extensive keloids appeared in patients whose pemphigus had been superinfected. Authors suggested that secondary infection induced keloids via dermal damage. However, the paucity of cases of keloids following PV flares contrasts with the frequency of superinfections which are common in pemphigus, suggesting the presence of other unknown factors [1,4].

Our case is notable due to the severity of the PV flare which was refractory to conventional treatment and then complicated by ecthyma gangrenosum. Re-epithelialization took more than one month and ended with extensive keloids after three months. It is reasonable that superinfection led to deep intradermal inflammation and keloid formation. In fact, keloids occurred in previously superinfected PV areas. In this context, there are reports of keloid onset following herpetic infections, with the keloids involving previously infected areas [6]. An alternative hypothesis is that erosions in our patient took exceptionally long to heal, which maintained a long inflammation and scarring process with a chronic fibroblast activation. This does not exclude the potential role of apoptotic cells as well as increased levels of tumor necrosis factor-alpha, found in PV, that may contribute to the dysregulation of fibroblasts, and lead to keloids in susceptible patients [7]. Intralesional steroids are generally proposed in patients with keloids. Yet, our patient is under high systemic doses of steroids, making this alternative debatable as we can opt for compression clothing and active monitoring.

## Conclusion

This case is being reported to highlight a very atypical and unusual course of PV with extensive keloid formation despite high systemic steroids. This could be promoted by the severe form of PV, the delay of scarring and the superinfection. Moreover, treatment in such cases may be challenging since patients are usually under systemic steroids making intralesional steroids contestable.

## CONFLICT OF INTEREST

None. This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

## AUTHOR CONTRIBUTIONS

Dorsaf Elinkichari is the guarantor of the content of the manuscript, including the data and analysis. Anissa Zaouak contributed to acquisition of data, conception, and interpretation of information and gave final approval of the version to be submitted. Amal Chamli revised data critically for important intellectual content. Houda Hammami and Samy Fenniche contributed to interpretation of data and revision of the manuscript.

## ETHICAL STATEMENT

Written informed consent was obtained from the patient.

## References

- [1] Sako EY, Workwick S. Extensive keloid formation after pemphigus vulgaris. *Dermatol Online J* 2015;21:13030/qt7sc4z88j.
- [2] Gupta SD, Khanna N, Rathi S. Extensive keloidal healing of pemphigus vulgaris. *Indian J Dermatol Venereol Leprol* 1997;63:199–200.
- [3] Shivaswamy K, Thappa D. Extensive keloids following cyclosporin therapy in a pemphigus vulgaris patient. *J Eur Acad Dermatol Venereol* 2007;21:111–112.
- [4] Nguyen J, Amber KT, Chapman LW. Pemphigus vulgaris complicated by extensive keloid scarring. *Clin Exp Dermatol* 2018;43:76–77.
- [5] Gira AK, Brown LF, Washington CV, Cohen C, Arbiser JL. Keloids demonstrate high-level epidermal expression of vascular endothelial growth factor. *J Am Acad Dermatol*. 2004;50(6):850-3.
- [6] Koley S, Saoji V, Salodkar A. Unusual formation of keloids after each episode of recurrent herpes zoster in an HIV positive patient. *Indian J Sex Transm Dis AIDS* 2009;30:109–111.
- [7] Chiapa-Labastida M, Zentella-Dehesa A, León-Dorantes G. Pemphigus vulgaris: accumulation of apoptotic cells in dermis and epidermis possibly relates to pathophysiology through TNF-alpha production by phagocytes. *Eur J Dermatol* 2011;21:874–888.







