

different treatment modalities for pulp stones

brigui foudad¹, amira kikly¹, Wided Askri¹, Ameni belghith¹, Neila Zokkar¹, and Nabiha Douki¹

¹Université de Monastir Faculté de Médecine Dentaire de Monastir

July 17, 2023

Type of article (Case Report)

different treatment modalities for pulp stones

Fouad Brigui¹, Amira Kikly^{1,3}, Wided Askri², Ameni Chadlia Belghith^{1,3}, Neila Zokkar¹ and Nabiha Douki Zbidi^{1,3}

¹ *Dental Medicine Department of Sahloul Sousse Hospital, Faculty of Dentistry of Monastir, Tunisia*

² *Dental Medicine Department of Hached Sousse Hospital, Faculty of Dentistry of Monastir, Tunisia*

³ *Research Laboratory LR12ES11*

Abstract

Pulpal calcifications or also called pulpal calcium degeneration are small masses of calcified tissue that obliterates part or almost all of the endodontic network, the presence of these masses can further complicate endodontic treatment and requires an adaptation of the therapeutic approach. The objective of this article is to illustrate the difficulties and the keys to success of endodontic treatment in the presence of pulpal calcifications. Three patients with three levels of difficulty; a free coronary pulp stone, an adherent coronary pulp stone and diffuse root canal calcification required well-adapted treatment in order to eradicate these calcified masses. The discovery of pulpal calcifications during daily practice is particularly frequent. These calcifications can present different clinical forms and pose problems of diagnosis and management. The indication for endodontic treatment cannot be based solely on the presence of pulp stone but on a diagnosis confirmed by a clinical and radiological examination showing the pulp status of the tooth.

Keywords

Pulp-Stone, obliteration, dental microscope, catheterization

Introduction


Pulpal calcifications or also called pulpal calcium degenerations are small masses of calcified tissue that obliterates part or almost all of the endodontic network [1]. the exact cause of their formation remains unknown. Therefore, it is currently unclear whether their presence reflects a pathological manifestation or a normal physiological development of the dental pulp [2]. Thus, many factors predisposing to the development of pulpal calcification have been suggested, such as aging, caries, periodontal disease and orthodontic movements. On the other hand, some studies have tried to establish a relationship with other general diseases such as cardiovascular and renal diseases [3]. These masses of calcified tissue can be found at the level of the pulp chamber and/or at the level of the root canals, but the accessibility to the canal orifices which is an essential element for the success of any endodontic treatment can become more complicated by the presence of these pulpolites [4].

The objective of this article is to illustrate, through three clinical cases, the difficulties and the keys to the success of endodontic treatment in the presence of pulpal calcifications.

Materials and Methods


Clinical case N°1

A 22-year-old patient in good general condition consulted following spontaneous pain related to the first left mandibular molar (36). The clinical examination reveals a Sista 1.3 carious lesion. The radiological examination shows the presence of calcification which obliterates almost all of the pulp chamber with a cleavage plane (clear X-ray line) between the calcified tissue and the healthy dental tissue (Fig.1).









After carious eviction and opening of the access cavity by connecting the projection of the pulp horns (Fig.2), the extent of the pulpolite is delimited by a hemorrhagic colorimetric demarcation between the dentin of the cameral walls and the calcified tissue center (Fig.3). Subsequently, this demarcation was widened using an ultrasonic diamond insert (ET18D) (Fig.4).






Once released and mobilized, the calcified mass is completely removed by an endodontic excavator (Fig.5.6.7).



After removal of the pulpstone, abundant irrigation with Naocl 5.25% was performed to better visualize the root canal entries (Fig.8), and from here endodontic treatment was continued as if of a normal tooth without particularities (Fig.9).





Clinical case N°2 :

A 66-year-old patient, hypertensive, diabetic under treatment, consulted urgently for symptomatic irreversible pulpitis related to the first maxillary right molar (16). The clinical examination shows a deep mesial caries reaching the inner third of the dentin, the radiological examination refines the report and the extent of the carious lesion and highlights the presence of a pulp polyp which completely obliterates the cameral pulp (pulp polyp adhering to the dentinal walls and to the pulpal floor). (Fig.10)



The reference points cited in the first clinical case when creating the access cavity are not applicable in this case. For example, there is no longer the feeling of emptiness during pulp effraction, there is no cleavage plane and the pulpolite cannot be mobilized.

In this case, we proceeded by deepening the perimeter of the access cavity step by step with a long-necked ball burr, then began circumferential pulpolite clearance with ultrasonic diamond inserts (ET18D / ETBD), giving priority to freeing the canal entrances (Fig. 11.12.13).


Using the same long-necked ball burr, the calcified tissue was abraded under direct visual control, using the color difference (between cameral dentin and calcified tissue) as a milling guide. This was continued until complete removal of the pulpolite and release of the canal orifices. Once free, root canal negotiation and instrumentation were carried out step by step, with continuous renewal of the irrigation solution (Fig.14.15.16).














Clinical case N°3 :

Hypertensive 65-year-old patient referred by a colleague for non-localization of root canal orifices on 46 with radiological evidence of diffuse calcifications obliterating the entire ductal network (Fig13).



Root canal negotiation began with pre-curved, small-diameter K files (K6/ K8/ K10), without immediately attempting full root canal catheterization (Fig 17,18).


We also opted for abundant irrigation with sodium hypochlorite, which has a solvent action on the organic substances involved in pulpitis adhesion. The addition of a lubricating and chelating gel (EDTA) facilitated instrument insertion and eliminated the mineral phase of the pulpitis.

Once negotiation to the working length had been achieved, advancement into the canal became easier with rotary Niti instruments always under abundant irrigation (Fig 19,20,21,22).











Discussion :

A comparative study carried out in 2016 "KARIMI.Z" attempted to investigate the prevalence of pulpal calcifications and study their impact on the quality of endodontic treatment [1]. It was shown that calcific degeneration appears to be a source of difficulty that needs to be taken into consideration to improve the quality of endodontic treatment. This can be achieved by adopting an appropriate clinical procedure for each type of calcification.

Preoperative X-rays can be used to determine the location of pulp calculi:

- pulp chamber
- root canal
- chamber and root canal,

and their relationship to the dentinal walls:

*Free: not attached to the dentinal walls and completely surrounded by pulp tissue. Radiographically, it appears as a radiopacity surrounded by a radiolucent halo.

*Adherent: attached to the root or cameral dentinal walls. On radiography, the radiolucent halo separating the dentinal walls from the calcified mass is absent.

*Root canal obliteration: complete disappearance of the root canal passage in a portion or all of the canal.

Depending on the radiological type of pulp calculus, we can orientate our therapeutic strategy:

- VIS-A-VIS of free coronal pulpstones (clinical case N°1); the success of endodontic treatment depends on the correct design of the access cavity. This involves establishing a contour shape linking the projections of the pulp horns at occlusal level. An ordinary cavity is then created and deepened using long-neck tungsten carbide burs. The sensation of emptiness, long associated with entering the pulp chamber, does not occur in the presence of calcified hard tissue. The extent of the pulpstone is delimited by a colorimetric demarcation between the dentin of the canal walls and the central calcified tissue. This demarcation may be manifested by a hemorrhagic line of underlying pulp tissue when the tooth in

question is vital, otherwise it is manifested by a line of necrotic pulp tissue debris. This line of separation is then extended using ultrasonic diamond inserts, under direct visual control (operating microscope / dental magnifying glass) to better distinguish dentin tissue from calcified tissue. Once the calcification has been completely circumscribed, the aim is to remove it in a single block, simply mobilize it with an endodontic excavator to detach it in a single block from the access cavity. The removal of this calcified mass reveals an intact, clean pulp floor and clearly distinguished canal entrances [4,5]

- VIS-A-VIS of adherent coronal pulpitis; the problem is that this type cannot be mobilized despite circumferential clearance. So, once the access cavity has been deepened, the first step is to liberate the canal orifices. Then, slowly abrade the central calcified tissue. The challenge is to know exactly how thick the pulp-stone can be abraded without damaging the floor. Care should be taken with a long-necked ball bur on a low-speed counter-angle or with an ultrasonic diamond insert [4].
- VIS-A-VIS root canal calcifications; the idea in this situation is not to negotiate all the way to the apical construction. This is an iatrogenic error and can lead to the formation of dentinal plugs and abutments, further complicating the management of this type of canal. Once the root canal entrances have been located, root canal penetration can begin. It is always important to remember that instrument fractures are more difficult to treat on mineralized teeth. Full canal catheterization should not be performed from the beginning, preferring instead a crown-down approach [6]. Manual K8 or K10 files, and sometimes even K6 files, are used to progress through the canal, but these instruments are very fragile and should be replaced regularly. After initial penetration, rotary pre-enlargement instruments can be used. It is important not to forget to pre-curve manual instruments over the last few millimetres, to enable them to bypass any mineralization present in the canal [7]. Advancing into the canal can sometimes become easier as the instrument progresses, since secondary dentinogenesis is more marked coronally than apically. However, if the instrument no longer seems to be advancing correctly, you should stop and try to feel how the instrument is engaged in the canal. If there is a sensation of resistance to withdrawal, then the canal is narrowed and you can continue as normal, opening the canal progressively. If, on the other hand, the instrument appears to be free in the canal, we are faced either with total apical calcification, which is very rare, or with an abutment, in which case we need to pre-curve the instrument further to find a zone where we can obtain this sensation of resistance to withdrawal [8]. Abundant irrigation with sodium hypochlorite should always be maintained, as the solvent action of NaOCl enables digestion of the organic substances involved in pulpitis adhesion. Manual filing helps to mobilize the pulpitis in a coronal direction, by establishing low-amplitude back-and-forth movements. The use of chelating agents facilitates progression within the canal by demineralizing the canal walls and eliminating the mineral phase of calcification adhesions [9]. It should also be remembered that, in the long-term, cement deposits form at the apex, altering the position of the radiological apex in relation to the anatomical apex. Root canal preparation must therefore sometimes be carried out away from this radiological apex [4].
- In the case of voluminous root canal calcification and if the clinical situation allows (visibility, accessibility) ultrasonic inserts (ET20, ETD25 Satelec®) are used to destroy this calcific obstruction, always under direct visual microscopic control [4,10,11].

Conclusion

Pulpal calcifications are frequently encountered in the during dental practice, and can take a variety of clinical forms, posing problems of diagnosis and management. The indication for endodontic treatment cannot be based on the presence of pulpitis alone, but rather on a diagnosis supported by a clinical and radiological examination of the pulp status of the tooth. NAOCla thorough understanding of these particularities, the choice of appropriate therapies and the use of optical aids are the keys to successful treatment of mineralized teeth.

Acknowledgements

there are no acknowledgements.

Conflict of interest

Any financial interest or any conflict of interest exists.

Consent

Written informed consent was obtained from the patient to publish this report in accordance with the journal's patient consent policy

Bibliography

1. **Karimi Z et al** ; Les dégénérescences calciques pulpairees et leur impact sur la qualité du traitement endodontique : étude radiographique ; **EDP Sciences AOS** 2016
2. **Goga R et al** ; Pulp stones: a review. *Int Endod J* 2008;41:457–68
3. **Gabardo et al** ; Association between Pulp Stones and Kidney Stones: A Systematic Review and Meta-analysis; **JOE** 2019
4. **Grégory Caron, Dominique Martin** ; Oblitérations canalaires les pulpolithes ; **L'INFORMATION DENTAIRE** n° 1 - 6 janvier 2010
5. **Machtou P** . Réaliser une cavité d'accès idéale dans toutes les situations; **Les dix points clés en endodontie** 2007
6. **McCabe PS, Dummer PMH** . Pulp canal obliteration: an endodontic diagnosis and treatment challenge: Diagnosis and treatment of pulpal obliteration. *Int Endod J*. 2012;45(2):177-197.
7. **Amir FA** , Gutmann JL, Witherspoon DE. Calcific metamorphosis: a challenge in endodontic diagnosis and treatment. *Quintessence Int Berl Ger* 1985. 2001;32(6):447- 455.
8. **Allen PF, Whitworth JM** . Endodontic considerations in the elderly. *Gerodontology* . 2004;21(4):185-194.
9. Julieta Juárez-Gallegos, Characterization of pulp calcifications and changes in their composition after treatments with citric acid and ethylenediaminetetraacetic acid solutions, *Microscopy Research and Technique* 86.1 (2023): 41-52.
10. **andolo A. et al** , Calcifications detection and management in the obliterated endodontic space, *J Osseointegr* 2023;15(1): 48-52.
11. **Chen B. et al** , Removal of calcifications from distal canals of mandibular molars by a non-instrumentational cleaning system: A micro-CT study, *Australian Endodontic Journal* , 2020 46(1)