

Influenza versus COVID-19: Comparison of clinical characteristics and outcomes in a pediatric hospital in Mexico City.

Almudena Laris González¹, Martha Avilés-Robles¹, Clemen Domínguez-Barrera², Israel Parra-Ortega², José Luis Sánchez Huerta¹, Karla Ojeda-Diezbarroso¹, Sergio Bonilla Pellegrini¹, Victor Olivar-López¹, Adrián Chávez López¹, and RODOLFO N Jimenez-Juarez¹

¹Hospital Infantil de Mexico Federico Gomez

²Hospital Infantil de México Federico Gomez

January 30, 2024

Abstract

Introduction: Respiratory viruses are among the leading causes of disease and death among children. Co-circulation of influenza and SARS-CoV2 can lead to diagnostic and management difficulties given the similarities the clinical picture. **Methods:** This is a cohort of all children hospitalized with SARS-CoV2 infection from March to September 3rd 2020, and all children admitted with influenza throughout five flu-seasons (2013-2018) at a pediatric referral hospital. Patients with influenza were identified from the clinical laboratory database. All hospitalized patients with confirmed SARS-CoV2 infection were followed-up prospectively. **Results:** A total of 295 patients with influenza and 133 with SARS-CoV2 infection were included. The median age was 3.7 years for influenza and 5.3 years for SARS-CoV2. Comorbidities were frequent in both groups, but they were more common in patients with influenza (96.6% vs 82.7%, $p < 0.001$). Fever and cough were the most common clinical manifestations in both groups. Rhinorrhea was present in more than half of children with influenza but was infrequent in those with COVID-19 (53.6 vs 5.8%, $p < 0.001$). Overall, 6.4% percent of patients with influenza and 7.5% percent of patients with SARS-CoV2 infection died. In-hospital mortality and the need for mechanical ventilation among symptomatic patients were similar between groups in the multivariate analysis. **Conclusions:** Influenza and COVID-19 have a similar picture in pediatric patients, which makes diagnostic testing necessary for adequate diagnosis and management. Even though most cases of COVID-19 in children are asymptomatic or mild, the risk of death among hospitalized patients with comorbidities may be substantial, especially among infants.

INFLUENZA VERSUS COVID-19: COMPARISON OF CLINICAL CHARACTERISTICS AND OUTCOMES IN A PEDIATRIC HOSPITAL IN MEXICO CITY

Running title: Influenza versus COVID-19 in children.

Authors:

Almudena Laris-González (1), Martha Avilés-Robles (1), Clemen Domínguez-Barrera (1), Israel Parra-Ortega (2), José Luis Sánchez-Huerta (2), Karla Ojeda-Diezbarroso (1), Sergio Bonilla-Pellegrini (1), Víctor Olivar-López (3), Adrián Chávez-López (4), Rodolfo Jiménez-Juárez (1,5).

Affiliation:

1. Hospital Infantil de México Federico Gómez. Department of Infectious Diseases.
2. Hospital Infantil de México Federico Gómez. Department of Clinical Laboratory.
3. Hospital Infantil de México Federico Gómez. Department of Emergency Medicine.

4. Hospital Infantil de México Federico Gómez. Department of Pediatric Intensive Care.

5. Hospital de Infectología "Daniel Méndez Hernández", UMAE CMN La Raza, IMSS. Department of Pediatrics.

Corresponding author : Rodolfo Jiménez-Juárez

Hospital Infantil de México Federico Gómez. Department of infectious Diseases.

Dr. Márquez 162, Cuauhtemoc, 06720, Ciudad de México, México.

Phone number: (52) 55 4342 8293. Email: dr.jimenezjuarez@gmail.com

Acknowledgments: This research was partially supported by a grant from Sanofi Pasteur.

Conflicts of Interest: The authors declare no conflict of interest.

ABSTRACT

Introduction: Respiratory viruses are among the leading causes of disease and death among children. Co-circulation of influenza and SARS-CoV2 can lead to diagnostic and management difficulties given the similarities the clinical picture.

Methods: This is a cohort of all children hospitalized with SARS-CoV2 infection from March to September 3rd 2020, and all children admitted with influenza throughout five flu-seasons (2013-2018) at a pediatric referral hospital. Patients with influenza were identified from the clinical laboratory database. All hospitalized patients with confirmed SARS-CoV2 infection were followed-up prospectively.

Results: A total of 295 patients with influenza and 133 with SARS-CoV2 infection were included. The median age was 3.7 years for influenza and 5.3 years for SARS-CoV2. Comorbidities were frequent in both groups, but they were more common in patients with influenza (96.6% vs 82.7%, $p < 0.001$). Fever and cough were the most common clinical manifestations in both groups. Rhinorrhea was present in more than half of children with influenza but was infrequent in those with COVID-19 (53.6 vs 5.8%, $p < 0.001$). Overall, 6.4% percent of patients with influenza and 7.5% percent of patients with SARS-CoV2 infection died. In-hospital mortality and the need for mechanical ventilation among symptomatic patients were similar between groups in the multivariate analysis.

Conclusions: Influenza and COVID-19 have a similar picture in pediatric patients, which makes diagnostic testing necessary for adequate diagnosis and management. Even though most cases of COVID-19 in children are asymptomatic or mild, the risk of death among hospitalized patients with comorbidities may be substantial, especially among infants.

Keyword: Influenza, SARS-CoV2, COVID-19, children, clinical characteristics, outcome.

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV2), the virus causing COVID-19, has caused a devastating pandemic with over 100 million cases and 2 million deaths globally so far.¹ To date, children have been relatively spared from severe disease and death, representing around 2% of hospitalizations and less than 1 in every 1,000 COVID-19 deaths.² However, pediatric patients with underlying conditions have higher rates of hospitalization and death compared with previously healthy children.³

Other respiratory viral infections such as influenza and respiratory syncytial virus, are among the leading causes of disease and death among pediatric populations.⁴⁻⁷ Co-circulation of influenza and SARS-CoV2 can lead to diagnostic and management difficulties given the similarities in the clinical picture. However, data directly comparing influenza and Covid-19 are scarce, especially in pediatric populations.

The objective of our study was to describe the clinical characteristics and outcomes of children hospitalized with SARS-CoV2 in a pediatric referral hospital and compare them to a cohort of hospitalized children with influenza in the same center.

METHODS

This is a cohort of all children hospitalized with SARS-CoV2 infection from March to September 3rd 2020, and all children admitted with influenza throughout five flu-seasons (2013-2018) at Hospital Infantil de México Federico Gómez, a national referral pediatric hospital with 290 beds and more than 7,000 discharges per year.

Patients with influenza were identified from the clinical laboratory database. Detection of respiratory viruses by PCR multiplex started in 2012 but was established as standard care of children with Severe Acute Respiratory Infection in 2013. We reviewed all records of the subjects with detection of influenza by PCR and included only those that had been hospitalized.

All hospitalized patients with confirmed SARS-CoV2 infection were followed-up prospectively. PCR test was performed on nasopharyngeal swabs of suspect cases beginning in March 2020, and to all admitted patients regardless of symptoms from June 2020.

We collected demographic and clinical variables from the clinical records, including age, sex, comorbidities, symptoms, admission to PICU (pediatric intensive care unit), need for mechanical ventilation and death, among others.

Microbiology

Nasopharyngeal samples were kept at 2-8°C and processed within 24 hours in the SARS-CoV2 cohort and 24 hours (up to 72 hours in weekends) in the influenza cohort.

In the influenza cohort, we used an RT-PCR system with microarray visualization (CLART PneumoVir, Genomica, Spain) capable of detecting adenovirus, bocavirus, coronavirus, rhinovirus, enterovirus, influenza virus A (subtypes AH3N2, AH1N1), influenza virus B, human metapneumovirus (subtypes A and B), parainfluenza virus 1, 2, 3, and 4 (subtypes A and B) and respiratory syncytial virus type A (RSV-A) and B (RSV-B), with a sensitivity of 83.3–100% depending on the virus. In the case of influenza, the sensitivity and specificity was 91% and 99% for influenza A, 82% and 99% for influenza B, respectively.⁸

Two different platforms were used for the detection of SARS-CoV2 throughout the duration of the study: TaqMan 2019-nCoV Assay Kit v1 (Thermo Fisher Scientific) and 2019 Novel Coronavirus (2019- nCoV) RNA (PCR-Fluorescence Probing) (Da An Gene Co. Ltd. Sun Yat-Sen University). RNA was extracted and purified from all samples using the QIAcube (QIAGEN) system and the QIAamp(r) Viral RNA mini kit (QIAGEN) following standard manufacturer procedure.

Statistical analysis

Categorical data are presented as frequencies and percentages, and quantitative data are provided as median and interquartile range (IQR). Demographic, clinical and outcome variables were compared between the two groups of patients, using Fisher's exact test for qualitative variables and Mann-Whitney U test for quantitative variables. The comparisons were carried out for all patients and by age group (younger than 1 year, 1 to 9 years and 10 years or older).

We estimated the crude and adjusted odds ratio and 95% confidence intervals (CI) for death and invasive mechanical ventilation using logistic regression. The final model included age a priori; other variables (sex, nutritional status, comorbidities) were included based on reverse stepwise regression (i.e., stopping rule Wald test <0.20). All analyses were performed using STATA 15 statistical package. (College Station, TX, USA: StataCorp).

RESULTS

A total of 295 patients with influenza and 133 with SARS-CoV2 infection were included, of whom 52% and 55% were male, respectively. The median age was 3.7 years (IQR 1.4-7.7) for influenza and 5.3 years (IQR 1.1-11.2) for SARS-CoV2. See table 1 and table S1 for the comparison between the two groups overall and by age group.

Comorbidities were frequent in both groups, but they were more common in patients with influenza (96.6% vs 82.7%, $p < 0.001$). Cancer, immunodeficiency states, and heart disease were the most common comorbidities in both groups. Chronic pulmonary disease and congenital malformations were more common among children hospitalized with influenza.

Twenty-two percent of patients with SARS-CoV2 had no symptoms attributable to the infection, while only 2 patients with influenza had subclinical infection. When comparing different age groups, infants were more likely to have asymptomatic SARS-CoV2 infection (48.3% versus 16.7% among children 1 to 9 years old, and 13.6% among those 10 years of age and older, $p=0.002$ by Fisher's exact test).

Among symptomatic children, fever and cough were the most common clinical manifestations in both groups, but cough was more than twice as common in patients with influenza (75.8% vs 30.1%, $p < 0.001$). Rhinorrhea was present in more than half of children with influenza but was infrequent in those with Covid-19 (53.6 vs 5.8%, $p < 0.001$). On the contrary, vomiting (29.1 vs 14.3%, $p = 0.002$) and oxygen saturation $< 92\%$ at hospital admission (40.8% vs 29.3%, $p = 0.04$) occurred more commonly in children with SARS-CoV2 infection, but this difference was only observed among those 10 years and older when age subgroups were analyzed.

Admission to the intensive care unit was more common in patients with Covid-19, but the need for invasive mechanical ventilation was similar between groups (see table 2). Overall, 6.4% percent of patients with influenza and 7.5% percent of patients with SARS-CoV2 infection died. When considering different age groups, a fatal outcome occurred in 12.7% of infants, 4.3% of children 1 to 9 years old and 7.7% of those 10 years and older with influenza ($p=0.08$ for the difference between age groups). The rate was 10.3%, 6.7% and 6.8% in the same age groups among patients with SARS-CoV2 infection ($p=0.83$ for the difference between age groups).

In-hospital mortality among symptomatic patients was similar between groups in the multivariate analysis, with an odds ratio of death of 1.87 (0.78-4.45, $p=0.16$) among patients with Covid-19 as compared to those with influenza after adjusting for age group (< 1 , 1-9, > 10 years), pulmonary disease and malignant neoplasia (See table 3 and Table S2). Age 1 to 9 years was associated with lower odds of death (OR 0.27, 95% CI 0.19-0.75, $p=0.01$) in the logistic regression model for the whole population, while a diagnosis of cancer conveyed higher odds of in-hospital mortality (OR 2.72, 95% CI 1.06-6.95, $p=0.04$).

The same was true for the need for invasive mechanical ventilation, with an odds ratio of 0.86 (0.49-1.51) among patients with Covid-19 as compared to those with symptomatic influenza, after accounting for age group, pulmonary disease, and immunodeficiency (See table S3).

DISCUSSION

Respiratory viruses are among the leading causes of pediatric morbidity and mortality worldwide. Unlike other viral respiratory diseases, COVID-19 has had a relatively limited impact on children as compared to adults.⁹ However, as lockdowns are eased and schools reopened, the increased contact between unvaccinated children might change the scenario and increase the circulation of both SARS-CoV2 and other respiratory viruses, including influenza.¹⁰

In this cohort comparing the clinical characteristics and outcomes of SARS-CoV2 infection with epidemic influenza in hospitalized pediatric patients in a single referral hospital, we found a higher proportion of asymptomatic infection in children with SARS-CoV2 than in those with influenza. Almost half of infants with SARS-CoV2 infection had no symptoms of COVID-19. However, the greater proportion of asymptomatic infection may be explained by the different sampling approach, with PCR testing performed on all admissions beginning in June.

Fever and cough were the most common manifestations among both groups, but rhinorrhea was rare in patients with COVID-19 while it was common in children with influenza. The absence of nasal symptoms might be suggestive of COVID-19 in pediatric patients with fever and cough. However, most signs and symptoms were similar between the two groups, making clinical distinction unreliable.

In comparison, in a retrospective cohort of 315 pediatric patients with COVID-19 (median age 8.3 years) and 1402 with influenza (median age 3.9 years) in the United States, Song et al.¹¹ describe a higher frequency of fever, gastrointestinal symptoms, headache, myalgia and chest pain among hospitalized children with COVID-19. However, patients with COVID-19 were older than those with influenza (median age 8.3 years, versus 3.9 years) which might have biased the reporting of symptoms.

Another retrospective single-center cohort study found a lower Charlson index in the COVID-19 group, as well as a greater frequency of anosmia, dysgeusia, diarrhea and frontal headache, and lower prevalence of dyspnea, conjunctivitis and vomiting. No children were included in this cohort.¹²

Regarding outcome, hospitalized patients with symptomatic influenza and COVID-19 had a similar risk of invasive mechanical ventilation and death, and one in ten infants had a fatal outcome. This underscores that even though COVID-19 is usually mild in children, it can cause severe disease and death in children with chronic diseases, as has been recognized for influenza for a long time⁴. Mortality was lowest among children 1 to 9 years old with either influenza or SARS-CoV2 infection, but differences between age groups were not statistically significant. This is consistent with reports from the United States, where the greatest proportion of COVID-19 pediatric deaths occurred in infants and adolescents¹³ and deaths from influenza are more frequent among those younger than 6 months.¹⁴

The aforementioned study by Song et al.¹¹ reported a similar rate of hospitalization, intensive care unit admission and use of mechanical ventilation. Two deaths were reported among patients with influenza while no patients with COVID-19 died.

In France, Piroth et al.¹⁵ describe a higher in-hospital mortality for patients with COVID-19 in comparison with those with influenza, based in a nationwide retrospective cohort of hospitalized patients. The proportion of pediatric patients was smaller for COVID-19 than for influenza, but in-hospital mortality was ten-times greater for COVID-19 in those 11-17 years old, and a larger proportion of patients younger than 5 years needed intensive care support for COVID-19 than for influenza.

A systematic review and meta-analysis of studies describing individuals with either influenza or COVID-19 reported a lower frequency of nasal symptoms, pharyngodynia and dyspnea, and a greater prevalence of radiographic abnormalities among patients with COVID-19. Case fatality rate of hospitalized patients was 6.5%, 6% and 3% respectively for individuals with COVID-19, influenza A and influenza B. However, pediatric patients represented a small fraction of the cases and findings were limited by the heterogeneity of the studies.¹⁶

Zhang et al.¹⁷ describe two cohorts of patients hospitalized with influenza or COVID-19 in two separate locations. One in every five patients with COVID-19 was admitted to ICU and 13% died, while no severe or fatal cases were recorded in the influenza cohort. No data for pediatric patients was reported in this study.

Our study has several limitations worth noting. The single-center nature of the cohort, as well as the frequency of comorbidities among participants, may limit the generalizability of findings. Data on patients with influenza was collected retrospectively from clinical records while the COVID-19 cohort was followed-up prospectively. The different time periods between the two cohorts may account for differences in diagnostic and therapeutic approaches. As mentioned above, all patients admitted to hospital were screened for SARS-CoV2 infection from June 2020, which explains the greater proportions of asymptomatic infections. However, asymptomatic patients were excluded from the outcome analysis to account for this limitation. Non-invasive mechanical ventilation was discouraged for patients with suspected or confirmed COVID-19 to reduce aerosol generation, which might have increased the proportion of patients receiving invasive mechanical ventilation.

In conclusion, influenza and COVID-19 have a similar picture in pediatric patients, which makes diagnostic testing necessary for adequate diagnosis and management and will add to the challenge of co-circulation as SARS-CoV2 becomes endemic. Even though most cases of COVID-19 in children are asymptomatic or mild, the risk of death among hospitalized children with comorbidities is substantial, especially among infants, and is similar to that of patients with influenza. Thus, children should not be left out of preventive and

therapeutic development in the COVID-19 pandemic, including vaccine development programs.¹⁸

REFERENCES

1. World Health Organization. WHO (COVID-19) Homepage. Accessed February 1, 2021. <https://covid19.who.int/>
2. Wilder JL, Parsons CR, Growdon AS, Toomey SL, Mansbach JM. Pediatric Hospitalizations During the COVID-19 Pandemic. *Pediatrics* . 2020;146(6). doi:10.1542/peds.2020-005983
3. Stokes EK, Zambrano LD, Anderson KN, et al. Coronavirus Disease 2019 Case Surveillance - United States, January 22-May 30, 2020. *MMWR Morb Mortal Wkly Rep* . 2020;69(24):759-765. doi:10.15585/mmwr.mm6924e2
4. Chaves SS, Perez A, Farley MM, et al. The burden of influenza hospitalizations in infants from 2003 to 2012, United States. *Pediatr Infect Dis J* . 2014;33(9):912-919. doi:10.1097/INF.0000000000000321
5. Shi T, McAllister DA, O'Brien KL, et al. Global, regional, and national disease burden estimates of acute lower respiratory infections due to respiratory syncytial virus in young children in 2015: a systematic review and modelling study. *Lancet (London, England)* . 2017;390(10098):946-958. doi:10.1016/S0140-6736(17)30938-8
6. Bourgeois FT, Valim C, Wei JC, McAdam AJ, Mandl KD. Influenza and Other Respiratory Virus-Related Emergency Department Visits Among Young Children. *Pediatrics* . 2006;118(1):e1 LP-e8. doi:10.1542/peds.2005-2248
7. Troeger CE, Blacker BF, Khalil IA, et al. Mortality, morbidity, and hospitalisations due to influenza lower respiratory tract infections, 2017: an analysis for the Global Burden of Disease Study 2017. *Lancet Respir Med* . 2019;7:69-89. doi:10.1016/S2213-2600(18)30496-X
8. Huang HS, Tsai CL, Chang J, Hsu TC, Lin S, Lee CC. Multiplex PCR system for the rapid diagnosis of respiratory virus infection: Systematic review and meta-analysis. *Clin Microbiol Infect* . 2018;24:1055-1063.
9. Sisk B, Cull W, Harris JM, Rothenburger A, Olson L. National Trends of Cases of COVID-19 in Children Based on US State Health Department Data. *Pediatrics* . 2020;146(6). doi:10.1542/peds.2020-027425
10. Krishnaratne S, Pfadenhauer LM, Coenen M, et al. Measures implemented in the school setting to contain the COVID-19 pandemic: a rapid scoping review. *Cochrane Database Syst Rev* . 2020;(12). doi:10.1002/14651858.CD013812
11. Song X, Delaney M, Shah RK, Campos JM, Wessel DL, DeBiasi RL. Comparison of Clinical Features of COVID-19 vs Seasonal Influenza A and B in US Children. *JAMA Netw Open* . 2020;3(9):e2020495-e2020495. doi:10.1001/jamanetworkopen.2020.20495
12. Zayet S, Kadiane-Oussou NJ, Lepiller Q, et al. Clinical features of COVID-19 and influenza: a comparative study on Nord Franche-Comte cluster. *Microbes Infect* . 2020;22(9):481-488. doi:10.1016/j.micinf.2020.05.016
13. Bixler D, Miller AD, Mattison CP, et al. SARS-CoV-2 – Associated Deaths Among Persons Aged < 21 Years — United States , February 12 – July 31 , 2020. *MMWR Morb Mortal Wkly Rep* . 2020;69:1324-1329. doi:http://dx.doi.org/10.15585/mmwr.mm6937e4external icon
14. Shang M, Blanton L, Brammer L, Olsen SJ, Fry AM. Influenza-Associated Pediatric Deaths in the United States, 2010-2016. *Pediatrics* . 2018;141(4). doi:10.1542/peds.2017-2918
15. Piroth L, Cottenet J, Mariet A, et al. Comparison of the characteristics, morbidity, and mortality of COVID-19 and seasonal influenza: a nationwide, population-based retrospective cohort study. *Lancet Respir* . 2020;2600(20):1-9. doi:10.1016/S2213-2600(20)30527-0

16. Pormohammad A, Ghorbani S, Khatami A, et al. Comparison of influenza type A and B with COVID-19: A global systematic review and meta-analysis on clinical, laboratory and radiographic findings. *Rev Med Virol* . Published online October 9, 2020:e2179-e2179. doi:10.1002/rmv.2179
17. Zhang J, Ding D, Huang X, et al. Differentiation of COVID-19 from seasonal influenza: A multicenter comparative study. *J Med Virol* . Published online August 28, 2020:10.1002/jmv.26469. doi:10.1002/jmv.26469
18. Kao CM, Orenstein WA, Anderson EJ. The Importance of Advancing Severe Acute Respiratory Syndrome Coronavirus 2 Vaccines in Children. *Clin Infect Dis an Off Publ Infect Dis Soc Am* . 2021;72(3):515-518. doi:10.1093/cid/ciaa712

Table 1. Demographic and clinical characteristics of patients with influenza and SARS-CoV2 infection.

Male sex, n (%)
Age, median (IQR)
Any comorbidity Cardiovascular Asthma Pulmonary Immunodeficiency Kidney disease Cancer Congenital malformations
Nutritional status Undernourished Overweight/Obesity
Nosocomial acquisition
Symptoms at diagnosis Asymptomatic Fever Cough Pharyngodynia Headache Myalgia Laboured breathing Rhinorrhea Vom
SaO2 <92% at hospital admission
Hypotension

SaO2: Oxygen saturation.

+ Fisher's exact test or Mann-Whitney test, as appropriate.

Table 2. Main outcomes of patients with influenza and SARS-CoV2 infection.

	Total population Influenza (n=295)	Total population SARS-CoV2 (n=133)	Total population p value+	Symptomatic inf Influenza (n=293)
PICU admission	55 (18.6%)	45 (33.8%)	0.001	54 (18.4%)
Non-invasive mechanical ventilation	127 (43%)	12 (9%)	<0.001	126 (43%)
Invasive mechanical ventilation	67 (22.7%)	27 (20.3%)	0.62	67 (22.9%)
High-frequency oscillatory ventilation	12 (4.1%)	3 (2.3%)	0.41	12 (4.1%)
Severe ARDS (PaO2/FiO2 index <100)	19 (6.4%)	3 (2.3%)	0.096	19 (6.4%)
Death	19 (6.4%)	10 (7.5%)	0.68	19 (6.5%)

ARDS: Acute respiratory distress syndrome, PaO2: Partial pressure of oxygen, FiO2: Fraction of inspired oxygen.

+Fisher's exact test or Mann-Whitney test, as appropriate.

Table 3. Crude and adjusted estimates of the odds ratio for in-hospital death and mechanical ventilation estimated by logistic regression among patients with symptomatic SARS-CoV2 infection in comparison to patients with symptomatic influenza.

Outcome (%)	Crude OR	Adjusted OR	p value
-------------	----------	-------------	---------

Death Influenza (N=293)	19 (6.5%)	9 (8.7%)	1.38 (0.60-3.16)	1.87 (0.78-4.45) +	0.16
SARS-CoV2 (N=103)					
Mechanical ventilation Influenza (N=293)	67 (22.9%)	22 (21.4%)	0.92 (0.53-1.58)	0.83 (0.46-1.49) ++	0.53
SARS-CoV2 (N=103)					

+Adjusted for age group (<1, 1-10, >10 years), pulmonary disease and malignant neoplasia.

++Adjusted for age group (<1, 1-10, >10 years), pulmonary disease and immunodeficiency.

Supplementary Table 1 . Clinical characteristics and outcome of patients with influenza and SARS-CoV2 infection by age group.

Male sex, n(%)
Comorbidities Cardiovascular Pulmonary Immune deficiency Kidney disease Cancer Congenital malformations
Nosocomial acquisition
Symptoms at diagnosis Asymptomatic Fever Cough Pharyngodinia Headache Laboured breathing Rhinorrhea Vomiting Dia
SaO2 <92% at hospital admission
Hypotension
Outcome PICU admission NIMV IMV HFOV PaO2/FiO2 index <100 Death

SaO2: Oxygen saturation, NIV: Non-invasive mechanical ventilation, IMV: Invasive mechanical ventilation, HFOV: High-frequency oscillatory ventilation. PaO2: Partial pressure of oxygen, FiO2: Fraction of inspired oxygen, NA: Not applicable.

Supplementary table 2. Results of the logistic regression model for in-hospital mortality among patients with symptomatic SARS-CoV2 infection in comparison to patients with symptomatic influenza (baseline).

Variable	Odds Ratio	95% CI	p value
SARS-CoV2 vs influenza	1.77	0.73-4.29	0.15
Age group < 1 year 1 to 9 years [?]10 years	Reference 0.27 0.41	0.19-0.75 0.12-1.32	0.01 0.13
Sex	0.57	0.25-1.30	0.18
Pulmonary disease	2.58	0.89-7.46	0.08
Cancer	2.72	1.06-6.95	0.04

CI: Confidence interval.

Supplementary table 3 . Results of the logistic regression model for invasive mechanical ventilation among patients with symptomatic SARS-CoV2 infection in comparison to patients with symptomatic influenza (baseline).

	Odds Ratio	95% CI	p value
SARS-CoV2 vs influenza	0.83	0.46-1.49	0.53
Age group < 1 year 1 to 9 years [?]10 years	Reference 0.60 1.10	0.32-1.11 0.52-2.33	0.10 0.80

	Odds Ratio	95% CI	p value
Pulmonary disease	1.75	0.88-3.49	0.11
Immunodeficiency	0.42	0.22-0.80	0.008

CI: Confidence interval.