

Katrine Gorham¹, Sam Abernethy¹, Tyler R Jones², Peter Hess³, Natalie M Mahowald⁴, Daphne Meidan⁵, Matthew S Johnson⁶, Maarten M J W Van Herpen⁷, Yangyang Xu⁸, Alfonso Saiz-Lopez⁵, Thomas Röckmann⁹, Chloe Brashear⁹, Erika Reinhardt¹, and David Mann¹

¹Spark Climate Solutions

²Institute of Arctic and Alpine Research, University of Colorado

³Department of Biological and Environmental Engineering, Cornell University

⁴Department of Earth and Atmospheric Sciences, Cornell University

⁵Department of Atmospheric Chemistry and Climate, Institute of Physical Chemistry Blas Cabrera, CSIC

⁶Department of Chemistry, University of Copenhagen

⁷Acacia Impact Innovation BV

⁸Department of Atmospheric Sciences, Texas A&M University

⁹Department of Physics, Institute for Marine and Atmospheric Research Utrecht, Utrecht University

December 7, 2023

Opinion: Exploring potential atmospheric methane removal approaches: an example research roadmap for chlorine radical enhancement

5 Katrine A. Gorham¹, Sam Abernethy¹, Tyler R. Jones², Peter Hess³, Natalie M. Mahowald⁴, Daphne Meidan⁵, Matthew S. Johnson⁶, Maarten M.J.W. van Herpen⁷, Yangyang Xu⁸, Alfonso Saiz-Lopez⁵, Thomas Röckmann⁹, Chloe A. Brashear⁹, Erika Reinhardt¹, David Mann¹

¹Spark Climate Solutions, Covina, CA 91723, USA

²Institute of Arctic and Alpine Research, University of Colorado, Boulder, CO 80303, USA

³Department of Biological and Environmental Engineering, Cornell University, Ithaca, NY 14853, USA

10 ⁴Department of Earth and Atmospheric Sciences, Cornell University, Ithaca, NY 14853, USA

⁵Department of Atmospheric Chemistry and Climate, Institute of Physical Chemistry Blas Cabrera, Spanish National Research Council, 28006 Madrid, Spain

⁶Department of Chemistry, University of Copenhagen, DK-2100 Copenhagen Ø, Denmark

⁷Acacia Impact Innovation, Bernheze 5384 BB, The Netherlands

15 ⁸Department of Atmospheric Sciences, Texas A&M University, College Station, TX, 77843, USA

⁹Institute for Marine and Atmospheric Research Utrecht, Department of Physics, Faculty of Science, Utrecht University, 3584 CS Utrecht, The Netherlands

20 *Correspondence to:* Katrine A. Gorham (katrine@sparkclimate.org)

Abstract. The escalating climate crisis requires rapid action to reduce the concentrations of atmospheric greenhouse gases and lower global surface temperatures. Methane will play a critical role in near-term warming due to its high radiative forcing and short atmospheric lifetime. Methane emissions have accelerated in recent years and there is significant risk and uncertainty associated with the future growth in natural emissions. The largest natural sink of methane occurs through oxidation reactions with atmospheric hydroxyl and chlorine radicals. Enhanced atmospheric oxidation could be a potential approach to remove atmospheric methane. One method proposes the addition of iron salt aerosols (ISA) to the atmosphere, mimicking a natural process proposed to occur when mineral dust mixes with chloride from sea spray to form iron chlorides, which are photolyzed by sunlight to produce chlorine radicals. Under the right conditions, lofting ISA into the atmosphere could potentially reduce atmospheric methane concentrations and lower global surface temperatures. Recognizing that potential atmospheric methane removal must only be considered as an additive measure – in addition to, not replacing, crucial anthropogenic greenhouse gas emission reductions and carbon dioxide removal – roadmaps can be a valuable tool to organize and streamline interdisciplinary and multifaceted research to efficiently move towards understanding whether an approach may be viable and socially acceptable, or if it is nonviable and further research should be deprioritized. Here we present an example five-year research roadmap to explore whether ISA enhancement of the chlorine radical sink could be a viable and socially acceptable atmospheric methane removal approach.

25
30
35

1 Introduction

Driven by anthropogenic greenhouse gas emissions, Earth's average surface temperature has increased by 1.1 °C since 1850 (Forster et al., 2021). As global temperature increases, irreversible changes in the Earth system will likely occur, such as ice sheet collapses in Greenland and Antarctica, coral reef die-off, abrupt permafrost thaw, mountain glacier loss, and Amazon rainforest dieback (Lenton et al., 2008; McKay et al., 2022). To mitigate both near-term and long-term warming we must undertake rapid, sustained action to establish a diverse portfolio of approaches to slow and then reverse the increase of atmospheric greenhouse gas concentrations, ideally reducing them to preindustrial levels. Emissions reductions must be prioritized. In addition, as near-term warming threatens to trigger climate tipping points, negative emissions approaches may be used to remove greenhouse gases already in the atmosphere and counter rising natural and uncontrollable emissions.

Methane (CH₄) has contributed 0.5 °C to the current 1.1 °C temperature increase relative to preindustrial times, second only to carbon dioxide (CO₂) (IPCC, 2023). Methane lasts roughly a decade in the atmosphere (UNEP, 2021), with a Global Warming Potential 83 times that of carbon dioxide over 20 years and 30 times that of carbon dioxide over 100 years (Forster et al., 2021). The concentration of methane in the atmosphere is 2.5 times pre-industrial levels, and the growth rate has accelerated since 2006, with record increases in 2020 and 2021 (NOAA, 2023). Methane emissions come from anthropogenic sources (fossil fuel use, agriculture, waste and wastewater, biomass burning, etc.) and natural sources (wetlands, oceans, freshwaters, termites, permafrost, etc.), both of which are increasing (Jackson et al., 2020; Nisbet et al., 2023, Saunio et al., 2020). As the planet continues to warm and precipitation patterns change, natural methane emissions are expected to increase from wetlands, as well as from permafrost due to abrupt thaw, thermokarst lake formation and expansion, and bacterial processes (Dean et al., 2018; Neumann et al., 2019; Paudel et al., 2016; Peng et al., 2022; Zhang et al., 2023). One estimate is that natural emissions could increase by 20% (~50 Tg CH₄/yr) to 150% (~330 Tg CH₄/yr) by 2100, becoming larger than anthropogenic emissions in most modeled scenarios (Kleinen et al. 2021). There is evidence that wetland emissions increases are already underway, with roughly half of the 2020 atmospheric methane increase attributed to wetlands (Peng et al., 2022; Nisbet et al., 2023; Zhang et al., 2023).

The natural sinks for atmospheric methane are oxidation by gas phase radicals (~95 %) and uptake into soil by methanotrophic bacteria and archaea (~5 %) (Saunio et al., 2020). The atmospheric lifetime of methane is thus mainly determined by the oxidative capacity of the atmosphere. Approximately 550 Tg methane per year is removed by reactions with hydroxyl radicals (OH.); chlorine radicals (Cl·) destroy ~11 Tg methane per year (Saunio et al., 2020). However, the chlorine radical sink estimate is poorly constrained, ranging from 1 to 35 Tg methane per year (Saunio et al., 2020). Recent research suggests the atmospheric chlorine radical sink could be underestimated (van Herpen et al., 2023).

Complex and non-linear atmospheric chemistry dictates the oxidation of methane, which results in the production of formaldehyde and carbon monoxide which is further oxidized to carbon dioxide. Hydroxyl radical oxidation of methane can result in the formation or loss of ozone, depending on the nitrogen oxide (NO_x) concentrations. Chlorine radicals react to remove both methane and ozone, the principal precursor of hydroxyl radical production (Lelieveld et al., 2002; Seinfeld and Pandis, 2016). Therefore, chlorine radicals may either increase or decrease the atmospheric lifetime of methane depending on the concentration of chlorine radicals and atmospheric conditions (Li et al., 2022, van Herpen et al., 2023, Li et al., 2023). Volatile organic compounds and NO_x will change the methane response since they directly and indirectly affect the concentrations of hydroxyl and chlorine radicals. Furthermore, since reactions with carbon monoxide and methane are the primary sinks for hydroxyl radicals, the abundance of methane impacts the oxidative character of the atmosphere in a self feedback process (Lelieveld et al., 2002; Staniaszek et al., 2022).

Enhanced atmospheric oxidative sinks could increase the rate of atmospheric methane removal and therefore reduce near-term warming. One untested proposal involves iron salt aerosols (ISA). This potential approach involves lofting iron-based particles into the troposphere (e.g., from ships or towers) to catalytically produce chlorine radicals (Oeste, 2009; Oeste et al., 2017), mimicking a natural phenomenon proposed to occur when mineral dust combines with sea spray aerosols (van Herpen et al., 2023). Discussing natural analogues of this process and the current state of research, this paper presents a roadmap for research and development that is needed to understand whether ISA enhancement of the chlorine radical sink may be a feasible, scalable, and safe approach for atmospheric methane removal.

2 State of the research

Currently, research into ISA falls into three categories: laboratory experiments to quantify the details of the mechanism, observational analysis of the natural analogue of ISA, known as mineral dust sea spray aerosol (MDSA), and numerical modeling evaluating the potential impacts of ISA.

90 2.1 Laboratory experiments

In a series of papers, Wittmer et al. demonstrated the production of chlorine atoms from iron-doped salts and aerosols (Wittmer et al., 2015a; Wittmer et al., 2015b; Wittmer et al., 2017). Reproducing and expanding upon these laboratory studies is of the highest importance, since the mechanism of chlorine generation is poorly understood (van Herpen et al., 2023; Wittmer et al., 2015; Wittmer et al., 2017; Zhu et al., 1997). The ISA mechanism is catalytic in iron and may be catalytic in chlorine in natural environments (Wittmer et al., 2017). Understanding of catalytic efficiency requires study of dependencies on conditions such as aerosol size distribution and number density; humidity; rate of supply of acidity to the system; rate of coagulation of aerosols; effects of organic chelating agents on iron activity; changes in chlorine escape probability due to aqueous chemical conditions; changes in iron activity due to aerosol chemistry; and, understanding how the atmosphere behaves under high ClO_x

conditions (Pennacchio et al., submitted). Furthermore, studies of real atmospheric conditions are needed to better understand
100 suppression of the ISA mechanism in the presence of sulfate and oxalate (Wittmer et al., 2015; Wittmer et al., 2017).

2.2 The natural analogue of ISA: mineral dust sea spray aerosol

The natural mineral dust sea spray aerosol (MDSA) mechanism is proposed to occur when iron from mineral dust mixes with
the chlorine in sea spray, forming iron chlorides which are photolyzed to produce chlorine radicals (van Herpen et al., 2023).
ISA could mimic this natural MDSA phenomenon by only aerosolizing what is currently believed to be the key component of
105 the mineral dust: the photoactive iron.

van Herpen et al. (2023) provided the first evidence that methane may be removed over the North Atlantic by chlorine radicals
produced by the MDSA mechanism. The semi-arid regions of North Africa are the dominant source of iron-containing mineral
dust, with frequent transport over the North Atlantic (Prospero et al., 2021). Using air samples collected during North African
110 dust events from 1996-2000 in Barbados (Mak et al., 2003), a model parameterized with the MDSA mechanism of chlorine
radical production produced results consistent with a previously unexplained ^{13}C -depletion in the reaction product carbon
monoxide (CO) (van Herpen et al., 2023). Carbon monoxide produced from chlorine radical oxidation of methane is extremely
depleted in ^{13}C which makes $\delta^{13}\text{C}(\text{CO})$ a very sensitive indirect detection method of chlorine radicals (Röckmann et al.,
1999). The variability of $\delta^{13}\text{C}(\text{CO})$ in atmospheric air is the main evidence that the MDSA process is occurring, as there is no
115 other mechanism proposed that can explain the carbon monoxide isotope signature (Mak et al., 2003).

As a proxy for methane oxidation by chlorine radicals, studies of the isotopic composition of carbon monoxide in the mid-
Atlantic boundary layer are underway to further confirm the MDSA mechanism. This includes a regular air sampling program
at four ground-based stations in the North Atlantic, as well as Atlantic transect sampling onboard commercial vessels. Time
120 series at the ground-based stations provide high-resolution observations at the longitudinal margins of trans-Atlantic dust
transport, including the site where seasonal depletions in $\delta^{13}\text{C}(\text{CO})$ were previously observed (Mak et al., 2003). Air samples
collected in north-south ship transects may allow spatial correlation of the $\delta^{13}\text{C}(\text{CO})$ excursions with North African dust
plumes. Together, these samples provide an opportunity to investigate the formation process of MDSA, as well as the seasonal
and spatial influence of North African dust on tropospheric chlorine radical oxidation.

125 2.3 Modeling

Early modeling studies indicate that atmospheric chlorine additions can increase or decrease methane concentrations,
depending on the concentration of chlorine that is present in the atmosphere (Horowitz et al., 2020; Li et al., 2022; Saiz-Lopez
et al., 2023). Chlorine radicals readily oxidize ozone, the main precursor of hydroxyl radicals, which results in less methane
oxidation via hydroxyls; at low atmospheric concentrations, chlorine radicals reduce ozone concentrations without having a
130 commensurate impact on methane. Even though chlorine radicals react 16x faster with methane than the reaction of hydroxyl

radicals with methane (Atkinson, 2006), hydroxyl radicals dominate the methane oxidation sink because they are much more abundant than chlorine radicals. As more chlorine is emitted, ozone concentrations will be reduced so that proportionally more chlorine radicals react with methane. The increased methane destruction by chlorine radicals will eventually outcompete decreased destruction by hydroxyl. In an initial, highly simplified model scenario, Li et al. (2023) found that a reduction in methane concentration could be achieved if more than 90 Tg Cl/yr (three times the estimated present-day emission rate) was added evenly to the atmosphere over all ocean surfaces. However, assuming a uniform increase of chlorine over all ocean surfaces may underestimate the potential effectiveness of local chlorine radical generation where efficiency may be condition-dependent (e.g., NO_x, CO, and chlorine concentrations, humidity, temperature, altitude, etc.) (Meidan et al., submitted). For example, considering the MDSA natural analogue of ISA, van Herpen et al (2023) found that high dust concentrations in the North Atlantic corresponded with net methane removal, while globally lower dust concentrations led to a net increase of methane.

Current models may not accurately capture the speed and efficiency of producing chlorine radicals via the ISA mechanism due to assumptions of the percentage of photoactive iron in the emitted iron, aerosol pH, aerosol mixing rates, and more. Another challenge with Earth system models is that they instantaneously dilute emissions to model grid dimensions which could lead to underestimates or overestimates of the effectiveness of the ISA mechanism, especially when considering iron emission additions from point sources like ships (e.g. Meidan et al., submitted). For example, the mixing of the iron and sea salt within the aerosol is modeled to occur instantaneously (Meidan et al., submitted; van Herpen et al., 2023); however, in reality it would likely take hours to days, leading the global model to overestimate the rate of chlorine radical production. Furthermore, the ISA mechanism is likely to occur faster in high NO_x environments (Oeste et al., 2017) but could be less efficient in high sulfate environments (Bondy et al., 2017; Chen et al., 2020; Legrand et al., 2017; Pio et al., 1998), and both NO_x and sulfate may be co-emitted with iron (e.g. from a ship plume). Thus, models that instantaneously dilute emissions across the grid dimensions may misrepresent the ISA mechanism. Overall, it is unclear whether current Earth system models overestimate or underestimate the efficiency of the oxidation mechanism. Additional detailed box modeling focusing on deployment sites and constrained by field observations are necessary to assess the effectiveness of the mechanism. However, local box models are less reliable over timescales where mixing between different air masses is relevant. This motivates the need for high-resolution regional and seasonal modeling over ocean basins and variable resolution configurations embedded in global models.

Considering the difficulty of simulating variable atmospheric chemical conditions (e.g., atmospheric chemical composition, solar radiation, wind mixing, etc.) across different geographic locations, it is important to develop an ensemble of models that enable uncertainty assessment. Such a model ensemble will allow a comprehensive exploration of different iron salt aerosol deployment scenarios (magnitude each year), aerosol particle sizes, and deployment location and timing.

3 Roadmap

165 3.1 Roadmap framework

Roadmaps are used in climate research to define knowledge gaps, needs, and associated outputs as they relate to interdependencies and timelines (e.g. Wanser et al., 2022; Maesano et al., 2022; Aschwanden et al., 2021; Ocean Visions, 2023), particularly in instances that benefit from integrated, interdisciplinary research. A coordinated, thorough, and science-based approach is needed to ensure that resources are used efficiently, stakeholders and interdisciplinary teams are engaged on appropriate timelines, and efforts are focused towards sequenced research questions and milestones.

3.2 Viability assessment

The viability of an atmospheric methane removal approach can be assessed by considering its potential for feasibility, scalability, and social license to operate. A feasible approach must be climate beneficial, safe, acceptable for its side effects, and cost-plausible. Determination of scalability will be approach-specific, acknowledging that the scale of increased natural emissions is anticipated to be tens of millions of tons of methane per year (Kleinen et al., 2021).

The key milestone questions below can help determine the viability of ISA and whether it should continue to be prioritized. The research that informs the key milestones questions should be pursued in parallel (Table 1).

1. Is enhancement of the chlorine radical oxidative sink of methane via the ISA mechanism effective and climate beneficial? At what scale?
2. What impacts could the ISA approach have on Earth systems and human systems, both positive and negative? Is there a cost-plausible ISA deployment method?
3. What is needed to advance a structure of ethical governance and social license for utilizing atmospheric intervention to reduce atmospheric methane concentrations?

185 3.2.1 Milestone question #1: Is the ISA mechanism effective and climate beneficial and scalable?

The complexity and nonlinearity of atmospheric chemistry and meteorology requires laboratory, field, and plume and global modeling studies of the efficiency of chlorine radical production, its dependence on atmospheric conditions and other gases, and the impact on methane removal.

190 An important assumption in previous studies (van Herpen et al., 2023; Meidan et al., submitted) is that only 2% of iron is photoactive (van Herpen et al., 2023; Meidan et al., submitted). However, the amount of ISA that is photoactive may vary by emission source (e.g. ship emissions may have more photoactive iron relative to mineral dust; Ito, 2013; Rodriguez et al., 2021), geographical location, aerosol pH, the presence of other chemical constituents (e.g. sulfate and NO_x), and altitude

(Mahowald et al., 2018). The efficiency, cost, safety (e.g. air quality), and net radiative forcing of ISA will depend on the
195 percentage of iron that is photoactive.

Current studies assume that the chlorine radicals released from the photochemical reaction with iron will react (e.g., with
methane) to form hydrochloric acid, which will then be reabsorbed back into the aerosol and thus recycled (van Herpen et al.,
2023; Oeste et al., 2017). It is unclear under which atmospheric conditions this cycle occurs, but if some chlorine radicals are
200 lost (e.g. to reactions with ozone instead), then the cycling would be less efficient. After reacting with ozone there is a much
longer chain of reactions before hydrochloric acid is created, thus reducing the cycling efficiency (Wang et al., 2019).
Therefore, further laboratory measurements and detailed box models are needed to further study hydrochloric acid recycling
efficiency by ISA.

205 Better understanding of how sulfur dioxide and NO_x concentrations impact the ISA mechanism is also needed (Oeste et al.,
2017). Sulfur dioxide and NO_x concentrations vary regionally and locally and their emissions may sometimes coincide with
those of iron.

Smaller ISA particles have greater surface area to mass ratios and stay longer in the atmosphere, likely increasing the efficiency
210 of the ISA mechanism. Furthermore, smaller particles tend to be more acidic (Pye et al., 2020), which is required for the
mechanism to be active (Wittmer et al., 2015; 2017). However, smaller aerosols may be transported further to coastal or inland
locations where these particles could contribute directly to negative human health effects or deposit on terrestrial or ice-covered
surfaces with unintended consequences. The aerosol size may also affect marine cloud cover, thereby influencing local
radiative forcing. In addition, there tends to be more sulfate in smaller aerosols, which may reduce the effectiveness of the ISA
215 mechanism (Bondy et al., 2017; Chen et al., 2020; Legrand et al., 2017; Pio et al., 1998).

Studying MDSA (the natural analogue of ISA) through field studies is essential to understand the effectiveness of this
mechanism under different atmospheric conditions and its geographical extent, thereby better constraining atmospheric and
Earth system models. Early MDSA field studies are underway to explore the seasonality and spatial extent of methane
220 oxidation by chlorine radicals that may occur in natural dust plumes through proxy measurements of $\delta^{13}\text{C}(\text{CO})$. Further studies
– both natural analogue and *in-situ* ISA enhancement studies – would benefit from alternative ISA detection and quantification
approaches, including direct chlorine measurements or additional proxy measurements to reinforce existing observations.

Ideally, models will be developed to include the entire MDSA mechanism, including implementation of the isotope effect in
225 the chlorine radical reaction with methane, thus enabling direct comparison of model results to observations of $\delta^{13}\text{C}(\text{CO})$. At
present, some models (EMAC; Gromov 2014) include complete carbon monoxide isotope representation, but not the MDSA

mechanism, whereas other models (CAM-CHem; van Herpen et al., 2023) include an initial representation of the MDSA mechanism but do not incorporate the isotopic effects.

230 Isotopic signatures and dust from ice core paleo records may elucidate evidence of historical MDSA activity. Methane isotope
measurements of air trapped in polar ice cores have been used to constrain the methane budget in the past (Bock et al. 2017;
Sapart et al., 2012; Fischer et al., 2008; Ferretti et al., 2005), but possible variations in the chlorine-based methane sink have
not been taken into account in these studies. Dust levels have undergone strong changes in the past (e.g. Fischer et al., 2007;
Han et al., 2018; Yuan et al., 2020; Yue et al., 2023), and associated changes in MDSA may have affected the paleo records
235 of $\delta^{13}\text{C}(\text{CH}_4)$.

If modeling, laboratory studies, and natural analogue field studies prove that a promising, safe mechanism exists to produce
chlorine radicals at sufficient scale and under diverse atmospheric conditions, it may be appropriate to consider field studies
with intentional enhancement of ISA, using a broad suite of atmospheric measurements to understand how the mechanism
240 performs in the real atmosphere. Prior to performing any ISA enhancement field studies – even at a small and controlled scale
– it is essential to engage and work collaboratively with potentially impacted communities, policy and science leaders,
governmental bodies, NGOs, media, and other stakeholders to ensure that actions are conducted with social license and an
appropriate governance framework with free, prior, and informed consent (FAO, 2016). For example, a study could involve
controlled enhancement of dust or emitted aerosolized iron, or could investigate existing anthropogenic emissions of iron (e.g.
245 from a ship plume, power plant, iron foundry, etc). The scale of the study should be suitable to accommodate a likely non-
linear atmospheric response, where substantial increases in chlorine – thus iron emissions – may be needed before there is a
decrease in methane.

3.2.2 Milestone question #2: What are the potential Earth system and human systems impacts of ISA, and is it cost-plausible?

250 Lifecycle analyses are necessary to assess the potential benefits, tradeoffs, risks, uncertainties, and costs of ISA. As part of
this analysis, understanding the potential impacts of ISA enhancement on the Earth system and human system is imperative
before considering large scale deployment. Human system impacts may include human health outcomes, as well as indirect
human impacts from Earth system changes. For example, if ISA resulted in ocean acidification there could be marine life
implications which could have far reaching ecosystem impacts, as well as economic (e.g. fisheries) and cultural (e.g.
255 Indigenous practices) impacts for coastal communities.

Earth system modeling must be conducted to understand the impact of atmospheric conditions, aerosol size, and release
locations, magnitudes, and timing, on ocean systems, terrestrial systems, the cryosphere, the stratosphere, and clouds and
precipitation. Results from atmospheric modeling (plume and global), laboratory studies, and initial field studies will inform

260 priorities and research directions for Earth system studies and could potentially inform the design of field studies to verify
model results.

The potential broader impacts of ISA deployment beyond removing methane are not well understood. These may include co-
benefits such as reductions in atmospheric black carbon (Li et al., 2021, Oeste et al., 2017) and ozone (Li et al., 2023; van
265 Herpen et al., 2023), both of which are climate warming agents. However, there are also potential negative impacts that must
be further explored including ocean acidification, stratospheric ozone loss, adverse chemical side reactions (such as COCl₂
formation), and albedo changes from deposition on ice surfaces (Li et al., 2023). Iron aerosols absorb light and thus tend to
warm the planet, offsetting some of the lowered radiative forcing from oxidized methane (Matsui et al., 2018; Li et al., 2021;
Meidan et al., submitted). There are also potential side effects such as indirect radiative forcing due to marine cloud brightening
270 and carbon dioxide absorption by ocean iron fertilization (Emerson, 2019) that could be either favorable or unfavorable.

The potential effects of ISA on air quality and human health are also poorly understood. Enhanced chlorine could lead to
beneficial reductions in tropospheric ozone. However, the reduced hydroxyl radical production may increase lifetimes of
volatile organic compounds that are detrimental to human health. Moreover, iron aerosols themselves present a human health
275 risk, especially when small (O'Day et al., 2022). Therefore, more research is required to determine under which conditions
(including deployment location/timing, particle composition, and ISA size) the air quality impact is beneficial or detrimental.

Engineering deployment modalities and implementation scenarios is one of the later steps in this roadmap, only to be pursued
if earlier dependencies are addressed and ISA proves effective and climate beneficial with acceptable side effects.
280 Nevertheless, to avoid delaying potential future deployment-readiness and to iteratively refine design in advance of any *in-situ*
field studies, development and engineering of a nozzle sprayer delivery system could begin in parallel with early research
activities.

As part of a lifecycle analysis, the cost of materials (e.g., iron), infrastructure, and other implementation resources must be
285 assessed. To be cost-plausible, these costs at scale must be lower than the social cost of methane, a monetary valuation that
estimates the socioeconomic impact caused by an additional metric ton of methane (Azar et al., 2023).

3.2.3 Milestone question #3: What is needed to advance ethical governance and social license?

Policy and social science aspects of this roadmap must occur in parallel with the research and technical activities and should
be iterative, where outcomes from policy and social science efforts inform development of research and technical direction,
290 and vice versa. A misaligned or siloed approach may result in outcomes where physical science results are not answering the
social science questions to understand if there could be a safe, cost-plausible deployment method and pathway to ethical

governance, which may stall or halt progress. Conversely, detrimental outcomes could also be realized if research and technical aspects were delayed.

295 It is imperative that governance and social impacts be considered for any atmospheric methane removal approach. Addressing
the climate crisis requires engagement beyond technical solutions; collaboration and transparency between scientists,
behavioral scholars, media, the public, policy-makers, NGOs, Indigenous peoples, and other stakeholders is essential to ensure
that an ethical governance framework is established (Dowell et al., 2020; Diamond et al., 2022; Data for Progress, 2023;
Carbon180, 2022). There must be effective engagement and education to co-create research questions and iteratively
300 communicate research findings and results, as well as risks and co-benefits. Failure to do so jeopardizes the trust and sound
decision making of communities and governments (The Arctic Institute, 2021), threatening our ability to critically and openly
assess potential climate solution approaches through a scientific process. Ideally, an external governance framework is
developed which is enforceable and legally binding; however, there is also value in internal governance frameworks which
may, for example, be based around a code of conduct, advisory or review boards, or other non-binding structures.

305 If ISA remains promising, potential implementation scenarios could be considered. Some methane removal solutions may be
deployed immediately, whereas others may be higher risk and may only be suitable for emergency deployment.

Category	Research and development activity	Year in which work becomes a funding priority	Anticipated duration (variable depending on outcomes)	Academia	Non-profits	Government	Engineering consultant
Key milestone question #1: Is enhancement of the chlorine oxidative sink of methane via the iron salt aerosol (ISA) mechanism effective and climate beneficial? At what scale?							
Modeling studies	Atmospheric plume and high-resolution regional modeling to understand if ISA-driven chlorine radical production results in net methane loss using various ISA sizes and concentrations of iron, NO _x , and other chemical species.	underway	3 years	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	


	Global atmospheric modeling to understand if ISA-driven chlorine radical production results in net cooling, considering radiative forcing from methane, ozone, and iron aerosols, the photoactivity of the iron, meteorological and atmospheric chemistry conditions, and ISA size.	underway	4 years	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
	Paleo modeling to understand variability in the chlorine-based methane sink that may have been associated with past dust changes.	2024	2 years	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
Laboratory studies	Chamber studies to understand the photoactive iron fraction, chlorine radical production efficiencies, methane oxidation efficiencies, and reactions with other species	underway	3 years	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
Field studies	Natural analogue studies to (a) better constrain the atmospheric radical sinks; (b) understand the spatial extent and magnitude of the natural analogue, MDSA; and (c) understand the sources and distribution of photoactive iron (from MDSA as well as combustion iron sources)	underway	3 years	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
	<i>In-situ</i> ISA enhancement field studies (e.g. ship plume and/or ground based) at a controlled experiment site to measure changes in chlorine, methane, and other species	2025	2+ years	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
Engineering	Initial design study for ISA sprayer system.	2024	1 year	<input checked="" type="checkbox"/>			<input checked="" type="checkbox"/>
Key milestone question #2: What impacts could the ISA approach have on Earth systems and human systems, both positive and negative? Is there a cost-plausible deployment method?							
Modeling studies	Modeling of ocean system impacts considering different release locations/timing, ISA sizes, and atmospheric conditions.	2024	3+ years	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	

	Modeling of ISA deposition impacts on different land and ice surfaces considering different release locations/timing, ISA sizes, and atmospheric conditions.	2024	3+ years	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
	Modeling of atmospheric impacts including clouds, precipitation, and stratospheric ozone considering different release locations/timing, ISA sizes, and atmospheric conditions.	2024	3+ years	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
	Lifecycle analysis to quantify ISA climate benefits, tradeoffs, and cost.	2024	Ongoing	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
Governance and social impacts	Study of potential human health impacts considering different release locations/timing, ISA sizes, and atmospheric conditions.	2025	3 years	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
Engineering	Advanced design study for ISA sprayer system.	2026	1 year	<input checked="" type="checkbox"/>			<input checked="" type="checkbox"/>
	Design study of deployment modalities and implementation scenarios.	2027	2 years	<input checked="" type="checkbox"/>			<input checked="" type="checkbox"/>
Key milestone #3: What is needed to advance a structure of ethical governance and social license for utilizing atmospheric intervention to reduce atmospheric methane concentrations?							
Governance and social impacts	Engage with stakeholders regarding the research findings and results, risks, and potential.	underway	Ongoing	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
	Develop a collaborative governance framework to monitor and report on environmental and social impacts.	2024	Ongoing		<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
	Stakeholder engagement for identification of potential deployment locations and strategies.	2026	3 years			<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>

310 **Table 1: Key research and development activities for a five-year timeline beginning in 2023. Funding is needed for all research and development activities, including those that are already underway.**

3.3 Priorities and timeline

It may be advantageous to pursue multiple research questions in parallel because the output from one research question may inform the inputs for other research questions. As such, activities can be sequenced using a prioritized timeline (Figure 1), where later research activities and action areas often have multiple dependencies on earlier activities. For example, the engineering design study is suggested to begin in year 4 (2027) because it can start prior to having complete Earth system modeling results, human health study outcomes, or conclusions from ISA enhancement field studies. However, the engineering design study cannot advance to its later stages until earlier activities have been thoroughly addressed. This expedited schedule advances possible timelines to avoid delaying potential deployment-readiness, but is not meant to accelerate the timeline beyond appropriate caution and due diligence. Setbacks in addressing early research questions and action areas will likely result in timeline delays.



325 **Figure 1: Roadmap of ISA research and development needs. Duration of research and development timelines may be longer or shorter than depicted and/or exceed the five-year horizon.**

4 Conclusion

The activities outlined in this roadmap require coordinated efforts across multiple government agencies to financially support research and development, to ensure robust assessment and governance processes, and to foster international engagement. This work is valuable for multiple reasons:

1. We need to understand if ISA is a feasible, scalable, and safe methane removal method, or if it is nonviable and further research should be deprioritized.
2. Though this roadmap is ISA-specific, the research and development needs identified here contribute to fundamental understanding of processes and mechanisms that are broadly applicable to exploration of other methane removal approaches.
3. The research outlined in this roadmap will contribute to constraining the global methane budget and oxidative character of the atmosphere, which will improve our understanding of atmospheric chemistry, Earth system dynamics, and air quality.

Addressing the climate crisis requires a diverse portfolio of climate solutions. It is essential that atmospheric methane removal approaches are considered in addition to, not replacing, crucial anthropogenic greenhouse gas emission reductions and carbon dioxide removal. All atmospheric methane removal approaches are at a very early stage (Jackson et al., 2021; Ming et al., 2022; Spark, 2023); all require further research and none are ready for deployment. We hope that this ISA roadmap, and other atmospheric methane removal roadmaps that follow, will help accelerate, prioritize, and parallelize research that is essential to understanding which climate solutions to pursue.

References

- Aschwanden, A., Bartholomäus, T. C., Brinkerhoff, D. J., and Truffer, M.: Brief communication: A roadmap towards credible projections of ice sheet contribution to sea level, *The Cryosphere*, 15, 5705–5715, <https://doi.org/10.5194/tc-15-5705-2021>, 2021.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and IUPAC Subcommittee: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II – gas phase reactions of organic species, *Atmos. Chem. Phys.*, 6, 3625–4055, <https://doi.org/10.5194/acp-6-3625-2006>, 2006.
- Azar, C., Martin, J.G., Johansson, D.J.A., and Sterner, T.: The social cost of methane, *Climatic Change*, 176, <https://doi.org/10.1007/s10584-023-03540-1>, 2023.

- 360 Bock, M., Schmitt, J., Beck, J., Seth, B., Chappellaz, J., and Fischer, H.: Glacial/interglacial wetland, biomass burning, and geologic methane emissions constrained by dual stable isotopic CH₄ ice core records, *P. Natl. Acad. Sci. USA*, 114, E5778-E5786, <https://doi.org/10.1073/pnas.1613883114>, 2017.
- Bondy, A.L., Wang, B., Laskin, A., Craig, R.L., Nhliziyo, M.V., Bertman, S.B., Pratt, K.A., Shepson, P.B., and Ault, A.P.: Inland Sea Spray Aerosol Transport and Incomplete Chloride Depletion: Varying Degrees of Reactive Processing Observed during SOAS, *Environ. Sci. Technol.*, 51, 9533-9542, <https://doi.org/10.1021/acs.est.7b02085>, 2017.
- 365 Carbon180, Setting DAC on Track: Strategies for Hub Implementation: <https://static1.squarespace.com/static/5b9362d89d5abb8c51d474f8/t/6261d1890b76863f1047a2dd/1650577901659/Carbon180-SettingDAConTrack.pdf>, last access: 7 December 2023, 2022.
- 370 Chen, Y., Cheng, Y., Ma, N., Wei, C., Ran, L., Wolke, R., Größ, J., Wang, Q., Pozzer, A., Denier van der Gon, H. A. C., Spindler, G., Lelieveld, J., Tegen, I., Su, H., and Wiedensohler, A.: Natural sea-salt emissions moderate the climate forcing of anthropogenic nitrate, *Atmos. Chem. Phys.*, 20, 771–786, <https://doi.org/10.5194/acp-20-771-2020>, 2020.
- Data for Progress, Advancing Equitable Deployment of Regional DAC Hubs: <https://www.filesforprogress.org/memos/advancing-equitable-deployment-of-regional-dac-hubs.pdf>, last access: 7 December 2023, 2023.
- 375 Dean, J.F., Middelburg, J.J., Röckmann, T., Aerts, R., Blauw, L.G., Egger, M., Jetten, M.S.M., de Jong, A.E.E., Meisel, O.H., Rasigraf, O., Slomp, C.P., in't Zandt, M.H., and Dolman, A.J.: Methane Feedback to the Global Climate System in a Warmer World, *Rev. Geophys.*, 56, 207-250, <https://doi.org/10.1002/2017RG000559>, 2018.
- 385 Diamond, M.S., Gettelman, A., Lebsock, M.D., McComiskey, A., Russell, L.M., Wood, R., and Feingold, G.: To assess marine cloud brightening's technical feasibility, we need to know what to study - and when to stop, *P. Natl. Acad. Sci. USA*, 119, <https://doi.org/10.1073/pnas.2118379119>, 2022.
- Dowell, G., Niederdeppe, J., Vanucchi, J., Dogan, T., Donaghy, K., Jacobson, R., Mahowald, N., Milstein, M., and Zelikova, T.J.: Rooting carbon dioxide removal research in the social sciences, *Interface Focus*, 10, <https://doi.org/10.1098/rsfs.2019.0138>, 2020.
- 390 Emerson, D.: Biogenic Iron Dust: A Novel Approach to Ocean Iron Fertilization as a Means of Large Scale Removal of Carbon Dioxide From the Atmosphere, *Front. Mar. Sci.*, 6, <https://doi.org/10.3389/fmars.2019.00022>, 2019.

Fischer, H., Behrens, M., Bock, M., Richter, U., Schmitt, J., Loulergue, L., Chappellaz, J., Spahni, R., Blunier, T., Leuenberger, M., and Stocker, T.F.: Changing boreal methane sources and constant biomass burning during the last termination, *Nature*, 395 452, 864-867, <https://doi.org/10.1038/nature06825>, 2008.

Fischer, H., Siggaard-Andersen, M.L., Ruth, U., Rothlisberger, R., and Wolff, E.: Glacial/interglacial changes in mineral dust and sea-salt records in polar ice cores: sources, transport, and deposition. *Rev. Geophys.*, 45, <https://doi.org/10.1029/2005RG000192>, 2007.

Ferretti, D.F., Miller, J.B., White, J.W., Etheridge, D.M., Lassey, K.R., Lowe, D.C., Meure, C.M., Dreier, M.F., Trudinger, C.M., Van Ommen, T.D., and Langenfelds, R.L.: Unexpected changes to the global methane budget over the past 2000 years, *Science*, 309, 1714-1717, <https://doi.org/10.1126/science.1115193>, 2005.

405 Forster, P., T. Storelvmo, K. Armour, W. Collins, J.-L. Dufresne, D. Frame, D.J. Lunt, T. Mauritsen, M.D. Palmer, M. Watanabe, M. Wild, and H. Zhang: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity, *Climate Change 2021: The Physical Science Basis, Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, 410 J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 923–1054, doi:10.1017/9781009157896.009, 2021.

Gromov, S.S.: Stable isotope composition of atmospheric carbon monoxide: a modeling study: <https://openscience.ub.uni-mainz.de/handle/20.500.12030/4250>, last access: 7 December 2023, 2014.

415 Han, C., Do Hur, S., Han, Y., Lee, K., Hong, S., Erhardt, T., Fischer, H., Svensson, A.M., Steffensen, J.P., and Vallelonga, P.: High-resolution isotopic evidence for a potential Saharan provenance of Greenland glacial dust, *Sci. Rep.*, 8, <https://doi.org/10.1038/s41598-018-33859-0>, 2018.

420 Horowitz, H.M., Holmes, C., Wright, A., Sherwen, T., Wang, X., Evans, M., Huang, J., Jaegle, L., Chen, Q., Zhai, S., and Alexander, B.: Effects of sea salt aerosol emissions for marine cloud brightening on atmospheric chemistry: Implications for radiative forcing, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2019GL085838>, 2020.

IPCC, AR6 Synthesis Report: Climate Change 2023: <https://www.ipcc.ch/report/sixth-assessment-report-cycle/>, last access 7 425 December 2023, 2023.

- Ito, A.: Global modeling study of potentially bioavailable iron input from shipboard aerosol sources to the ocean, *Global Biogeochem. Cy.*, 27, <https://doi.org/10.1029/2012GB004378>, 2013.
- 430 Jackson, R.B., Abernethy, S., Canadell, J.G., Cargnello, M., Davis, S.J., Feron, S., Fuss, S., Heyer, A.J., Hong, C., Jones, C.D., Matthews, H.D., O’Conner, F.M., Pisciotta, M., Rhoda, H.M., de Richter, R., Solomon, E.I., Wilcox, J.L., and Zickfeld, K.: Atmospheric methane removal: a research agenda, *Phil. Trans. R. Soc. A*, 379, <https://doi.org/10.1098/rsta.2020.0454>, 2021.
- Jackson, R.B., Saunio, M., Bousquet, P., Canadell, J.G., Poulter, B., Stavert, A.R., Bergamaschi, P., Niwa, Y., Segers, A.,
435 and Tsuruta, A., Increasing anthropogenic methane emissions arise equally from agricultural and fossil fuel sources, *Environ. Res. Lett.*, 15, <https://doi.org/10.1088/1748-9326/ab9ed2>, 2020.
- Kleinen, T., Gromov, S., Steil, B., and Brovkin, V.: Atmospheric methane underestimated in future climate projections. *Environ. Res. Lett.*, 16, <https://doi.org/10.1088/1748-9326/ac1814>, 2021.
- 440
- Legrand, M., Preunkert, S., Wolff, E., Weller, R., Jourdain, B., and Wagenbach, D.: Year-round records of bulk and size-segregated aerosol composition in central Antarctica (Concordia site) – Part 1: Fractionation of sea-salt particles, *Atmos. Chem. Phys.*, 17, 14039–14054, <https://doi.org/10.5194/acp-17-14039-2017>, 2017.
- 445 Lelieveld, J., Peters, W., Dentener, F.J., and Krol, M.C.: Stability of tropospheric hydroxyl chemistry, *J. Geophys. Res.*, 107, <https://doi.org/10.1029/2002JD002272>, 2002.
- Lenton, T.M., Held, H., Kriegler, E., Hall, J.W., Lucht, W., Rahmstorf, S., and Schellnhuber, H.J.: Tipping elements in the Earth’s climate system, *P. Natl. Acad. Sci. USA*, 105, 1786-1793, <https://doi.org/10.1073/pnas.0705414105>, 2008.
- 450
- Li, Q., Fernandez, R.P., Hossaini, R., Iglesias-Suarez, F., Cuevas, C.A., Apel, E.C., Kinnison, D.E., Lamarque, J.F., and Saiz-Lopez, A.: Reactive halogens increase the global methane lifetime and radiative forcing in the 21st century, *Nat. Commun.*, 13, <https://doi.org/10.1038/s41467-022-30456-8>, 2022.
- 455 Li, L., Mahowald, N. M., Miller, R. L., Pérez García-Pando, C., Klose, M., Hamilton, D. S., Gonçalves Ageitos, M., Ginoux, P., Balkanski, Y., Green, R. O., Kalashnikova, O., Kok, J. F., Obiso, V., Paynter, D., and Thompson, D. R.: Quantifying the range of the dust direct radiative effect due to source mineralogy uncertainty, *Atmos. Chem. Phys.*, 21, 3973–4005, <https://doi.org/10.5194/acp-21-3973-2021>, 2021.

- 460 Li, Q., Meidan, D., Hess, P., Anel, J.A., Cuevas, C.A., Doney, S., Fernandez, R.P., van Herpen, M., Hoglund-Isaksson, L., Johnson, M.S., Kinnison, D.E., Lamarque, J.F., Röckmann, T., Mahowald, N.M., and Saiz-Lopez, A.: Global environmental implications of atmospheric methane removal through chlorine-mediated chemistry-climate interactions, *Nat. Commun.*, 14, <https://doi.org/10.1038/s41467-023-39794-7>, 2023.
- 465 Li, Y., Song, Y., Li, X., Kaskaoutis, D.G., Gholami, H., and Li, Y.: Disentangling variations of dust concentration in Greenland ice cores over the last glaciation: An overview of current knowledge and new initiative, *Earth-Science Reviews*, 242, <https://doi.org/10.1016/j.earscirev.2023.104451>, 2023.
- Maesano, C.N., Campbell, J.S., Foteinis, S., Furey, V., Hawrot, O., Pike, D., Aeschlimann, S., Reginato, P.L., Goodwin, D.R.,
470 Looger, L.L., Boyden, E.S., and Renforth, P.: Geochemical Negative Emissions Technologies: Part II. Roadmap, *Front. Clim.*, 4, <https://doi.org/10.3389/fclim.2022.945332>, 2022.
- Mahowald, N.M., Hamilton, D.S., Mackey, K.R.M., Moore, J.K., Baker, A.R., Scanza, R.A., and Zhang, Y.: Aerosol trace metal leaching and impacts on marine microorganisms, *Nat. Commun.*, 9, <https://doi.org/10.1038/s41467-018-04970-7>, 2018.
475
- Mak, J., Kra, G., Sandomenico, T., and Bergamaschi, P.: The seasonality varying isotopic composition of the sources of carbon monoxide at Barbados, West Indies, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2003JD003419>, 2003.
- Matsui, H., Mahowald, N.M., Moteki, N., Hamilton, D.S., Ohata, S., Yoshida, A., Koike, M., Scanza, R.A., and Flanner, M.G.:
480 Anthropogenic combustion iron as a complex climate forcer, *Nat. Commun.*, 9, <https://doi.org/10.1038/s41467-018-03997-0>, 2018.
- McKay, D.I.A., Stall, A., Abrams, J.F., Winkelmann, R., Sakschewski, B., Loriani, S., Fetzer, I., Cornell, S.E., Rockstrom, J.,
485 and Lenton, T.M.: Exceeding 1.5°C global warming could trigger multiple climate tipping points, *Science*, 377, <https://doi.org/10.1126/science.abn7950>, 2022.
- Meidan, D.: Evaluating the potential of iron-based interventions in methane reduction and climate mitigation, submitted to *Environ. Res. Lett.*, <https://doi.org/10.17632/v6mmbm75wd.1>.
- 490 Ming, T., Li, W., Yuan, Q., Davies, P., de Richter, R., Peng, C., Deng, Q., Yuan, Y., Caillol, S., and Zhou, N.: Perspectives on removal of atmospheric methane., *Advances in Applied Energy*, 5, <https://doi.org/10.1016/j.adapen.2022.100085>, 2022.

Neumann, R.B., Moorberg, C.J., Lundquist, J.D., Turner, J.C., Waldrop, M.P., McFarland, J.W., Euskirchen, E.S., Edgar, C.W., and Turetsky, M.R.: Warming Effects of Spring Rainfall Increase Methane Emissions From Thawing Permafrost, 495 *Geophys. Res. Lett.*, 46, 1393-1401, <https://doi.org/10.1029/2018GL081274>, 2019.

Nisbet, E.G., Manning, M.R., Dlugokencky, E.J., Michel, S.E., Lan, X., Röckmann, T., Denier van der Gon, H.A.C., Schmitt, J., Palmer, P.I., Dyonisius, M.N., Oh, Y., Fisher, R.E., Lowry, D., France, J.L., White, J.W.C., Brailsford, G., and Bromley, T.: Atmospheric Methane: Comparison Between Methane's Record in 2006-2022 and During Glacial Terminations, Global 500 *Biogeochem. Cy.*, 37, <https://doi.org/10.1029/2023GB007875>, 2023.

NOAA Global Monitoring Laboratory, Earth System Research Laboratories, Trends in Atmospheric Methane: https://gml.noaa.gov/ccgg/trends_ch4/, last access: 7 December 2023.

505 Ocean Visions, Ocean-Based Carbon Dioxide Removal: Road Maps: <https://www2.oceanvisions.org/roadmaps/>, last access: 7 December 2023.

O'Day, P.A., Pattammattel, A., Aronstein, P., Leppert, V.J., Forman, H.J.: Iron Speciation in Respirable Particulate Matter and Implications for Human Health, *Environ. Sci. and Technol.*, 56, 7006-7016, <https://doi.org/10.1021/acs.est.1c06962>, 2022. 510

Oeste, F. D., Method for cooling the troposphere: <https://patentimages.storage.googleapis.com/ac/64/72/49f3c79762e44f/AU2010203265B2.pdf>, last access: 7 December 2023, 2009.

515 Oeste, F. D., de Richter, R., Ming, T., and Caillol, S.: Climate engineering by mimicking natural dust climate control: the iron salt aerosol method, *Earth Syst. Dynam.*, 8, 1–54, <https://doi.org/10.5194/esd-8-1-2017>, 2017

Paudel, R., Mahowald, N.M., Hess, P.G.M., Meng, L., and Riley, W.J.: Attribution of changes in global wetland methane emissions from pre-industrial to present using CLM4.5-BGC, *Environ. Res. Lett.*, 11, <https://doi.org/10.1088/1748-9326/11/3/034020>, 2016. 520

Pennacchio, L., van Herpen, M., Meidan, D., Saiz-Lopez, A., and Johnson, M.S.: Catalytic efficiencies for atmospheric methane removal in the high-chlorine regime, submitted to *Environ. Res. Lett.*, <https://doi.org/10.26434/chemrxiv-2023-3r8sf>.

- 525 Peng, S., Lin, X., Thompson, R.L., Xi, Y., Liu, G., Hauglustaine, D., Lan, X., Poulter, B., Ramonet, M., Saunois, M., Yin, Y., Zhang, Z., Zheng, B., and Ciais, P.: Wetland emission and atmospheric sink changes explain methane growth in 2020, *Nature*, 612, 477–482, <https://doi.org/10.1038/s41586-022-05447-w>, 2022.
- Pio, C.A., and Lopes, D.A.: Chlorine loss from marine aerosol in a coastal atmosphere, *J. Geophys. Res.*, 103, 25,263-25,272, <https://doi.org/10.1029/98JD02088>, 1998.
- 530 Prospero, J.M., Delany, A.C., Delany A.C., and Carlson, T.N.: The Discovery of African Dust Transport to the Western Hemisphere and the Saharan Air Layer, *B. Am. Meteorol. Soc.*, 102, E1239-E1260, <https://doi.org/10.1175/BAMS-D-19-0309.1>, 2021.
- 535 Pye, H. O. T., Nenes, A., Alexander, B., Ault, A. P., Barth, M. C., Clegg, S. L., Collett Jr., J. L., Fahey, K. M., Hennigan, C. J., Herrmann, H., Kanakidou, M., Kelly, J. T., Ku, I.-T., McNeill, V. F., Riemer, N., Schaefer, T., Shi, G., Tilgner, A., Walker, J. T., Wang, T., Weber, R., Xing, J., Zaveri, R. A., and Zuend, A.: The acidity of atmospheric particles and clouds, *Atmos. Chem. Phys.*, 20, 4809–4888, <https://doi.org/10.5194/acp-20-4809-2020>, 2020.
- 540 Röckmann, T., Brenninkmeijer, C.A.M., Crutzen, P.J., and Platt, U.: Short term variations in the $^{13}\text{C}/^{12}\text{C}$ ratio of CO as a measure of Cl activation during tropospheric ozone depletion events in the Arctic, *J. Geophys. Res.*, 104, 1691–1697, <https://doi.org/10.1029/1998JD100020>, 1999.
- 545 Rodriguez, S., Prospero, J.M., Lopez-Darias, J., Garcia-Alvarez, M.I., Zuidema, P., Nava, S., Lucarelli, F., Gaston, C.J., Galindo, L., and Sosa, E.: Tracking the changes of iron solubility and air pollutants traces as African dust transits the Atlantic in the Saharan dust outbreaks, *Atmos. Environ.*, 246, <https://doi.org/10.1016/j.atmosenv.2020.118092>, 2021.
- 550 Saiz-Lopez, A., Fernandez, R.P., Li, Q., Cuevas, C.A., Fu, X., Kinnison, D.E., Times, S., Mahajan, A.S., Martin, J.C.G., Iglesias-Suerez, F., Hossaini, R., Plane, J.M.C., Myhre, G., and Lamarque, J.F.: Natural short-lived halogens exert an indirect cooling effect on climate, *Nature*, 618, 967-973, <https://doi.org/10.1038/s41586-023-06119-z>, 2023.
- Sapart, C.J., Monteil, G., Prokopiou, M., van de Wal, R.S.W., Kaplan, J.O., Sperlich, P., Krumhardt, K.M., Van der Veen, C., Houweling, S., Krol, M.C., Blunier, T., Sowers, T., Martinerie, P., Witrant, E., Dahl-Jensen, D., and Röckmann, T.: Natural and anthropogenic variations in methane sources during the past two millennia, *Nature*, 490, 85-88, <https://doi.org/10.1038/nature11461>, 2012.

Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguia-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth Syst. Sci. Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.

570

Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*. John Wiley & Sons, ISBN 0471720186, 2016.

Spark Climate Solutions, *Approaches to Atmospheric Methane Removal*: <https://www.sparkclimate.org/methane-removal/101/approaches>, last access: 7 December 2023.

580

Staniaszek, Z., Griffiths, P.T., Folberth, G.A., O'Connor, F.M., Abraham, N.L., and Archibald, A.T.: The role of future anthropogenic methane emission in air quality and climate. *NPJ Clim. Atmos. Sci*, 5, <https://doi.org/10.1038/s41612-022-00247-5>, 2022.

The Arctic Institute, *Sami Council resistance to SCoPEX highlights the complex questions surrounding geoengineering and consent*: <https://www.thearcticinstitute.org/sami-council-resistance-scopex-highlights-complex-questions-geoengineering-consent/>, last access: 7 December 2023, 2021.

585 FAO, Food and Agriculture Organization of the United Nations, *Free Prior and Informed Consent: An indigenous peoples' right and a good practice for local communities*: <https://www.fao.org/3/i6190e/i6190e.pdf>, last access: 7 December 2023, 2016.

UNEP, United Nations Environment Programme, *Global Methane Assessment: Benefits and Costs of Mitigating Methane Emissions*: <https://www.unep.org/resources/report/global-methane-assessment-benefits-and-costs-mitigating-methane-emissions>, last access: 7 December 2023, 2021.

590

- van Herpen, M.M.J.W., Li, Q., Saiz-Lopez, A., Liisberg, J.B., Röckmann, T., Cuevas, C.A., Fernandez, R.P., Mak, J.E., Mahowald, N.M., Hess, P., Meidan, D., Stuut, J.-B.W., and Johnson, M.S.: Photocatalytic Chlorine Atom Production on Mineral Dust-Sea Spray Aerosols over North Atlantic, *P. Natl. Acad. Sci. USA*, 120, 595 <https://doi.org/10.1073/pnas.2303974120>, 2023.
- Wang, X., Jacob, D. J., Eastham, S. D., Sulprizio, M. P., Zhu, L., Chen, Q., Alexander, B., Sherwen, T., Evans, M. J., Lee, B. H., Haskins, J. D., Lopez-Hilfiker, F. D., Thornton, J. A., Huey, G. L., and Liao, H.: The role of chlorine in global tropospheric chemistry, *Atmos. Chem. Phys.*, 19, 3981–4003, <https://doi.org/10.5194/acp-19-3981-2019>, 2019.
- 600 Wanser, K., Doherty, S.J., Hurrell, J.W., and Wong, A.: Near-term climate risks and sunlight reflection modification: a roadmap approach for physical science research, *Climatic Change*, 174, <https://doi.org/10.1007/s10584-022-03446-4>, 2022.
- Wittmer, J., Bleicher, S., Ofner, J., and Zetzsch, C.: Iron (III)-induced activation of chloride from artificial sea-salt aerosol. *Environ. Chem.*, 12, 461-475, <https://doi.org/10.1071/EN14279>, 2015a.
- 605 Wittmer, J., Bleicher, S., and Zetzsch, C.: Iron (III)-induced activation of chloride and bromide from modeled salt pans, *J. Phys. Chem. A*, 119, 19, 4373-4385, <https://doi.org/10.1021/jp508006s>, 2015b.
- 610 Wittmer, J., and Zetzsch, C.: Photochemical activation of chlorine by iron oxide aerosol. *J. Atmos. Chem.*, 74, 187-204, <https://doi.org/10.1007/s10874-016-9336-6>, 2017.
- Yuan, T., Yu, H., Chin, M., Remer, L.A., McGee, D., and Evan, A.: Anthropogenic decline of African dust: Insights from the Holocene records and beyond. *Geophys. Res. Lett.*, 47, 22, <https://doi.org/10.1029/2020GL089711>, 2020.
- 615 Zhang, Z., Poulter, B., Feldman, A.F., Ying, Q., Ciais, P., Peng, S., and Li, X.: Recent intensification of wetland methane feedback, *Nat. Clim. Change*, 13, 430-433, <https://doi.org/10.1038/s41558-023-01629-0>, 2023.
- Zhu, X. R., Prospero, J. M., and Millero, F. J.: Diel variability of soluble Fe(II) and soluble total Fe in North African dust in the trade winds at Barbados, *J. Geophys. Res.*, 102, 21297– 21305, <https://doi.org/10.1029/97JD01313>, 1997.
- 620