

1 **Factors that shape the elevational patterns of plant diversity in the Yatsugatake**
2 **Mountains, Japan**

3
4
5 **Yoshitaka Oishi***

6
7 **ORCID:** <https://orcid.org/0000-0002-7837-8746>

8
9 Fukui Prefectural University, Centre for Arts and Sciences, Fukui 910-1195, Japan

10
11 *oishiy@fpu.ac.jp

ABSTRACT

Elevation is involved in determining plant diversity in montane ecosystems. This study examined whether the species distribution of plants in the Yatsugatake Mountains, central Japan, substantiated hypotheses associated with an elevational diversity gradient. Species richness of trees, shrubs, herbs, ferns, and bryophytes was investigated in study plots established at 200-m elevational intervals from 1800 to 2800 m. The changes in plant diversity (alpha and beta diversities, plant functional types, and elevational ranges) with elevation were analyzed in relation to climatic factors and elevational diversity gradient hypotheses, that is, mass effect, mid-domain effect, and Rapoport's elevational rule. A comparison of alpha and beta diversities revealed that different plant groups respond variably to elevation; the alpha diversity of trees and ferns decreased, that of herbs increased, whereas the alpha diversity of shrubs and bryophytes showed a U-shaped relationship and a hump-shaped pattern. The beta diversity of shrubs, herbs, and bryophytes increased above the subalpine–alpine transition ecotone. In accordance with these changes, the dominance of evergreen shrubs and graminoids increased above this ecotone, whereas that of evergreen trees and liverworts decreased. None of the plant groups showed a wide elevational range at higher elevations. These elevational patterns of plant groups were explained by climatic factors, and not by elevational diversity gradient hypotheses. These patterns were further influenced by plant–plant interactions via competition for light availability and physical habitat alternation.

Keywords: climatic factor, elevational pattern, mass effect, mid-domain effect, plant–plant interaction, Rapoport's elevational rule

1. INTRODUCTION

Elevation is a key factor for determining the patterns of plant diversity in mountainous areas (Jiang, Ma, Liu, & Tang, 2018; Körner, 2007; Lee, Chun, Song, & Cho, 2013; Lee & La Roi, 1979). The patterns of plant diversity are often explained by several hypotheses for elevational diversity gradient such as mass effect, mid-domain effect, and Rapoport's elevational rule. According to the mass effect hypothesis, a species disperses from source areas occupied by larger populations to adjacent sink areas; however, the dispersed species cannot maintain viable populations in the sink areas (Grytnes, Heegaard, & Ihlen, 2006; Shmida & Wilson, 1985). These temporal populations increase species richness around ecotones where two different communities meet. Unlike this hypothesis, the mid-domain effect hypothesis assumes a random

distribution of species within a geometrically constrained domain such as that between the summit and base of a mountain. Consequently, species richness peaks in the middle of the domain because of the increasing overlap of species distribution toward the center (Colwell & Lees, 2000; Colwell, Rahbek, & Gotelli, 2004). Rapoport's elevational rule postulates that species at higher elevations can withstand a broad range of climatic conditions across a high range of elevations (Stevens, 1992).

However, explanation of plant elevational patterns using these hypotheses is often debatable. This is because plant groups show disparate responses to elevation, and even plants within a group present dissimilar elevational distribution in different areas (Bhattarai & Vetaas, 2003; Bruun et al., 2006; Grytnes, Heegaard, & Ihlen, 2006; Miyajima, Sato, & Takahashi, 2007; Sánchez-González & López-Mata, 2005). Hence, the applicability and limitations of these hypotheses for elevational diversity gradient should be further explored through careful examination of relationships between plant distribution and elevation.

Such an examination in mountainous areas in Japan is of interest because of the characteristic vegetation at higher elevations. These areas experience strong winds and heavy snowfall owing to the influence of jet streams and winter monsoons (Manabe, 1957; Riehl, 1962; Ueda, Kibe, Saitoh, & Inoue, 2015). The severe climatic conditions enable a single Japanese stone pine tree species (*Pinus pumila* Regel) to become the dominant species above the subalpine–alpine transition zone owing to its resistance to wind and snow (Okitsu, 1984). This dominance could alter plant distribution from that predicted based on the hypothesis of elevational diversity distribution. This is because the subalpine–alpine transition zone is otherwise dominated by a mixture of alpine and subalpine plants according to the mass effect. From other perspectives, the dominance of *P. pumila* could reduce the randomness of species distribution as postulated by the mid-domain effect and hinder the dispersal of alpine species into lower elevations against the assumption based on the Rapoport's elevational rule.

In this study, the elevational distribution of multiple plant groups in the Yatsugatake Mountains, Japan, was evaluated to determine whether the hypotheses for elevational diversity gradient (i.e., mass effect, mid-domain effect, and Rapoport's elevational rule) could substantiate the existing elevational patterns of the plant groups. If these hypotheses could explain the elevational patterns of plants, plant diversity can exhibit the following changes with elevation.

Hypothesis 1: If the elevational patterns of plants follow the mass effect hypothesis, the alpha diversity will be the highest around the subalpine and alpine ecotones.

Furthermore, beta diversity can be the highest in these ecotones owing to the coexistence of subalpine and alpine plants.

Hypothesis 2: If the mid-domain effect hypothesis is applied to the elevational patterns, the alpha diversity will be the highest around the middle elevations between the summit and the base of a mountain. Beta diversity can be uniform across the elevations based on the nature of random distribution.

Hypothesis 3: If the distribution of plant species can be explained by the Rapoport's elevational rule, alpine species will exhibit larger elevational ranges than species at lower elevations. Hence, the alpha diversity will be relatively high at lower elevations where both alpine and lowland species coexist.

Hypothesis 4: If the three elevational hypotheses do not apply to the observed patterns of plant distribution, other factors, such as climate and plant–plant interactions, will largely determine their distribution. Specifically, the influence of plant–plant interactions may be presented in relationships between the ecological traits of plant groups (plant functional types).

2. MATERIALS AND METHODS

2.1 Study site

This study was conducted in the Yatsugatake Mountains. The highest peak of these mountains is the 2899 m summit of Mount Akadake (35°58.259'N, 138°22.223'E). The vegetation in the Yatsugatake Mountains is largely classified into the following four types: temperate deciduous forest dominated by *Quercus crispula* and *Larix kaempferi* below ~1800 m; subalpine conifer forest dominated by *Tsuga diversifolia*, *Abies veitchii*, and *Abies mariesii* f. *hayachinensis* at ~1800–2600 m; alpine dwarf pine scrub dominated by *P. pumila* at ~2600–2800 m; and alpine meadow with turf-rock vegetation at ~2800–2900 m. The highest and lowest mean temperatures in the nearest weather station at 1350 m are 19.2°C in August and -5.3°C in January, and the mean precipitation is 1439.9 mm/year (Japan Meteorological Agency, 2018).

2.2 Field survey

Twelve 10 × 10 m study plots were established at 200 m elevational intervals from 1800 to 2800 m, extending from the east to the west of the Yatsugatake Mountains. This plot size was determined using the minimal area method in areas with uniform microtopography. The presence of plant species in each plot was determined based on the taxonomic groups: trees, shrubs, herbs, ferns, and bryophytes. Shrubs are defined as woody plant species of height less than 2.0 m. Ferns and bryophytes included both

epiphytic and terrestrial species. Epiphytic species were recorded up to 2.0 m from the ground.

To examine the influence of climate on plant diversity, temperature and relative humidity (RH) were measured in each plot at ~5 cm above the ground at 4 h intervals throughout the year, obtained using HOBO U23 Pro v2 temperature/relative humidity data loggers (Onset Computer Corporation, Bourne, MA, USA). To reduce the influence of sunlight on the measurements, the data loggers were covered with a CO-RS1 solar radiation shield (Onset Computer Corporation). Using the temperature and RH data, the following climatic variables were calculated: the mean annual temperature ($Temp_{ann}$), mean temperature during the growing season ($Temp_{grow}$), mean temperature during the snow season ($Temp_{snow}$), mean annual RH (RH_{ann}), mean RH during the growing season (RH_{grow}), mean RH during the snow season (RH_{snow}), and duration of the snow season (snow cover). A series of snow-covered days with over 90% RH was defined as the snow season. This definition was based on a primary experiment in which data loggers were buried under a snow pack. The remaining days were defined as the growing season. RH_{snow} was not used in the following analysis because this variable is nearly uniform according to its definition.

2.3 Statistical analyses

Alpha diversity, beta diversity, dominance of functional types, and elevational range of each plant group were calculated to examine the validity of the hypotheses for elevational diversity gradient. Alpha diversity was defined as the species richness of each plant group within a plot, and its correlation with the elevation was fitted by generalized linear models (GLMs) with log-link function and Poisson distribution. Models were selected as follows (model selection 1). Both linear and second-order polynomial models were constructed, and the final model with significant coefficients and lower Akaike information criterion (AIC) values was selected. If a model did not have significant coefficients, then the model with the lowest AIC value was adopted.

The influence of climatic factors on alpha diversity was also analyzed using GLMs with log-link function and Poisson distribution. The final model was selected as follows (model selection 2). Climatic factors that strongly correlated with alpha diversity ($r > 0.5$) were selected as explanatory variables. The models with different combinations of these explanatory variables were then constructed. Among these models, the final model had significant coefficients and lower AIC values.

Beta diversity was evaluated using the β_{sim} index, considering that the performance of this metric was excellent under various criteria (Koleff, Gaston, & Lennon, 2003). The β_{sim} index was calculated between the adjacent plots as follows:

$$\beta_{sim} = 1 - \frac{a}{(\min[b, c] + a)} \quad (1)$$

where, a is the total number of species found in both plots, b is the number of species that occur in the first plot but not in the second plot, and c is the number of species that occur in the second plot but not in the first plot. The changes in beta diversity with elevation were fitted by linear models (LMs). The final model was selected according to model selection 1. The influences of climatic factors on beta diversity were analyzed using LMs. The final model was determined based on model selection 2.

The elevational patterns of plants were further discussed in relation to the changes in plant functional types. Plant functional types are groups of species that respond similarly to environmental and biotic changes (Duckworth, Kent, & Ramsay, 2000). These responses were used to characterize the elevational patterns of plant groups (Bruun et al., 2006; Sánchez-González & López-Mata, 2005; Zhou et al., 2019). In this study, trees, shrubs, herbs, and bryophytes were classified into the following functional types: evergreen and deciduous trees, evergreen and deciduous shrubs, forbs and graminoids, and mosses and liverworts, respectively. The dominance of plant functional groups was calculated as the ratio of the species richness of each functional type to the total species richness of the entire plant assemblage in each plot.

The changes in the dominance of plant functional types with elevation were fitted by LMs. Elevation was used as an explanatory variable and the final model was selected according to model selection 1. The relationships between the plant functional types were examined using Pearson correlation.

To test the applicability of the mid-domain effect to elevational distribution, the discrepancy between the observed results and those predicted based on the mid-domain effect was determined using the null model (discrete mid-domain effect model) (Colwell & Hurtt, 1994). The predicted mean species richness was calculated by 9999 Monte Carlo simulations with RangeModel software version 5.0 (Colwell, 2006). The predicted values were plotted against elevation and correlated with the observed alpha diversity using Pearson's correlation.

Finally, the applicability of the Rapoport's elevational rule to elevational distribution was examined based on the elevational range of each plant group. The elevational range was defined as the mean difference between the highest and lowest elevations of all the recorded species in each plot and was calculated as follows:

$$\text{Elevational range} = \frac{\sum_{i=1}^n (H_i - L_i)}{n} \quad (2)$$

where, n is the total number of species recorded in a plot, H_i is the highest elevation where the i th species was recorded, and L_i is the lowest elevation where the i th species was recorded. The highest and lowest elevations were based on elevation records in this study site and herbarium specimens (Nagoya University Museum; NUM-Bt) collected in central Japan (Chubu, Kinki, and Kanto regions), plant diversity databases (Biodiversity Centre of Japan, 2018), and plant survey reports from the studied region (Akiyama, 1983, 1984; Hattori, 1958; Inoue, 1981; Kodama, 1971, 1972; Masuzaki & Katagiri, 2010; Takaki, Amakawa, Osada, & Sakuma, 1970). The changes in elevational ranges with elevation were examined using LMs. This modeling was performed with the same procedure for LMs between beta diversity and elevation.

3. RESULTS

3.1 Species richness, elevational distribution, and climatic factors

Two hundred and sixty plant species (20 tree species, 17 shrubs, 49 herbs, 11 ferns, and 163 bryophytes) were recorded in the study area. The species richness per plot (mean \pm standard deviation) of trees, shrubs, herbs, ferns, and bryophytes was 5.00 ± 1.81 , 2.50 ± 2.32 , 9.42 ± 5.14 , 2.00 ± 1.86 , and 41.00 ± 17.00 , respectively (Appendix S1).

The changes in climatic factors with elevations are presented in Appendix S2. The Pearson's correlation analysis between climatic factors and elevation showed that elevation was positively correlated with the snow cover ($r = 0.769$; $p < 0.01$) but negatively correlated with Temp_{ann} ($r = -0.862$; $p < 0.01$) and Temp_{grow} ($r = -0.822$; $p < 0.01$). In contrast, no significant correlations were observed between elevation and the RH variables (RH_{ann} and RH_{grow}).

3.2 Alpha diversity

The alpha diversity of the plant groups varied with elevational distribution (Figure 1). All coefficients, except those for ferns, in the GLMs were significant (Appendix S3). The alpha diversity of both total plants and trees decreased with elevation (Figure 1a, 1b), whereas that of herbs increased (Figure 1d). The alpha diversity of bryophytes showed a hump-shaped pattern (Figure 1f) and that of shrubs showed a U-shaped relationship with the highest value at an elevation of 2800 m (Figure 1c). The alpha diversity of herbs increased with elevation; however, the highest alpha diversity on an average was recorded in the 2400-m plot.

The results of alpha diversity and climatic factors obtained using GLMs are shown in Table 1. The variables Temp_{ann} and Temp_{snow} presented significant positive effects on the alpha diversity of trees and ferns. The alpha diversity of shrubs negatively correlated with the RH_{grow}, whereas that of herbs positively correlated with snow cover and that of bryophytes positively correlated with Temp_{ann} and RH_{grow}.

3.3 Mid-domain effect

Species richness (alpha diversity) was predicted using a discrete mid-domain effect model to examine the applicability of this hypothesis to the sampling site (Figure 1, Appendix S4). In all plant groups, the correlations between the observed species richness and those predicted based on the mid-domain effect were not positively significant (Table 2).

3.4 Beta diversity

When beta diversity (based on the B_{sim} index) was plotted against elevation (Figure 2, Appendix S5), the β_{sim} index of shrubs and ferns was not calculated in several plots because of the absence of these plant groups (Figure 2c, 2e). The β_{sim} of shrubs and herbs substantially increased and that of bryophytes slightly increased above the subalpine–alpine transition zone (Figure 2c, 2d, 2f).

The results of LMs between the β_{sim} index and environmental factors are presented in Table 3. Similar to the results of alpha diversity, snow cover and RH_{grow} had a significant influence on the beta diversity of herbs and bryophytes. No LMs with significant coefficients were constructed for the beta diversity of trees and ferns.

3.5 Functional types

Elevational patterns of dominance of plant functional types are shown in Figure 3 and Appendix S6. The changes in the dominance of plant functional types were notable around 2600–2800 m (Figure 3). The dominance of evergreen trees and liverworts decreased in this elevation range (Figure 3a, 3d), whereas that of evergreen shrubs and graminoids increased (Figure 3b, 3c).

Regarding the relationships between functional types, the dominance of evergreen trees had a negative influence on that of evergreen shrubs and graminoids whereas it positively affected the dominance of liverworts (Table 4).

3.6 Elevational range

The mean elevational range of each plant group was calculated and plotted as a function of elevation (Figure 4, Appendix S7). None of the plant groups presented a wider elevational range at higher elevations.

4. DISCUSSION

In this study, elevational distribution of multiple plant groups was investigated in the Yatsugatake Mountains to determine whether the hypotheses for elevational diversity distribution could substantiate the existing elevational patterns of plant groups. The plant groups showed different responses to the elevational gradients and climatic factors in the study site. Regarding the elevational changes in alpha diversity, only shrubs presented higher alpha diversity in the subalpine–alpine ecotone than that predicted based on the mass effect. However, none of the plant groups exhibited elevational distribution patterns corresponding to those predicted based on the mid-domain effect or Rapoport’s elevational rule. In contrast, the alpha diversity of all plant groups was explained by the climatic factors. The dominance of plant functional types correlated with each other.

4.1 Elevational patterns of plants

The alpha diversity of trees decreased with elevation (Figure 1b), and this elevational pattern was explained by $Temp_{ann}$. These results are supported by the findings of a previous study that the elevational range of trees can be determined by temperature, which strongly affects their physiochemical activities (Körner, 2003). This obvious influence of $Temp_{ann}$ on tree diversity may have led to the relatively constant changes in the beta diversity compared with that of other plant groups (Figure 2a) owing to a linear decrease in $Temp_{ann}$ with elevation.

Similar to the results for trees, the alpha diversity of ferns decreased with elevation (Figure 1e), which can be attributed to $Temp_{snow}$ in the GLM. These results are consistent with the findings of previous reports, that is, fern richness decreased at higher elevation and lower temperature in central Japan (Tanaka & Sato, 2013, 2014).

In contrast to the results for the trees and ferns, the alpha diversity of herbs positively correlated with elevation (Figure 1d). Given that snow cover positively affected herb diversity in the GLM, the species richness of herbs increased with the environmental changes related to snow cover, such as soil moisture and canopy closure. Long periods of snow augment soil moisture (Hardy et al., 2001) essential for the proliferation of herbs in arctic–alpine habitats (Litaor, Williams, & Seastedt, 2008; Nabe–Nielsen et al., 2017; Roux, Aalto, & Luoto, 2013). On the contrary, snow cover

decreases canopy closure by suppressing tree canopy growth (Song, Hogan, Brown, Cao, & Yang, 2017). This decrease improves light intensity and reduces litter accumulation on the forest floor. These changes in soil moisture and canopy closure enhance herb richness by guaranteeing the photosynthesis and germination of herbs.

The alpha diversity of shrubs rapidly increased from the subalpine–alpine to the alpine zone (Figure 1c). This elevational pattern was explained by the RH_{grow} in the GLM. Considering that canopy closure reduces shrub diversity (Speziale, Ruggiero, & Ezcurra, 2010) and intensifies air humidity (Jung et al., 2017), RH_{grow} may be selected in the GLM as a surrogate to represent the negative influence of canopy closure on shrub richness. Accordingly, this change in canopy closure from dense to open can be attributed to the rapid increase in the beta diversity of shrubs above the subalpine–alpine transition zone (Figure 2c).

Only the bryophytes showed a hump-shaped change in alpha diversity with an increase in elevation (Figure 1f). The GLM results demonstrated that this pattern could be mainly attributed to RH_{grow} . This relationship is consistent with the results of a previous study, which suggested that water stress is a major determinant of bryophyte diversity (Grau, Grytnes, & Birks, 2007). The negative correlation between RH_{grow} and bryophyte beta diversity implies that water stress facilitates species turnover via the disappearance of drought-sensitive species.

4.2 Functional types and plant–plant interactions

Here, the elevational patterns of plant diversity are further discussed in relation to plant functional types. Evergreen shrub species (*P. pumila*) have started to dominate the subalpine–alpine ecotone instead of evergreen conifer trees. This replacement has caused a reduction in canopy closure above the ecotone, as indicated by the negative correlation between the dominance of evergreen trees and evergreen shrubs (Table 4).

In accordance with the decrease in evergreen trees, the dominance of graminoids increased but that of liverworts decreased (Figure 3c, 3d). These correlations can be attributed to improved light availability and increased drought stress owing to the change from a closed to an open canopy. Graminoids dominate open forests (Roberts & Zhu, 2002; Thomas, Halpern, Falk, Liguori, & Austin, 1999) and are more resistant to drought stress than forbs (Rosbakh et al., 2017), whereas liverworts are more shaded adapted (Marschall & Proctor, 2004) and are more vulnerable to drought stress than mosses (Grau Grytnes, & Birks, 2007; Jan & Wolf, 1993). These plant–plant interactions such as competition for light availability and alternation of physical habitat impose a certain influence on elevational patterns of plants shaped by climatic factors.

These interactions facilitate species turnover around subalpine–alpine ecotones, increasing the beta diversity of shrubs, herbs, and bryophytes.

4.3 Mass effect on elevational patterns

None of the plant groups, except shrubs, presented the highest alpha diversity around the subalpine–alpine ecotone (Figure 1) and all the elevational patterns, including those of shrubs, could be explained by climatic factors and plant–plant interactions (section 4.1 and 4.2). These results contradict the expected hump-shaped pattern predicted based on the mass effect (Hypothesis 1). Specifically, plant–plant interactions might strongly affect these unexpected patterns because the dominance of stone pine trees inhibits the establishment of small plants by shading and curbs the increase in species richness in the subalpine–alpine ecotone.

4.4 Mid-domain effect and the Rapoport’s elevational rule on elevational patterns

The alpha diversity of all plant groups differed from that predicted based on the mid-domain effect hypothesis (Figure 1, Table 2). Furthermore, none of the plant groups showed a wider elevational range in the alpine area against that predicted using the Rapoport’s elevational rule (Figure 4). The lack of applicability of the mid-domain effect and Rapoport’s rule to elevational patterns can be attributed to the simple assumptions of these hypotheses; the former excludes any effects of gradients on expected patterns (Colwell, Rahbek, & Gotelli, 2004), whereas the latter fails when species distribution is affected by the interaction of several factors (Bhattarai & Vetaas, 2006; Grau, Grytnes, & Birks, 2007). Climate and vegetation at higher elevations in this study site are characterized by severe winter climates and the resultant dominance of *P. pumila* above the subalpine–alpine ecotones. These characteristics affect the elevational distribution of plants, and thereby increase discrepancies between actual and predicted elevational patterns.

4.5 Determinants of plant elevational patterns

In summary, the elevational patterns of plant diversity in the Yatsugatake Mountains were explained by climate and plant–plant interactions, and not by hypotheses for elevational diversity distribution. Therefore, these results correspond to Hypothesis 4, indicating that the applicability of hypotheses for elevational diversity distribution largely depends on the environment and biota in the targeted study site. Given that elevational hypotheses are proposed based on simple assumptions such as random distribution of species and adaptation of alpine species to a broad range of

environments, the applicability is further reduced in areas with large differences in climatic conditions and plant–plant interactions within elevation ranges.

This study also emphasizes the importance of studies on elevational patterns using multiple plant groups and multiple indices of plant diversity (e.g., alpha and beta diversities, and functional types). These groups and indices are necessarily to understand the characteristics of plants’ responses to elevation and their interactions that influence elevational diversity distribution.

Acknowledgments

The author is grateful to Taichi Watanabe (Shinshu University) for his support in the field survey of vascular plants. The author acknowledges the financial support for this research received from the Japan Society for the Promotion of Science–Grant-in-Aid for Young Scientists (B) (grant number 24710029).

Competing Interest

The author declares no competing interests.

Author Contributions

Yoshitaka Oishi: Conceptualization, Methodology, Formal analysis, Investigation, Visualization, and Funding acquisition.

Data Accessibility Statement

The datasets used in this study are available in the Supporting information.

REFERENCES

- Akiyama, H. (1983). Bryophytes of Mt. Hakusan 1. Hepaticopsida & Anthocerotopsida. *Annual report of Hakusan Nature Conservation Center*, 9, 13–24.
- Akiyama, H. (1984). Bryophytes of Mt. Hakusan 2. Bryopsida. *Annual report of Hakusan Nature Conservation Center*, 11, 25–46.
- Bhattarai, K. R., & Vetaas, O. R. (2003). Variation in plant species richness of different life forms along a subtropical elevation gradient in the Himalayas, east Nepal.

410 *Global Ecology and Biogeography*, 12(4), 327–340.
 411 <https://doi.org/10.1046/j.1466-822X.2003.00044.x>

412 Bhattarai, K. R., & Vetaas, O. R. (2006). Can Rapoport's rule explain tree species
 413 richness along the Himalayan elevation gradient, Nepal? *Diversity and*
 414 *Distributions*, 12(4), 373–378. <https://doi.org/10.1111/j.1366-9516.2006.00244.x>

415 Biodiversity centre of Japan (2018). National vegetation survey database. Retrieved
 416 from http://gis.biodic.go.jp/webgis/files/veg_survey_db.zip

417 Bruun, H. H., Moen, J., Virtanen, R., Grytnes, J. A., Oksanen, L., & Angerbjörn, A.
 418 (2006). Effects of altitude and topography on species richness of vascular plants,
 419 bryophytes and lichens in alpine communities. *Journal of Vegetation Science*,
 420 17(1), 37–46. <https://doi.org/10.1111/j.1654-1103.2006.tb02421.x>

421 Colwell, R. K. (2006). RangeModel A Monte Carlo simulation tool for assessing
 422 geometric constraints on species richness. Version 5. Retrieved from
 423 <http://viceroy.eeb.uconn.edu/rangemodel>.

424 Colwell, R. K., & Hurtt, G. C. (1994). Nonbiological gradients in species richness and a
 425 spurious Rapoport effect. *The American Naturalist*, 144(4), 570–595.

426 Colwell, R. K., & Lees, D. C. (2000). The mid-domain effect: geometric constraints on
 427 the geography of species richness. *Trends in Ecology & Evolution*, 15(2), 70–76.
 428 [https://doi.org/10.1016/S0169-5347\(99\)01767-X](https://doi.org/10.1016/S0169-5347(99)01767-X)

429 Colwell, R. K., Rahbek, C., & Gotelli, Nicholas J. (2004). The mid-domain effect and
 430 species richness patterns: what have we learned so far? *The American Naturalist*,
 431 163(3), E1–E23. <https://doi.org/10.1086/382056>

432 Duckworth, J. C., Kent, M., & Ramsay, P. M. (2000). Plant functional types: an
 433 alternative to taxonomic plant community description in biogeography?
 434 *Progress in Physical Geography*, 24, 515–542.
 435 <https://doi.org/10.1177/030913330002400403>

436 Grau, O., Grytnes, J. A., & Birks, H. (2007). A comparison of altitudinal species
 437 richness patterns of bryophytes with other plant groups in Nepal, Central

438 Himalaya. *Journal of Biogeography*, 34(11), 1907–1915. [https://doi.org/10.1111/](https://doi.org/10.1111/j.1365-2699.2007.01745.x)
439 [j.1365-2699.2007.01745.x](https://doi.org/10.1111/j.1365-2699.2007.01745.x)

440 Grytnes, J. A., Heegaard, E., & Ihlen, P. G. (2006). Species richness of vascular plants,
441 bryophytes, and lichens along an altitudinal gradient in western Norway. *Acta*
442 *Oecologica*, 29(3), 241–246. <https://doi.org/10.1016/j.actao.2005.10.007>

443 Hardy, J. P., Groffman, P. M., Fitzhugh, R. D., Henry, K. S., Welman, A. T., Demers, J.
444 D., . . . Nolan, S. (2001). Snow depth manipulation and its influence on soil frost
445 and water dynamics in a northern hardwood forest. *Biogeochemistry*, 56(2),
446 151–174. <https://doi.org/10.1023/a:1013036803050>

447 Hattori, S. (1958). The hepaticae of Ontake Mountain, middle Japan. *Journal of Hattori*
448 *Botanical Laboratory*, 20, 33–53.

449 Inoue, H. (1981). Hepaticae of Mt. Fuji, central Japan. *Memoirs of the National Science*
450 *Museum*, 14, 59–74.

451 Jan, H., & Wolf, D. (1993). Diversity patterns and biomass of epiphytic bryophytes and
452 lichens along an altitudinal gradient in the Northern Andes. *Annals of the*
453 *Missouri Botanical Garden*, 80(4), 928–960. <https://doi.org/10.2307/2399938>

454 Japan Meteorological Agency. (2018). Past meteorological data (Nobeyama). Retrieved
455 from [http://www.data.jma.go.jp/obd/stats/etrn/view/nml_amd_ym.php?](http://www.data.jma.go.jp/obd/stats/etrn/view/nml_amd_ym.php?prec_no=48&block_no=0415&year=&month=&day=&view=)
456 [prec_no=48&block_no=0415&year=&month=&day=&view=](http://www.data.jma.go.jp/obd/stats/etrn/view/nml_amd_ym.php?prec_no=48&block_no=0415&year=&month=&day=&view=)

457 Jiang, Z., Ma, K., Liu, H., & Tang, Z. (2018). A trait-based approach reveals the
458 importance of biotic filter for elevational herb richness pattern. *Journal of*
459 *Biogeography*, 45(10), 2288–2298. <https://doi.org/10.1111/jbi.13398>

460 Jung, S. H., Lim, C. H., Kim, A. R., Woo, D. M., Kwon, H. J., Cho, Y. C., & Lee, C. S.
461 (2017). Edge effects confirmed at the clear-cut area of Korean red pine forest in
462 Uljin, eastern Korea. *Journal of Ecology and Environment*, 41(1), 36.
463 <https://doi.org/10.1186/s41610-017-0051-2>

- 464 Kodama, T. (1971). *Liverworts of Kinki District, central Japan, preserved in the Osaka*
 465 *Museum of Natural History. Part 1.* Osaka, Japan: Osaka Museum of Natural
 466 History (in Japanese).
- 467 Kodama, T. (1972). *Liverworts of Kinki District, central Japan, preserved in the Osaka*
 468 *Museum of Natural History. Part 2.* Osaka, Japan: Osaka Museum of Natural
 469 History (in Japanese).
- 470 Koleff, P., Gaston, K. J., & Lennon, J. J. (2003). Measuring beta diversity for presence–
 471 absence data. *Journal of Animal Ecology*, 72(3), 367–382.
 472 <https://doi.org/10.1046/j.1365-2656.2003.00710.x>
- 473 Körner, C. (2003). Carbon limitation in trees. *Journal of Ecology*, 91(1), 4–17.
 474 <https://doi.org/10.1046/j.1365-2745.2003.00742.x>
- 475 Körner, C. (2007). The use of ‘altitude’ in ecological research. *Trends in Ecology &*
 476 *Evolution*, 22(11), 569–574. <https://doi.org/10.1016/j.tree.2007.09.006>
- 477 Lee, C.-B., Chun, J.-H., Song, H.-K., & Cho, H.-J. (2013). Altitudinal patterns of plant
 478 species richness on the Baekdudaegan Mountains, South Korea: mid-domain
 479 effect, area, climate, and Rapoport’s rule. *Ecological Research*, 28(1), 67–79.
 480 <https://doi.org/10.1007/s11284-012-1001-1>
- 481 Lee, T. D., & La Roi, G. H. (1979). Bryophyte and understory vascular plant beta
 482 diversity in relation to moisture and elevation gradients. *Vegetation*, 40(1), 29–
 483 38.
- 484 Litaor, M. I., Williams, M., & Seastedt, T. R. (2008). Topographic controls on snow
 485 distribution, soil moisture, and species diversity of herbaceous alpine vegetation,
 486 Niwot Ridge, Colorado. *Journal of Geophysical Research: Biogeosciences*, 113,
 487 G02008. <https://doi.org/10.1029/2007JG000419>
- 488 Manabe, S. (1957). On the modification of air-mass over the Japan Sea when the
 489 outburst of cold air predominates. *Journal of the Meteorological Society of*
 490 *Japan. Ser. II*, 35(6), 311–326. https://doi.org/10.2151/jmsj1923.35.6_311

- 491 Marschall, M., & Proctor, M. C. F. (2004). Are bryophytes shade plants? Photosynthetic
492 light responses and proportions of chlorophyll a, chlorophyll b and total
493 carotenoids. *Annals of Botany*, 94(4), 593-603. <https://doi:10.1093/aob/mch178>
- 494 Masuzaki, H., & Katagiri, T. (2010). Liverworts on Mt. Kitadake & Mt. Ainodake. In T.
495 Masuzawa (Ed.), *Minami alps: geography and biota* (pp. 257–276). Shizuoka,
496 Japan: Shizuoka Prefecture (in Japanese).
- 497 Miyajima, Y., Sato, T., & Takahashi, K. (2007). Altitudinal changes in vegetation of
498 tree, herb and fern species on Mount Norikura, central Japan. *Vegetation*
499 *Science*, 24(1), 29–40. <https://doi.org/10.15031/vegsci.24.29>
- 500 Okitsu, S. (1984). Comparative studies on the Japanese alpine zone with special
501 reference to the ecology of *Pinus pumila* thickets. *Geographical Review of*
502 *Japan Ser. A*, 57(11), 791–802. https://doi.org/10.4157/grj1984a.57.11_791 (in
503 Japanese)
- 504 Riehl, H. (1962). *Jet streams of the atmosphere*. Fort Collins, Colorado: Colorado State
505 University.
- 506 Roberts, M. R., & Zhu, L. (2002). Early response of the herbaceous layer to harvesting
507 in a mixed coniferous–deciduous forest in New Brunswick, Canada. *Forest*
508 *Ecology and Management*, 155(1), 17–31. [https://doi.org/10.1016/S0378-](https://doi.org/10.1016/S0378-1127(01)00544-8)
509 [1127\(01\)00544-8](https://doi.org/10.1016/S0378-1127(01)00544-8)
- 510 Rosbakh, S., Leingärtner, A., Hoiss, B., Krauss, J., Steffan-Dewenter, I., & Poschlod, P.
511 (2017). Contrasting effects of extreme drought and snowmelt patterns on
512 mountain plants along an elevation gradient. *Frontiers in plant science*, 8, 1478–
513 1478. <https://doi:10.3389/fpls.2017.01478>
- 514 Roux, P. C., Aalto, J., & Luoto, M. (2013). Soil moisture's underestimated role in
515 climate change impact modelling in low–energy systems. *Global Change*
516 *Biology*, 19(10), 2965–2975. <https://doi.org/10.1111/gcb.12286>

517 Sánchez-González, A., & López-Mata, L. (2005). Plant species richness and diversity
 518 along an altitudinal gradient in the Sierra Nevada, Mexico. *Diversity and*
 519 *Distributions*, 11(6), 567–575. <https://doi.org/10.1111/j.1366-9516.2005.00186.x>
 520 Shmida, A., & Wilson, M. V. (1985). Biological determinants of species diversity.
 521 *Journal of Biogeography*, 1–20. <https://doi.org/10.2307/2845026>
 522 Song, X., Hogan, J. A., Brown, C., Cao, M., & Yang, J. (2017). Snow damage to the
 523 canopy facilitates alien weed invasion in a subtropical montane primary forest in
 524 southwestern China. *Forest Ecology and Management*, 391, 275–281.
 525 <https://doi.org/10.1016/j.foreco.2017.02.031>
 526 Speziale, K. L., Ruggiero, A., & Ezcurra, C. (2010). Plant species richness–environment
 527 relationships across the Subantarctic–Patagonian transition zone. *Journal of*
 528 *Biogeography*, 37(3), 449–464. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2699.2009.02213.x)
 529 [2699.2009.02213.x](https://doi.org/10.1111/j.1365-2699.2009.02213.x)
 530 Stevens, G. C. (1992). The elevational gradient in altitudinal range: an extension of
 531 Rapoport's latitudinal rule to altitude. *The American Naturalist*, 140(6), 893–
 532 911. <https://doi.org/10.1139/X10-156>
 533 Takaki, N., Amakawa, T., Osada, T., & Sakuma, E. (1970). bryophytes flora of Mt.
 534 kaikoma, Mt. Senjo and Mt. Kitadake (Southern Japan Alps). *Journal of Hattori*
 535 *Botanical Laboratory*, 33, 171–202.
 536 Tanaka, T., & Sato, T. (2013). Elevational patterns of fern species assemblages and
 537 richness in central Japan. *Plant Ecology*, 214(9), 1189–1197.
 538 <https://doi.org/10.1007/s11258-013-0243-5>
 539 Tanaka, T., & Sato, T. (2014). Species richness of seed plants and ferns along a
 540 temperate elevational gradient in central Japan. *Plant Ecology*, 215(11), 1299–
 541 1311. <https://doi.org/10.1007/s11258-014-0388-x>
 542 Thomas, S. C., Halpern, C. B., Falk, D. A., Liguori, D. A., & Austin, K. A. (1999). Plant
 543 diversity in managed forests: understory responses to thinning and fertilization.

Ecological Applications, 9(3), 864-879. <https://doi:10.1890/1051->

0761(1999)009[0864:Pdimfu]2.0.Co;2

Ueda, H., Kibe, A., Saitoh, M., & Inoue, T. (2015). Snowfall variations in Japan and its linkage with tropical forcing. *International Journal of Climatology*, 35(6), 991–998. <https://doi.org/10.1002/joc.4032>

Zhou, Y., Ochola, A. C., Njogu, A. W., Boru, B. H., Mwachala, G., Hu, G., . . . Wang, Q. (2019). The species richness pattern of vascular plants along a tropical elevational gradient and the test of elevational Rapoport's rule depend on different life-forms and phytogeographic affinities. *Ecology and Evolution*, 9(8), 4495–4503. <https://doi.org/10.1002/ece3.5027>

TABLE 1.

Generalized linear model results between alpha diversity and climate variable.

Res. var	Exp. var	Estimate	Std. Error	Z-value	P-value	R ²
Total	Temp _{ann}	0.133	4.58×10^{-2}	2.91	< 0.01	< 0.01
	Temp _{snow}	-5.73×10^{-2}	4.03×10^{-2}	-1.42	0.155	
	RH _{ann}	0.126	3.92×10^{-2}	3.22	< 0.01	
Trees	Temp _{ann}	0.206	9.48×10^{-2}	2.18	0.0296	0.779
	Intercept	-55.4	26.2	-2.11	0.0347	
Shrubs	RH _{grow}	-0.234	6.89×10^{-2}	-3.39	< 0.01	0.686
	Intercept	22.4	6.28	3.57	< 0.01	
Herbs	Snow cover	1.54×10^{-2}	4.91×10^{-3}	3.15	< 0.01	0.592
	Intercept	-0.414	0.863	-0.480	0.632	
Ferns	Temp _{snow}	0.360	0.164	2.20	0.0280	0.519
	Intercept	-97.0	44.5	-2.18	0.0294	
Bryophytes	Temp _{ann}	0.113	3.51×10^{-2}	3.20	< 0.01	0.977
	RH _{grow}	8.60×10^{-2}	2.01×10^{-2}	4.29	< 0.01	
	Intercept	-4.69	1.85	-2.54	0.0111	

575 Res. var = response variable, Exp. var = explanatory variable, Std. error = standard error, R² =
576 Nagelkerke's R², Temp_{ann} = mean annual temperature, Temp_{snow} = mean temperature during the snow
577 season, RH_{ann} = mean annual RH, RH_{grow} = mean RH during the growing season, and Snow cover =
578 the duration of the snow season

579

580

581

582

583

584

585

586

587

588

589 **TABLE 2.**

Pearson correlation coefficients between species richness and that predicted using a discrete mid-domain effect model

	Total	Trees	Shrubs	Herbs	Ferns	Bryophytes
Cor.	0.160	−0.031	−0.738**	−0.074	0.110	0.276

**·: p < 0.01, *·: < p< 0.05

Cor. = Pearson correlation coefficient

TABLE 3.

Linear model results between beta diversity and climate variable.

Res. var	Exp. var	Estimate	Std. Error	t-value	p-value	R ²
Total	RH _{ann}	-9.21×10^{-2}	2.67×10^{-2}	-3.45	0.011	0.666
	Snow cover	6.09×10^{-3}	1.82×10^{-3}	3.34	0.012	
	Intercept	8.27	2.54	3.26	0.014	
Trees	RH _{ann}	7.56×10^{-2}	5.07×10^{-2}	1.49	0.174	0.217
	Intercept	-6.95	4.88	-1.43	0.192	
Shrubs	Tem _{ann}	-0.496	0.126	-3.94	0.029	0.784
	Intercept	1.70	0.329	5.17	0.014	
Herbs	Temp _{grow}	-9.44×10^{-2}	6.78×10^{-2}	-1.39	0.213	0.941
	RH _{ann}	-0.191	2.34×10^{-2}	-8.15	< 0.01	
	Snow cover	1.29×10^{-2}	2.62×10^{-3}	4.91	< 0.01	
	Intercept	17.2	2.55	6.74	< 0.01	
Ferns	RH _{grow}	-8.71×10^{-2}	5.32×10^{-2}	-1.64	0.349	0.456
	Intercept	8.69	5.04	1.72	0.335	
Bryophytes	RH _{grow}	-5.32×10^{-2}	2.11×10^{-2}	-2.52	< 0.01	0.374
	Intercept	5.39	1.96	2.74	< 0.01	

Res. var = response variable, Exp. var = explanatory variable, Std. error = standard error, R² = adjusted R², Temp_{ann} = mean annual temperature, Temp_{grow} = mean temperature during the growing season, RH_{ann} = mean annual RH, RH_{grow} = mean RH during the growing season, and Snow cover = the duration of the snow season

TABLE 4.

Pearson correlation coefficients of dominance among plant functional types

	Eve. trees	Eve. shrubs	Graminoids	Liverworts
Eve. trees	1.000	−0.899**	−0.766**	0.596*
Eve. shrubs		1.000	0.639	−0.388
Graminoids			1.000	−0.499
Liverworts				1.000

** $: p < 0.01$, * $: p < 0.05$

Eve. trees; evergreen trees, Eve. shrubs; evergreen shrubs

FIGURE LEGENDS

Figure 1. Elevational changes in the alpha diversity of plant groups in the Yatsugatake Mountains, central Japan: (a) total, (b) trees, (c) shrubs, (d) herbs, (e) ferns, and (f) bryophytes. The calculated alpha diversity based on the mid-domain effect is denoted with an asterisk (*). To illustrate the alpha diversity, generalized linear models were used to explore the pertinent changes with elevation. When the coefficient was significant, a 95% confidence interval was calculated and colored.

Figure 2. Elevational changes in the beta diversity (β_{sim} indices) of plant groups in the Yatsugatake Mountains, central Japan: (a) total, (b) trees, (c) shrubs, (d) herbs, (e) ferns, and (f) bryophytes. To illustrate the beta diversity, linear models were used to explore the pertinent changes with elevation. When the coefficient was significant, a 95% confidence interval was calculated and colored.

Figure 3. Elevational changes in the dominance of plant functional types in the Yatsugatake Mountains, central Japan: (a) evergreen trees, (b) evergreen shrubs, (c) graminoids, and (d) liverworts. To illustrate the dominance of plant functional types, linear models were used to explore the pertinent changes with elevation. When the coefficient was significant, a 95% confidence interval was calculated and colored.

Figure 4. Elevational changes in the distribution range of plant groups in the Yatsugatake Mountains, central Japan: (a) total, (b) trees, (c) shrubs, (d) herbs, (e) ferns, and (f) bryophytes. To illustrate the elevational changes, linear models were used to explore the pertinent changes with elevation. When the coefficient was significant, a 95% confidence interval was calculated and colored.

