

Table 3 Differences in chemotherapy regimens

Chemotherapy agent ¹	Germany and Austria since 2016	Germany and Austria until 2016 ²			United Kingdom	France and Lausanne ³			Israel	Netherlands	Czech Republic (COG ARET0332)	Hospital Vall d'Hebron, Spain
	VEC	RB-A	RB-B	RB-C	JOE ⁴	EC	VCy	VCyHR	VEC	VEC	VEC	VEC
Vincristine in mg/m ²	1.5	1.5	1.5	1.5	1.5	-	1.5	1.5	1.5	2	1.5	1.5
Carboplatin in mg/m ²	560	-	300	300	600	800	-	-	560	560	560	560
Etoposide in mg/m ²	300	450	450	-	300	500	-	-	300	150	300	300
Cyclophosphamide in mg/m ²	-	1200	-	1200	-	-	1500	3000	-	-	-	-

¹Calculated per body surface area. Most regimens include calculation per body weight for body weight < 10-12 kg that are not displayed in this table.

²alternating cycles of RB-A, RB-B, RB-C

³ alternating cycles Standard risk: EC and VCy, High Risk: EC and VCyHR

⁴JOE and VEC stand for a chemotherapy containing Vincristin, Etoposide and Carboplatin