

Table 1. Basic characteristics of patients in the three groups ( ( $\bar{x} \pm s$ , n=30))

	<b>SAB</b>	<b>ESP</b>	<b>RIB</b>	<b>P</b>
<b>Age (years)</b>	52.10±11.50	54.73±13.60	56.06±11.15	0.439
<b>BMI ( kg/m<sup>2</sup> )</b>	23.52±3.12	23.40±3.05	23.71±4.03	0.104
<b>Procedure duration (min)</b>	100.96±29.30	97±21.77	104.8±17.33	0.436
<b>Duration of anesthesia (min)</b>	113.3±30.59	109.36±21.91	112.75±24.20	0.606
<b>ASA class I /II</b>	12 /18	13/17	14/16	0.271

Statistical tests: Pairwise comparisons of groups analyzed by one-way ANOVA were made using *post hoc* analyses and the Student–Newman–Keuls *Q*-test.

Table 2. Intraoperative anesthetic dosage and recovery time in the three groups ( $\bar{x} \pm s$ , n=30))

	<b>SAB</b>	<b>ESP</b>	<b>RIB</b>	<b>F</b>	<b>P</b>
<b>Remifentanyl (<math>\mu\text{g}</math>)</b>	315 $\pm$ 94.31	311 $\pm$ 80.61	309.33 $\pm$ 80.61	0.047	0.954
<b>Propofol (mg)</b>	195 $\pm$ 51.57	192.66 $\pm$ 39.21	190.66 $\pm$ 20.16	0.092	0.912
<b>Recovery time (min)</b>	13.56 $\pm$ 1.07	13.26 $\pm$ 1.11	13.06 $\pm$ 1.04	1.635	0.201

Statistical tests: Pairwise comparisons of groups analyzed by one-way ANOVA were made using *post hoc* analyses and the Student–Newman–Keuls *Q*-test as well as the chi-square test.

Table 3. Prevalence of adverse events in the three groups ( ( $\bar{x} \pm s$ , n=30))

	<b>SAB</b>	<b>ESP</b>	<b>RIB</b>	$\chi^2$	<b>P</b>
<b>Nausea (n/%)</b>	4 (13.33%)	3 (10%)	3 (10%)	0.225	0.894
<b>Vomiting (n/%)</b>	2 (6.67%)	1 (3.33%)	1 (3.33%)	0.523	0.770
<b>block-related complications (n/%)</b>	0	0	0	–	–

Statistical test: Chi-square test.