

1
2
3
4
5 **Characterizing Peridotite Xenoliths from Southern Vietnam: Insight into the Underlying**
6 **Lithospheric Mantle**
7

8 Kirby P Hobbs¹, Lynne J. Elkins¹, John C. Lassiter², Nguyen Hoang³, Caroline M. Burberry¹
9

10 ¹ University of Nebraska-Lincoln, Lincoln, NE, USA

11 ² Jackson School of Geoscience, University of Texas at Austin, Austin, TX, USA

12 ³ Institute of Geological Sciences, Vietnam Academy of Science and Technology (VAST),
13 Hanoi, Vietnam
14
15
16
17
18
19
20
21

22 For submission to *Geochemistry, Geophysics, Geosystems*

Abstract

Extrusion tectonics has been invoked to explain recent magmatism across Indochina (e.g., Hoang & Flower, 1998). Lithospheric mantle xenoliths sampled from two alkali basalts in south-central Vietnam, Pleiku and Xuan Loc, comprise fertile spinel peridotites. To better determine the origins of the Indochinese subcontinental lithospheric mantle (SCLM), including impacts of tectonic extrusion, we present major and trace elements, and $^{87}\text{Sr}/^{86}\text{Sr}$, $^{143}\text{Nd}/^{144}\text{Nd}$, $^{206}\text{Pb}/^{204}\text{Pb}$, $^{207}\text{Pb}/^{204}\text{Pb}$, and $^{208}\text{Pb}/^{204}\text{Pb}$ in xenolith mineral separates.

Most peridotites from Pleiku and Xuan Loc have fertile major element compositions, “depleted” and “spoon-shaped” rare earth element (REE) patterns, interpreted to record prior melt depletion followed by melt metasomatism, and depleted isotopic signatures (e.g., $^{87}\text{Sr}/^{86}\text{Sr} = 0.70238\text{--}0.70337$ and $^{143}\text{Nd}/^{144}\text{Nd} = 0.512921\text{--}0.514190$). A small group of refractory peridotites have “enriched” REE patterns suggesting more extensive metasomatism, and enriched isotope ratios ($^{87}\text{Sr}/^{86}\text{Sr} = 0.70405$ and $^{143}\text{Nd}/^{144}\text{Nd} = 0.512755\text{--}0.512800$). The presence of both fertile and refractory xenoliths records a heterogeneous SCLM beneath Vietnam. Based on geothermobarometry calculations, fertile xenoliths have equilibrium temperatures of 923-1,034 °C and pressures of 11.7-15.8 kbar, while refractory xenoliths have comparable temperatures of 923-1,006 °C, but lower pressures of 7.1-10.0 kbar, suggesting refractory rocks are dominantly present at shallower depths.

We suggest that the regional lithospheric mantle has experienced variable melt extraction, producing heterogeneous compositions. While we cannot rule out partial removal and replacement of the lithosphere, large-scale delamination is not necessary to explain observed characteristics. The entire SCLM was more recently metasomatized by melts resembling

Cenozoic basalts, suggesting recent asthenospheric melting has modified the SCLM by melt infiltration.

Key Points

Main point #1: Lithospheric mantle xenoliths in Vietnam record a heterogeneous SCLM beneath Indochina, containing fertile and refractory compositions

Main point #2: Refractory xenoliths overall derive from shallower depths than fertile xenoliths, suggesting a chemically stratified Indochinese SCLM

Main point #3: The SCLM was metasomatized by melts resembling Cenozoic basalts, suggesting one melt infiltration event that does not require delamination

1. Introduction

The diffuse igneous province of Indochina is a collision-adjacent, complex tectonic region with extensive basaltic magmatism (Hoang & Flower, 1998; Hoang et al., 1996). Vietnamese Cenozoic plateau basalts record two-stage eruptive cycles that initiated ~17 Ma, and consist of large volumes of tholeiitic followed by smaller quantities of alkaline basalt eruptions, several of which host crustal and lithospheric mantle xenoliths (e.g., Anh et al., 2021; Hoang & Flower, 1998; Hoang et al., 1996; Nguyen & Kil, 2019, 2020). Prior geochemical analysis of Vietnamese basalts has included major element and trace element compositions and $^{87}\text{Sr}/^{86}\text{Sr}$, $^{143}\text{Nd}/^{144}\text{Nd}$, $^{206}\text{Pb}/^{204}\text{Pb}$, $^{207}\text{Pb}/^{204}\text{Pb}$, and $^{208}\text{Pb}/^{204}\text{Pb}$ isotopes, and suggests that early-stage tholeiitic volcanism may record partial melt contributions from a subcontinental lithospheric mantle (SCLM) source, while later-stage volcanism derives from an asthenospheric source (Hoang & Flower, 1998; Hoang et al., 1996). Hoang et al. (2013) further suggested that some early-stage volcanism may have been moderately chemically contaminated by crustal assimilation. Traditional tectonic regimes associated with mantle upwelling and partial melting, such as regional extension, do not adequately explain the observed abundance (70,000 km²) of volcanic activity in Indochina (Hoang & Flower, 1998). Previous studies have instead proposed a role for extrusion tectonics, positing that the adjacent Himalayan collision extruded Southeast Asia eastward and caused accompanying local mantle upwelling beneath Indochina (Figure 1) (Flower et al., 1998; Hoang & Flower, 1998; Hoang et al., 1996, 2013; Jolivet et al., 2018). However, the origin and character of the proposed mantle upwelling and associated partial melting have not been clearly explained.

Lithospheric mantle xenoliths provide an opportunity to constrain the depth, temperature, and composition of the local SCLM, and thereby better understand part of the local partial

melting and melt transport environment beneath a continent (e.g., Byerly & Lassiter, 2012). Subcontinental lithospheric mantle is typically an ancient, cold layer, variably enriched in trace elements, that separates the continental crust from the underlying convecting asthenosphere (McDonough, 1990). Lithospheric mantle xenoliths are typically thought to represent either the ancient SCLM or, in some circumstances, younger, more recently emplaced asthenosphere beneath continents, and may track underlying magmatic processes such as partial melting, melt enrichment, and metasomatic events. For example, major and trace element concentrations in mantle xenolith mineral phases can be used to calculate the equilibration temperature and pressure history of a xenolith, which can in turn be used to infer the thickness and thermal evolution of the SCLM (Ballhaus et al., 1991; Brey & Kohler, 1990; Liang et al., 2013; Lierman & Ganguly, 2003; Putirka, 2008). Mantle xenoliths thus provide a “window” into the SCLM, and by characterizing the SCLM spatially and temporally, we can further constrain local lithospheric and mantle dynamics to inform more realistic tectonic and dynamical models.

To date, mantle xenoliths from Vietnam have not clearly constrained the regional tectonics and dynamics in Indochina. Recent studies in central Vietnam at the Pleiku and Dalat volcanic centers have suggested that the mantle beneath Indochina experienced both a prior melt depletion event and various chemical re-enrichment processes (Anh et al., 2021; Nguyen & Kil, 2019, 2020). Pleiku lithospheric mantle xenoliths from Nguyen and Kil (2019) have estimated two-pyroxene equilibrium temperatures ranging from 825-1131°C and may have experienced 1-20% fractional melting, followed by silicate melt metasomatism. Xenoliths from locations farther to the south in the Dalat volcanic center, including Ba Ria and Nui Nua, have likewise experienced low degrees of melt extraction (1-13%), equilibrated between 714-1211°C, and experienced varying degrees of metasomatism (Anh et al., 2021; Nguyen & Kil, 2020). Our

current study expands this xenolith dataset and further characterizes the local Vietnamese SCLM using additional major and trace element and $^{87}\text{Sr}/^{86}\text{Sr}$, $^{143}\text{Nd}/^{144}\text{Nd}$, $^{206}\text{Pb}/^{204}\text{Pb}$, $^{207}\text{Pb}/^{204}\text{Pb}$, and $^{208}\text{Pb}/^{204}\text{Pb}$ compositions, for additional xenoliths collected from the Pleiku and Xuan Loc (south-central Vietnam) volcanic centers, comprising a total of 25 new spinel peridotite samples (Table 1, Figure 1b). The three objectives of this study are to: (1) track magmatic processes in the local SCLM; (2) compare the SCLM sampled by xenoliths across two different sites in Vietnam, to better characterize its spatial variability (Pleiku and Xuan Loc); and (3) better constrain local mantle evolution beneath a tectonically extruding Indochina.

2. Background

2.1. Geologic Background

Indochina was rifted from Gondwana and subsequently sutured to Asia during the closure of a series of Tethyan ocean basins in the Permian and Triassic (Metcalf, 2013). Subsequent subduction-related magmatism occurred over Indochina during the Cretaceous as a result of northward subduction of the Tethyan seafloor, emplacing Cordilleran-type granitic batholiths (Gibbons et al., 2015; Shellnutt et al., 2013). The initial collision of Indian continental lithosphere with Eurasia occurred in the Miocene at ~50 Ma (Gibbons et al., 2015). The continued movement of the Indian block northward (~3000 km) after the initial continental collision led to the onset of extrusion tectonics in Asia by 35 Ma (Rohrmann et al., 2012; Royden et al., 2008). The extrusion and clockwise rotation of Indochina occurred as left-lateral movement along large-scale block-bounding, regional transform faults (Figure 1a) (the Ailao Shan-Red River Fault Zone and Mae Ping Fault Zone) from ~35 to ~17 Ma, when motion along the transform faults became right-lateral, coeval with the cessation of South China Sea/East

Vietnam Sea rifting (Li et al., 2015; Zhu et al., 2009). This reconfiguration event was accompanied by the onset of diffuse volcanic activity in Indochina at ~17 Ma, which has continued through the Holocene and peaked within the last 3 Ma (Hoang & Flower, 1998; Hoang et al., 1996). The dominant driving mechanism of anomalous Cenozoic basaltic magmatism across Indochina has alternatively been attributed to either ongoing, reorganized extrusion tectonics, or the onset of a new phase of regional continental extension (Cullen et al., 2010; Flower et al., 1998; Hoang & Flower, 1998). The continued extrusional tectonic scenario, accompanied by coupled mantle flow and local upwelling, was suggested as an explanation for the observed local sequences of 1) early-stage tholeiites, proposed to be partial melts of refractory SCLM due to the arrival of extruded, possibly upwelling asthenospheric mantle and associated lithospheric heating, followed by 2) alkali basalts, thought to be partial melts sourced from the decompressing asthenosphere.

The basaltic plateaus across central and southern Vietnam cover an area of 23,000 km² and have an estimated volume of ~8,000 km³ (Hoang & Flower, 1998). The Pleiku basaltic plateau (Figure 1b) covers an area of ~4,000 km² and is characterized by the two-stage eruptive cycle typical of most eruptive centers in southern Vietnam (Hoang & Flower, 1998; Hoang et al., 2013). The Xuan Loc plateau basalts cover a smaller area of ~2400 km² (Figure 1b) (Hoang et al., 2013). The earlier stage eruptive units at both localities consist of quartz and olivine tholeiites with relatively high SiO₂ (48-55 wt.%) and low FeO (8.0-10.5 wt.%) and have been dated at Pleiku from 6.5-3.4 Ma using ⁴⁰Ar/³⁹Ar methods (Hoang & Flower, 1998; Hoang et al., 1996, 2013). In both locations, a series of tholeiitic flows is overlain by smaller eruptions of predominantly alkali basalt, with relatively low SiO₂ (40-50 wt.%) and high FeO (9.0-14.5 wt.%) that have been dated at Pleiku from 2.4-0.2 Ma (Hoang & Flower, 1998; Hoang et al., 1996,

2013). Although the individual flows have not been measured in as much detail, Xuan Loc plateau basalts span a similar range of measured ages to the Pleiku lavas (5.0-0.3 Ma) (Hoang et al., 2013).

Alkali basalts from both volcanic plateaus host numerous mantle-derived xenoliths including spinel lherzolites, spinel harzburgites, wehrlites, websterites, and pyroxenites (Hoang & Flower, 1998; Hoang et al., 2013, Nguyen & Kil, 2019, 2020). Basalts from both localities have $^{206}\text{Pb}/^{204}\text{Pb}$, $^{207}\text{Pb}/^{204}\text{Pb}$, $^{208}\text{Pb}/^{204}\text{Pb}$, $^{87}\text{Sr}/^{86}\text{Sr}$, and ϵ_{Nd} compositions that extend from the Indian MORB field towards a more enriched composition that resembles the Enriched Mantle 2 (EM2) mantle component (e.g., Sims & Hart, 2006; Workman et al., 2004) (where “enriched” refers to elevated time-integrated incompatible element concentrations recorded as relatively high $^{206}\text{Pb}/^{204}\text{Pb}$, $^{207}\text{Pb}/^{204}\text{Pb}$, $^{208}\text{Pb}/^{204}\text{Pb}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and low ϵ_{Nd}), with slightly more enriched isotopic compositions in Pleiku basalts relative to Xuan Loc; tholeiitic basalts likewise exhibit more enriched isotopic compositions overall relative to the alkali basalts (Hoang et al., 1996; 2013).

2.2. Tracking magmatic processes in the SCLM

The compositions of mantle xenoliths provide insight into magmatic processes that have affected the local lithospheric mantle, such as the nature of prior melt extraction (e.g., the degree of melting), which depletes residual peridotites in incompatible elements (Frey & Green, 1974; Michael & Bonatti, 1985). Melting models can help to infer degrees of prior melting, given the measured incompatible element concentrations of the residual rock. To apply these methods to the study of residual rocks, Johnson et al. (1990) revised the basic melting equations of Gast (1968) and Shaw (1970) to describe melting as the change of element concentration in

clinopyroxene (cpx) during melting of mantle peridotites, because cpx hosts the highest concentrations of incompatible trace elements.

Metasomatism, the process of fluid (aqueous or carbonitic)-rock or silicate melt-rock reaction, can also affect incompatible element concentrations in lithospheric mantle rocks. Metasomatism is considered “modal” when the metasomatic agent introduces new phases (i.e., precipitates new minerals along reactive pathways) and “cryptic” when reactions are only recorded in trace element compositions (Dawson, 1984). Because metasomatism can partially or fully overprint previous magmatic events, however, simultaneously deciphering the characteristics (e.g., source, timing, and melt/fluid compositions) of both metasomatism and partial melting processes is potentially complex (Ionov, 2002). For instance, the trace element makeup of the metasomatic agent itself is a controlling factor over the resulting trace element patterns in the host rock. Highly variable and heterogeneous metasomatic enrichment of trace elements in local rocks can likewise be produced by chromatographic fractionation of elements around, e.g., interstitial fluid-filled veins in peridotites; counterintuitively, such chromatographic processes may preferentially re-enrich harzburgites over lherzolites in trace elements (Navon & Stolper, 1987; Toramaru & Fujii, 1986). From modelling results, Ionov et al. (2002) suggested that such chromatographic fractionation effects could cause wide ranges of trace element patterns in host rocks during a single metasomatic event, where some rocks develop trace element patterns primarily controlled by the composition of the metasomatic agent, while others are controlled by trace element fractionation between coexisting phases. Despite the described complexity, we can nonetheless interpret the trace element concentrations and radiogenic isotope compositions of lithospheric mantle xenoliths to better understand the magmatic history of the

SCLM using a combination of partial melting and temperature and pressure equilibration models, as explored further below.

3. Analytical Methods

3.1. Electron microprobe analysis

The analytical techniques for this study were conducted at the University of Texas at Austin. Major elements for spinel (sp), orthopyroxene (opx), cpx, and olivine (ol) were measured on mineral separates in epoxy mounts with a JEOL JXA-8200 electron microprobe analyzer (EPMA) with wavelength dispersive spectrometry (WDS). Generally, at least one core and rim measurement were analyzed per grain, and typically five grains were measured for each sample. Analytical spot size was a diameter of 2 μm with a beam voltage of 15 kV and an operating current of 50 nA. Elemental concentrations were acquired using $K\alpha$ peak signals with analyzing crystal detectors, assigned as follows: LiFH – Mn, Ti, Cr; LiF – Fe, Ni; PETH – Ca; TAP – Na, Mg, Al, Si. The on-peak count times were variable for elements in each mineral (40-120 s on-peak, with equal time spent measuring background signals) and correspond to expected element concentrations in a given mineral, with minor elements having longer count times than major elements. The off-peak correction method was linear for all elements in ol and opx (except Ti in ol, for which we used an exponential correction). Spinel was calibrated using the mean atomic number (MAN) background intensity calibration curve. For cpx, we conducted the measurements in two rounds, one using the linear off-peak correction method and the other using the MAN background intensity calibration curve. Natural and synthetic crystalline solids with known major element compositions were used as calibration standards, and the matrix corrections used were ZAF and $\phi(\rho Z)$ (Armstrong, 1988). Average calculated major element

concentrations have standard deviations broadly correlated with elemental concentration. Major elements with concentrations > 5 wt.% typically have uncertainties (1σ) < 2% (except for Al), with higher uncertainties ranging from 3-20% in elements with concentrations < 5 wt.%. These uncertainties incorporate both analytical error and intra-sample heterogeneity (Table S1). Such natural heterogeneity within a sample is recorded, for example, in systematic core-rim variations in clinopyroxene Al and Cr content, and in the MgO and FeO variations between grains for sample VN-2018-21-PL-4 (see below).

3.2. In-situ trace element analysis

Trace element measurements were performed for the same cpx and opx epoxy mounts prepared for major element analysis, using an Agilent 7500ce laser-ablation inductively-coupled plasma mass spectrometer (LA-ICP-MS) with a NWR193-FX laser system. The analytical spot size had a diameter of 150 μm , to increase the signal intensity of low trace element concentrations in opx, and overlapped with EPMA spot locations. Samples were pre-ablated, with a dwell time of 60 s and washout time of 30 s. The Si wt.% from the EPMA measurements was used as the internal standard, NIST 612 was used as the primary analytical standard, and NIST 610, BHVO-2G, and NIST 616 were used as secondary standards. Analysis of NIST 610 and BHVO2G for all elements was generally within 5% of accepted values (for NIST 610, [Tm] was < 10 %; for BHVO2G, [Na] and [Nd] were within <10% and [Ca] and [Pb] were <20%). Results of NIST 616 analysis for all elements were within 10 % of expected values, except for elements with concentrations near detection limits (i.e., Nd, Gd, Dy, Ho, and Er, which were within 20 % of expected values). [Ti] and [Yb] for NIST 616, however, did not agree with expected values within 20%. Analyzed NIST 610 concentrations for all measured trace elements

are precise within 4% (1σ standard error, with $n = 27$), with reproducibilities within 6% of expected certified concentrations for all elements, with a mean difference of 2%.

3.3. Sr-Nd-Pb isotope geochemistry

Strontium, Nd, and Pb isotopes were measured in cpx separates. After hand-picking, the cpx separates were leached using a 2 N HCl solution in an ultrasonic bath for 5 minutes to remove surface coatings. 100 mg of cpx mineral separates from each sample were then dissolved in 5 mL concentrated HF and 1 mL concentrated HNO₃ in an oven at 105°C. After the initial dissolution, samples were dried and then dissolved in 6 N HCl in an oven at 105°C. The samples were redissolved in HCl + HBr and passed through AG1-X8 anion exchange resin to separate Pb. The washes containing Sr and Nd were then dried down and redissolved in HNO₃. The washes were then passed through Sr-Spec resin to separate Sr followed by RE-Spec resin to separate the rare earth elements (REE). Finally, the rare earth portions were converted to a HCl solution and passed through LN-Spec resin to separate Nd from other REE (procedures after Lassiter et al., 2003).

Lead isotopes were measured using a Nu Plasma 3D multi-collector ICP-MS. Lead separates were diluted to concentrations of 10 ppb in 2% HNO₃ and doped with 2 ppb Tl for pseudo-internal standard normalization. ²⁰³Tl and ²⁰⁵Tl were used to correct for mass fractionation of Pb and Hg isotopes using an exponential law. NBS 981 was used for standard-sample-bracketing to correct for analytical drift using the accepted values of ²⁰⁶Pb/²⁰⁴Pb = 16.9405, ²⁰⁷Pb/²⁰⁴Pb = 15.4967, and ²⁰⁸Pb/²⁰⁴Pb = 36.7220. Standard-sample-bracketing was used to linearly interpolate and correct for analytical drift. BCR-2 and BHVO-2 were measured as unknowns to assess external reproducibility; our measured values are largely consistent with values reported by Baker et al. (2004), except for ²⁰⁶Pb/²⁰⁴Pb in BHVO-2, where we measured a

very minor difference in values from reported compositions (Table 2). Five samples with low Pb concentrations were additionally diluted to a concentration of 0.25 ppb and analyzed using a multi-Daly detector array using sample-standard bracketing with NBS 981 as the bracketing standard (Table 2).

Strontium and Nd isotopes were measured using a Triton thermal ionization mass spectrometer as metals precipitated on Re filaments. We measured NBS 987 $^{87}\text{Sr}/^{86}\text{Sr} = 0.701254 \pm 0.000009$ (2σ ; $n=33$). BCR-2 and BHVO-2 were measured as unknowns for external reproducibility and are within 0.000008 of accepted values (Table 2). We measured JNdi-1 $^{143}\text{Nd}/^{144}\text{Nd} = 0.512114 \pm 0.000013$ (2σ ; $n=15$). The cpx mineral separates analyzed generally contained 2.5-50 ng of Pb, 1,000-10,000 ng of Sr, and 100-1,000 ng of Nd, making the impact of procedural blanks (<130 pg Pb; <140 pg Sr; <200 pg Nd) negligible.

4. Results

4.1. Petrology of Vietnamese mantle xenoliths

Mantle peridotite xenoliths from Pleiku and Xuan Loc are classified as group-1 xenoliths based on the classification scheme by Frey and Prinz (1978) and consist of sp, cpx, opx, and ol (Figure S1). By definition, group-1 xenoliths are typically lherzolites, harzburgites, and dunites that contain Cr-rich, Al_2O_3 -poor and TiO_2 -poor sp and pyroxenes (Frey & Prinz, 1978). All samples from Pleiku for this study are spinel lherzolites; however, the Pleiku basaltic plateau has also produced harzburgite and dunite xenoliths (Hoang et al., 2013; Nguyen & Kil, 2019). Xuan Loc xenolith samples are more diverse, containing lherzolites, harzburgites, and dunites. Sample VN-2018-36-XL-1 is a dunite and only contains ol with minor cpx and sp. Samples have been further subdivided into two groups based on sp Cr# (molar Cr / (Al + Cr)) after methods from

Byerly and Lassiter (2012): type-F (fertile) samples have a sp Cr# <0.25 and type-R (refractory) samples have sp Cr# \geq 0.25, where the relative “fertility” of the peridotite refers to how readily and productively it can generate magma upon partial melting (Figure 3). Based on this Cr# definition, Pleiku samples from this study except for VN-2018-21-PL-4 are all type-F peridotites, while the more southerly Xuan Loc samples contain several samples that are both type-F and type-R peridotites. For comparison, xenoliths from other eruptive centers in the southern Dalat province (including Nui Nua and Ba Ria in the Dalat volcanic plateau) and from Pleiku include both type-R and type-F peridotites (Anh et al., 2021; Nguyen & Kil, 2019, 2020), indicating that both fertile and refractory types have been identified from both sampled regions.

4.2. Major element compositions

Major element data for ol, opx, cpx, and sp mineral separates from Pleiku and Xuan Loc peridotites are provided in Table S1. Generally, all samples exhibit low heterogeneity on the grain scale between core measurements, except the major element data for sample VN-2018-21-PL-4, which indicate that it is heterogeneous on a grain scale (e.g., FeO and MgO in ol and opx separates of < 32% and < 7% RSD, respectively). Clinopyroxene and opx display variations between rim and core measurements in all samples except VN-2018-21-PL-5 and VN-2018-36-XL-11. The core-to-rim variation is present across all localities and sample types (Figure 2). Clinopyroxene Al₂O₃ and Cr₂O₃ concentrations are lower in the rims by ~0.2 to 1.0 wt. % and ~0.1 to 0.2 wt. %, respectively. Similarly, opx grains have rims that are lower by ~0.1 to 0.6 wt. % in Al₂O₃ and by up to 0.1 wt. % in Cr₂O₃. Samples VN-2018-21-PL-1 and VN-2018-36-XL-14 also display rims with slightly higher CaO (0.75 wt. % and 0.94 wt. %, respectively). In light of this variation, we present average core and rim compositions for each mineral as separate data sets (Table S1). Averaged core measurements are expected to best preserve the compositions of

the peridotites prior to entrainment and exhumation and, thus, have been used here to characterize the SCLM.

The Mg# (molar Mg / (Mg + Fe²⁺)) of ol from Pleiku and Xuan Loc xenoliths are between 0.86 and 0.91 (Table S1), which lies within the compositional range of SCLM (Arai, 1994). Lherzolites have overlapping but slightly lower core ol Mg# (0.86-0.91) than harzburgites and dunites (0.89-0.91). Olivine Mg# correlates with cpx Mg#, opx Mg#, and sp Cr#. The CaO content of ol is less than 0.06 wt.% and NiO is between 0.3 to 0.42 wt. % for all samples.

We find that sp compositions are highly variable across Vietnam peridotites. Spinel grains in lherzolites have Al₂O₃ and MgO ranges of 52.9 to 60.0 wt.% and 19.3 to 20.9 wt.%, respectively. Harzburgite and dunite samples have lower Al₂O₃ (32.3-34.6 wt.%) and lower MgO (12.9-16.7 wt.%) than lherzolites. One dunite sample, VN-2018-36-XL-1, has notably higher FeO (26.1 wt.%) and lower Al₂O₃ (24.0 wt.%) than the other refractory, i.e., harzburgite and dunite samples (13.5 wt. % and 33.1 wt. %, respectively). Spinel Cr# for the lherzolites (0.08-0.17) and harzburgites/dunites (0.40-0.50) all lie within the compositional range previously documented for SCLM-derived xenoliths (Arai, 1994).

Lherzolites from Pleiku and Xuan Loc have cpx cores with high Al₂O₃ (5.48-7.37 wt.%), low Cr# (0.06-0.12), and high Na₂O (1.43-1.93 wt.%) (Figure 2b). Clinopyroxene cores from harzburgites generally exhibit comparatively low Al₂O₃ (1.49-2.73 wt.%), high Cr# (0.18-0.38), and low Na₂O (0.57-1.60 wt.%). Orthopyroxene displays a similar pattern to cpx, with lherzolite-hosted grains exhibiting high Al₂O₃ (3.26-4.43 wt.%) and low Cr# (0.04-0.08 wt.%), while opx in harzburgites exhibits relatively low Al₂O₃ (1.86-2.21 wt.%) and high Cr# (0.11-0.16).

4.3. Trace element concentrations

Average trace element data for cpx and opx from Pleiku and Xuan Loc peridotites are provided in Table S2. The primitive mantle normalized trace element concentrations and REE patterns (REE_{PM}) for cpx are shown in Figure 3a-b. Type-F cpx from Pleiku and Xuan Loc display depleted, spoon-shaped, and enriched REE_{PM} patterns. The “depleted” and “spoon-shaped” patterns for Pleiku xenoliths are generally more steeply sloped than those for Xuan Loc samples, with a median $(Ce/Yb)_{PM}$ of 0.17 for Pleiku versus 0.51 for Xuan Loc. Most type-F xenoliths have a relatively flat slope from the middle rare earth elements (MREE) to heavy rare earth elements (HREE), with Pleiku and Xuan Loc having $(Sm/Yb)_{PM}$ from 0.53 to 1.04 and 0.60 to 1.22, respectively. “Depleted” patterns show a positive steep slope from light rare earth elements (LREEs) to MREEs with Pleiku and Xuan Loc showing $(La/Sm)_{PM}$ from 0.10 to 0.14 and 0.33 to 0.61, respectively. “Spoon-shaped” patterns are similar, but also exhibit a LREE enrichment with $(La/Ce)_{PM} = 1.08$ to 1.79 (Pleiku) and 1.22 and 1.38 (Xuan Loc). VN-2018-36-XL-8 and VN-2018-21-PL-4 have “enriched” REE_{PM} patterns exhibiting negative slopes from LREE to MREE ($(La/Sm)_{PM} = 1.55$). Similar types of REE patterns have been observed in type-F xenoliths from prior studies in both the Pleiku and Dalat volcanic plateaus (Figure 3) (Anh et al., 2021; Nguyen & Kil, 2019, 2020).

The Type-R cpx have enriched REE_{PM} patterns and, in this study, are only present in the Xuan Loc suite of samples, but similar REE patterns have likewise been reported for published xenolith samples from Dalat and Pleiku (Figure 3a,b) (Anh et al., 2021; Nguyen & Kil, 2019, 2020). Clinopyroxenes from the refractory xenoliths studied here have elevated LREE and low HREE, with steep negative slopes from LREEs to HREEs. These incompatible element enriched patterns exhibit $(Ce/Yb)_{PM}$ from 6.14 to 9.15. Clinopyroxene from sample VN-2018-36-XL-1

has an S-shaped REE_{PM} pattern, with a positive slope from the MREEs to HREEs and (La/Sm)_{PM} of 0.71.

Additional trace element compositions show that other incompatible elements in cpx (Rb, Nb, Pb, Sr, Ti) are overall fractionated compared to the REE (Figure 3c,d). Rubidium in cpx is depleted in almost all samples, with concentrations near or below analytical detection limits. Titanium, Pb, and Nb display negative anomalies across all samples compared to neighboring REE in Figure 3c-d. Sample VN-2018-21-PL-2 has a steep negative slope from U_{PM} to Pb_{PM}. Strontium is also variably enriched and depleted in the samples when compared to REE_{PM}, and high-Sr samples overall tend to exhibit spoon-shaped REE_{PM} patterns.

Orthopyroxene grains have REE_{PM} concentration patterns that strongly correlate with cpx patterns (Figure 3e). Type-F opx overall have REE_{PM} that exhibit depleted and spoon-shaped patterns. Some opx samples have very low LREE concentrations near analytical detection limits. Type-R opx have convex downward REE_{PM} patterns, with a negative slope from LREE to MREE and a positive slope from MREE to HREE. Moderately incompatible high-field strength elements (HFSE) Ti, Zr, and Hf have positive anomalies relative to REE. The Lu/Hf ratios in opx are also systematically higher by a factor of 3.8 than those in coexisting cpx.

4.4. Radiogenic isotopes

The Sr, Nd, and Pb isotopic compositions of cpx mineral separates from Pleiku and Xuan Loc are provided in Table 2 and Figure 4. Samples VN-2018-21-PL-1, VN-2018-21-PL-3, VN-2018-21-PL-6, VN-2018-21-PL-7, and VN-2018-21-PL-9 all contained low Pb abundances which necessitated analysis via multi-Daly measurement. Although the analytical uncertainty for these measurements is ~10x greater than for the measurements using TI for mass fractionation correction (Table 2), these samples fall along the same ²⁰⁶Pb/²⁰⁴Pb-²⁰⁷Pb/²⁰⁴Pb and ²⁰⁶Pb/²⁰⁴Pb-

$^{208}\text{Pb}/^{204}\text{Pb}$ correlations as other samples (Figure 4a,c). Clinopyroxenes from Pleiku lherzolites exhibit highly variable isotopic compositions: $^{206}\text{Pb}/^{204}\text{Pb} = 16.69\text{--}18.86$, $^{207}\text{Pb}/^{204}\text{Pb} = 15.34\text{--}15.58$, $^{208}\text{Pb}/^{204}\text{Pb} = 36.48\text{--}38.85$, $^{87}\text{Sr}/^{86}\text{Sr} = 0.70238\text{--}0.70337$, and $^{143}\text{Nd}/^{144}\text{Nd} = 0.513091\text{--}0.514190$ (Figure 4). Clinopyroxene grains from the Xuan Loc peridotites display a similar range of isotopic compositions to those from Pleiku: $^{206}\text{Pb}/^{204}\text{Pb} = 17.13\text{--}18.37$, $^{207}\text{Pb}/^{204}\text{Pb} = 15.44\text{--}15.57$, $^{208}\text{Pb}/^{204}\text{Pb} = 37.08\text{--}38.64$, $^{87}\text{Sr}/^{86}\text{Sr} = 0.70257\text{--}0.70405$, and $^{143}\text{Nd}/^{144}\text{Nd} = 0.512755\text{--}0.513371$, but the more enriched values (e.g., higher $^{87}\text{Sr}/^{86}\text{Sr}$ and lower $^{143}\text{Nd}/^{144}\text{Nd}$) are observed in the refractory samples for that locality.

The cpx Pb isotopic compositions mostly lie near or within the range observed in Indian MORB (Gale et al., 2013) and form positive correlations between $^{206}\text{Pb}/^{204}\text{Pb}$ and both $^{207}\text{Pb}/^{204}\text{Pb}$ and $^{208}\text{Pb}/^{204}\text{Pb}$ that plot above the northern hemisphere reference line (NHRL) (Figure 4a,c). The Xuan Loc cpx compositions also overall exhibit systematically more radiogenic $^{208}\text{Pb}/^{204}\text{Pb}$ than Pleiku cpx on a plot of $^{208}\text{Pb}/^{204}\text{Pb}$ versus $^{206}\text{Pb}/^{204}\text{Pb}$ (Figure 4c).

$^{206}\text{Pb}/^{204}\text{Pb}$, $^{207}\text{Pb}/^{204}\text{Pb}$, $^{208}\text{Pb}/^{204}\text{Pb}$ and $^{87}\text{Sr}/^{86}\text{Sr}$ in the xenoliths largely do not exhibit well-defined correlations with trace element data, though there is a broad correlation between $^{87}\text{Sr}/^{86}\text{Sr}$ and Sm/Nd ratios. Clinopyroxene $^{87}\text{Sr}/^{86}\text{Sr}$ and ϵ_{Nd} exhibit a negative correlation extending from the Indian-MORB field to highly radiogenic $^{143}\text{Nd}/^{144}\text{Nd}$ and unradiogenic $^{87}\text{Sr}/^{86}\text{Sr}$, most clearly in Pleiku fertile xenoliths (Figure 4d). $^{143}\text{Nd}/^{144}\text{Nd}$ also exhibits a positive correlation with Sm/Nd ratios and a strong negative correlation with Ce/Yb_{PM} (Figure 5). Type-R xenoliths with $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ measurements plot at the far right side of the type-F xenolith range, at the enriched limit of the Indian MORB field. Type-F xenoliths also have highly variable $^{143}\text{Nd}/^{144}\text{Nd}$ (0.512921–0.348419) with many samples plotting to the left of and above the MORB field in Figure 4d.

4.5. Thermometry and barometry of Vietnam xenoliths

Calculated equilibrium temperatures and pressures for Vietnam xenoliths from this study are presented in Table 3 and Figure 6. Temperatures based on major element composition were calculated using the two-pyroxene thermometer (T_{BKN}) of Brey and Kohler (1990) and the two-pyroxene thermometer (equation 36) of Putirka (2008) (T_{36}), and temperatures based on REE + Y compositions were calculated using the two-pyroxene thermometer (T_{REE}) of Liang et al. (2013) (Figure S2). For T_{BKN} , we assumed a pressure of 15 kbar, and a ± 5 kbar change in pressure results in a 10 °C difference in calculated temperature. Temperatures and pressure were also calculated iteratively using equation 36 for temperature (T_{36}) and equation 38 for pressure from Putirka (2008) (P_{38}). The two-pyroxene barometer we used (P_{38}) is temperature-independent. The temperature-dependent barometer of Putirka (2008; equation 39) in principle provides greater precision; however, pressures determined with this barometer were geologically unreasonable, providing either negative values or placing many of the mantle peridotites within the crust (< 5 kbar). The pressures calculated using equation P_{38} , in contrast, largely appear geologically reasonable, generally straddling the calculated geotherm estimated from local heat flow (see Section 5.2). Based on our observations and calculated results, the cpx and opx from this study appear to have experienced protracted cooling and subsolidus exsolution. As a result, the cpx and opx compositions have higher and lower wollastonite components, respectively, than the pyroxenes used by Putirka (2008) to calibrate the 2-pyroxene thermobarometers (Putirka, pers. comm.). We thus consider the temperature-independent pressure estimates (P_{38}) to be more accurate for the xenoliths analyzed here.

Equilibration pressures and temperatures of the xenolith samples were determined using the equations described above and by pairing either average cpx and opx core compositions or average cpx and opx rim compositions together, to evaluate multiple possible episodes of

equilibration under different conditions. For most samples, core and rim P and T estimates are very similar: the average absolute differences between core and rim temperatures are 10 °C for T_{36} and 20 °C for T_{BKN} . Likewise, absolute differences in estimated core and rim pressures average 0.8 kbar, which is well within the ± 3.7 kbar uncertainty of the barometer (Putirka, 2008). However, although small on average, differences in estimated core and rim temperatures were systematically larger using T_{BKN} than T_{36} , and several samples had estimated rim temperatures over 50 °C less than core estimates. In contrast, all rim-core estimated temperatures were within 25 °C for T_{36} calculations, except for two samples (VN-2018-36-XL-15 and VN-2018-36-XL-19), which had lower estimated rim than core temperatures by differences of 37 and 35 °C, respectively.

The equilibrium T_{BKN} temperatures for Pleiku and Xuan Loc xenoliths range from 891-1067 °C and 765-1026 °C, respectively, calculated using mineral core compositions (Figure 6). Calculated T_{36} values span a narrower range, from 956-1034 and 926-1006 °C in the Pleiku and Xuan Loc xenoliths, respectively. T_{REE} spans a similar range, with Pleiku xenoliths ranging from 887 to 1032 °C and Xuan Loc xenoliths from 821 to 1076 °C. Although on average, T_{REE} and T_{36} are in good agreement (with an average absolute difference of 56 °C for the full data set), and the differences between the two calculators do not appear to be systematic, several samples have calculated T_{REE} values that are either higher or lower than T_{36} by over 100 °C. Significantly, both T_{REE} and the absolute difference between T_{REE} and T_{36} are negatively correlated with the calculated pressure, P_{38} . This negative correlation is the opposite of the expected temperature profile for the mantle lithosphere, and we thus consider it to likely be an artifact of the rare earth thermometer inadequately incorporating the pressure dependence of REE partitioning between cpx and opx.

Calculated pressures (P_{38}) for Pleiku xenoliths range from 11.7-15.8 kbar, similar to type-F peridotites from Xuan Loc, which have calculated equilibration pressures of 12.1-16.1 kbar. In contrast, type-R xenoliths have lower pressures ranging from 7.1 to 10.0 kbar. No correlation is observed between P_{38} and the temperatures calculated using the Putirka two-pyroxene thermometer (T_{36}). However, a broad positive correlation between P_{38} and temperature is observed for T_{BKN} . This pressure-temperature correlation roughly parallels the geothermal gradient calculated below (Section 5.2, Figure 7) using the local estimated surface heat flow.

5. Discussion

5.1. Partial melting of the lithospheric mantle beneath Indochina

The observed variations in major element, trace element, and isotopic compositions among Vietnam xenoliths are indicative of mantle residues that have undergone variable degrees of progressive melt extraction and other magmatic processes (e.g., melt addition, refertilization, and metasomatism). The presence of two compositional groups of xenoliths, type-F and type-R, also suggests that the SCLM beneath Indochina may record two distinct histories, although the presence of some overlapping and intermediate compositions among all samples from Vietnam (e.g., Figures 3, 4, and 5) (Anh et al., 2021; Nguyen & Kil, 2019, 2020) makes it less certain that this is the case.

Xenoliths from Pleiku and Xuan Loc span a wide range in composition, from relatively fertile lherzolites characterized by high (>5 wt.%) Al_2O_3 in cpx and low (<0.2) spinel Cr#, to refractory harzburgites with lower Al_2O_3 (<3 wt.%) in cpx and high sp Cr# (>0.4) (Figure 2, 5). Although the additional published data from Anh et al. (2021) and Nguyen and Kil (2019, 2020) include a few intermediate or overlapping compositions, the reported xenolith compositions are

largely bimodal, defining fertile (type-F) and refractory (type-R) populations. We interpret these major element variations to primarily indicate that the two populations of xenoliths have experienced different extents of prior partial melting.

Type-R harzburgites and dunites exhibit major element and trace element mineral compositions similar to other mantle xenoliths thought to derive from ancient SCLM (Griffin et al., 2008; McDonough, 1990), namely low Al_2O_3 (1.5-2.8 wt.%), high Mg# (0.918-0.938), and high $(\text{LREE}/\text{HREE})_{\text{PM}}$ in cpx. Low modal cpx in these samples is also indicative of previous partial melting that consumes clinopyroxene (Herzberg, 1999). Type-R samples from Xuan Loc have high sp Cr#s and strong depletions in HREE (Table S1), which also likely indicate a high degree of partial melt extraction. There is also a strong negative correlation between Ti and sp Cr# across all type-R xenoliths, which can be generated by high degrees of partial melt extraction (Figure 5). The refractory harzburgites from Pleiku (Nguyen & Kil, 2019) also have major and trace element compositions that resemble type-R refractory samples from Xuan Loc, suggesting that both the Pleiku and Xuan Loc volcanic centers overlie SCLM containing highly-depleted residues, while harzburgites from Ba Ria and Nui Nua appear more intermediate, with higher Al_2O_3 that overlaps with the fertile lherzolites from the region (Nguyen & Kil, 2020). Relatedly, refractory xenoliths from Ba Ria and Nui Nua have REE patterns that more closely resemble fertile xenoliths (Nguyen & Kil, 2020), while all type-R xenoliths from both Pleiku and Xuan Loc have relatively high LREE enrichment and are characterized by low ϵ_{Nd} (+2.3 to +3.2) (Nguyen & Kil, 2019) (Table 2), suggesting an overall complex history of metasomatism or melt-rock interaction.

In contrast, Type-F xenoliths from Vietnam contain cpx with relatively fertile major element compositions (e.g., $\text{Al}_2\text{O}_3 = 5.5 - 7.4$ wt.%) that fall within the range of abyssal

peridotites, though Vietnamese xenoliths are slightly less fertile than estimates for depleted mantle cpx compositions (~7.9 wt. % Al_2O_3 in cpx) (Warren, 2016; Workman & Hart, 2005). The depleted and spoon-shaped REE_{PM} patterns exhibited by some type-F lherzolites from Pleiku, Xuan Loc, and Dalat are consistent with both partial melting, and modest subsequent enrichment during silicate melt metasomatism (see Section 5.3 for further discussion of metasomatism). Clinopyroxenes from abyssal peridotites display similar LREE patterns to our depleted xenoliths, although they typically exhibit even more extreme LREE depletion (Figure 3) (Warren, 2016, and references therein). While Type-F lherzolites from Pleiku, Xuan Loc, and Dalat all exhibit a range of depleted, spoon-shaped, and LREE-enriched REE_{PM} patterns in cpx, our Pleiku xenoliths are generally more LREE-depleted, extending to lower $(\text{LREE}/\text{HREE})_{\text{PM}}$ than our Xuan Loc samples (Figure 5) and comparable to the most LREE-depleted Dalat samples from the literature (Anh et al., 2021; Nguyen & Kil 2020). Rather than recording a partial melting history, the trace element depletion patterns in some type-F xenoliths may reflect moderately high degrees of metasomatic enrichment, in agreement with the more radiogenic $^{87}\text{Sr}/^{86}\text{Sr}$ and unradiogenic ε_{Nd} in many of the fertile xenoliths, and as explored further in Section 5.3. Sample VN-2018-36-XL-8 appears to be the most affected by this character of metasomatic enrichment among the Vietnamese type-F xenoliths: it displays notable LREE-enrichment and has the most enriched radiogenic isotopic composition of the F-type xenoliths analyzed in this study.

Previous studies have shown that sp Cr# and cpx Yb content are well correlated in many mantle peridotites, and can be used to estimate the extent of prior partial melting (e.g., Hellebrand et al., 2001). However, although incompatible trace element abundances and ratios (e.g., La/Sm) should also be strongly affected by partial melting, numerous studies have shown

that highly incompatible elements like the LREE are particularly susceptible to subsequent metasomatic re-enrichment (e.g., Warren, 2016), and so may not directly record prior melt extraction in lithospheric mantle peridotites. The spoon-shaped and LREE-enriched REE patterns observed in many of the xenoliths examined here thus suggest they have experienced metasomatic enrichment. However, this metasomatism does not appear to have significantly affected either sp compositions or HREE abundances in cpx, as evidenced by the strong correlation between cpx [Yb] and spinel Cr# in our samples (Figure 5).

The melting model of Hellebrand et al. (2001) calculates the degree of fractional melting experienced by mantle peridotites as a function of sp Cr#, assuming an initial LREE-depleted lherzolite composition after Loubet et al. (1975). The Hellebrand et al. (2001) empirical function was derived using melt fractions calculated from Dy, Er, and Yb concentrations, after the approach of Johnson et al. (1990). Calculations using this model for Vietnam xenoliths yield melt fractions of 15-17% for type-R xenoliths from Xuan Loc, and up to 6.1% for type-F xenoliths from both localities (Figure 5). The different xenolith populations are also characterized by notably different pressures of equilibration, with the refractory xenoliths constrained to shallower pressures overall. Our Xuan Loc type-R xenoliths all have pressures < 10 kbar, and type-R xenoliths from the nearby Dalat volcanic plateau have calculated pressure estimates of 9.3 to 12.4 kbar (Nguyen & Kil, 2020). Applying this melting model to the published Pleiku dataset from Nguyen and Kil (2019) yields melt fractions of 10.3-18.5% and up to 7.1% for type-R and type-F xenoliths, respectively, ranges which overlap with or approach results for our samples. However, unlike Xuan Loc xenoliths, the refractory Pleiku xenoliths are not entirely restricted to low (< 10 kbar) equilibration pressures. Nguyen and Kil (2020) likewise estimated partial melting of 1-13% for their harzburgite and lherzolite xenolith samples, and Anh

et al. (2021) calculated that their type-F lherzolite xenoliths from Dalat experienced up to 5% partial melting, consistent with our type-F samples.

5.2. Thermal and chemical structure of the SCLM beneath Pleiku and Xuan Loc

Our pressure and temperature estimates for the Pleiku and Xuan Loc xenoliths, combined with their chemical and isotopic compositions, provide a more detailed picture of the thermal and chemical structure of the SCLM beneath these regions. Figure 7 shows the estimated pressures and temperatures of the Xuan Loc and Pleiku xenoliths for our temperature estimates, using both the Brey & Kohler (1990) two-pyroxene thermometer (T_{BKN}) (Figure 7a) and the two-pyroxene thermometer of Putirka (2008) (T_{36}) (Figure 7b). Using the same methods, we also determined pressure and temperature estimates for the Dalat and Pleiku xenoliths analyzed by Nguyen and Kil (2019, 2020) and Anh et al. (2021), and these estimates are included in Figure 7 for comparison.

Figure 7 indicates that the calculated xenolith equilibration pressures and temperatures generally cluster near geothermal gradients calculated using regional heat flow data, suggesting they are broadly in agreement with those estimated gradients. Unfortunately, there are currently no local heat flow data in the immediate vicinity of the study area, so best estimates come from regional datasets, which have been compiled into a heat flow map (Hall, 2002). From this map, the estimated heat flow values for our study areas are 85 mW/m² and 90 mW/m² for Pleiku and Xuan Loc, respectively. The measured thickness of the crust is 32 km at Pleiku and 28 km at Xuan Loc, from broadband seismic data (Yu et al., 2017a). We assumed average densities (ρ) of 2800 kg/m³ and 3250 kg/m³ for the crust and mantle, respectively, with thermal conductivities of 2.7 W/(m·K) for the crust and 4.0 W/(m·K) for the mantle (Turcotte & Schubert, 2002). The heat generation (HG) of the crust was then calculated using the model of Hasterok and Chapman (2011):

$$HG = 10^{-5} \rho [3.5C_{K_2O} + 9.67C_U + 2.63C_{Th}]$$

where HG is expressed in units of $\mu W/m^3$, and C_i values indicate concentrations for each species *i*. Here we used the estimated abundances of heat-producing elements in the average continental crust from Rudnick and Fountain (1995): 1.88 w.t% K_2 , 5.6 ppm Th, and 1.4 ppm U.

Xenolith P-T estimates using T_{36} are located above and below the estimated geothermal gradients and show no clear correlation between depth and temperature (Figure 7b). In contrast, the P-T estimates using T_{BKN} are generally closer to the geothermal gradients, though shifted to moderately lower temperatures at a given pressure, especially at higher pressures (Figure 7a). One possible explanation for this mismatch is that regional surface heat flow does not reflect a semi-steady-state conductive gradient, but may have been elevated due to heat deposition in the crust during magma transport and storage. Thus, the actual conductive heat flow within the local SCLM may be lower than required to support a measured surface heat flow of 85-90 mW/m².

Regardless of the thermometer used, the Dalat and Xuan Loc xenoliths record a chemically stratified lithospheric mantle. Specifically, the population of refractory (type-R) xenoliths from Dalat and Xuan Loc generally have lower estimated pressures than fertile-type xenoliths from Dalat, Xuan Loc, or Pleiku, though the populations do overlap (Figure 7). However, similar stratification is not as clear for the type-R xenoliths sampled at Pleiku, which is located to the north of Dalat and Xuan Loc: although the current study only retrieved fertile (type-F) xenoliths from Pleiku, data from Nguyen and Kil (2019) include type-R Pleiku xenoliths that are not as restricted to shallow equilibration depths. That said, although Pleiku xenoliths do exhibit more overlap, they nonetheless appear to have some depth dependence. For example, in Figure 7, 5-6 Pleiku samples with shallower equilibration depths (< 40 km) are refractory, while the majority of those with greater equilibration pressures are fertile.

The shallower, type-R Xuan Loc xenoliths are also LREE-enriched with relatively unradiogenic $^{143}\text{Nd}/^{144}\text{Nd}$ signatures compared to the deeper, type-F xenoliths. Figure 7c shows the resulting variations in $^{143}\text{Nd}/^{144}\text{Nd}$ with estimated depth. Although two type-F Dalat samples have somewhat shallower equilibration pressures relative to their $^{143}\text{Nd}/^{144}\text{Nd}$ compositions (Anh et al., 2021; Nguyen & Kil, 2020), the other samples record a broad compositional correlation with depth. Correlations between depth and isotopic compositions are weaker for $^{87}\text{Sr}/^{86}\text{Sr}$ and absent for Pb-isotopes, however, perhaps reflecting a greater sensitivity of Sr and Pb to metasomatic overprinting than Nd in the local lithospheric mantle, whether because of the initial composition of the unmetasomatized type-R peridotites or due to the composition of the metasomatizing fluid, which is explored in further detail in Section 5.3. Although these isotopes may reflect a more complex history, at Xuan Loc and Dalat, the refractory-type lithosphere does appear to be underlain by more fertile, LREE-depleted mantle. One possible explanation for the apparent depth-dependent compositional distribution in Figure 7c, which is consistent with the presence of two largely distinct xenolith populations, is that the refractory samples represent fragments of the original lithospheric mantle, which has since been partially replaced by more fertile material, an interpretation that is consistent with the model suggested by Anh et al. (2021). These more fertile peridotites, in turn, may have been emplaced more recently in a younger, deeper layer, and were derived from the more recently convecting asthenospheric mantle. However, in contrast, both the handful of more intermediate xenolith compositions (e.g., harzburgites with elevated Al_2O_3 and $\text{sp Cr\#} = \sim 0.25$; Nguyen & Kil, 2020), and the correlations shown in Figure 5 suggest that the xenoliths were all affected by a single metasomatic episode or event, which would not necessitate the proposed history of thinning and replacement of the deep lithospheric mantle layer.

The following section explores the metasomatic histories recorded by F- and R-type xenoliths further, and aims to clarify which genetic models are most consistent with all of the observed geochemical correlations and variations in mantle composition with depth.

5.3. Metasomatism of the SCLM beneath Vietnam

Given that LREE are highly incompatible, they should be strongly depleted by removal from the source rock during partial melting. As discussed above, the apparent enrichment in these elements in both fertile and refractory xenoliths with “spoon-shaped” and LREE-enriched REE_{PM} patterns requires an additional explanation, and likely indicates metasomatism of incompatible-element depleted residues by the later addition of a LREE-enriched melt or fluid. This evidence for variable metasomatic enrichment in all sampled regions (Pleiku, Dalat, and Xuan Loc) raises several related questions: 1) Do the different enrichment patterns observed in the fertile and refractory xenoliths reflect two distinct metasomatic episodes, or are they related to a single event? 2) When did metasomatic enrichment occur? and 3) What was the nature of the metasomatic agent(s)?

One scenario that could explain the character of type-F xenoliths is more recent emplacement of fertile (type-F), young lithospheric mantle from the convecting asthenosphere, similar to the lithospheric replacement model suggested by, e.g., Anh et al. (2021). As discussed previously, refractory xenoliths at Xuan Loc and Dalat are restricted to relatively shallow equilibration pressures (≤ 12.4 kbar; Figure 7). In this scenario, the type-F xenoliths from Pleiku and from the deeper portions of the Xuan Loc and Dalat lithosphere derived from convecting asthenospheric mantle material that was relatively recently emplaced as younger lithosphere, beneath the older refractory rocks. This model is similar to that proposed to explain the presence of a bimodal xenolith population with similar refractory/LREE-enriched and fertile/LREE-

depleted characteristics within the Rio Grande Rift, North America (Byerly and Lassiter, 2012) and was previously proposed by Nguyen and Kil (2019) to explain the presence of both refractory and fertile mantle beneath Pleiku. In this scenario, the metasomatic enrichment of the older, refractory xenoliths could reflect addition of subduction-derived melts or fluids during the Mesozoic. The modest metasomatic enrichment observed in some fertile xenoliths with spoon-shaped REE profiles then perhaps reflect a second metasomatic episode, unrelated to that observed in the older refractory xenoliths; this second event could have occurred either before (i.e., still within the convecting asthenosphere) or after emplacement as new lithosphere. In either case, this second metasomatic episode could potentially indicate the addition of low-degree melts generated during tectonic extrusion or otherwise related to Cenozoic volcanism in Vietnam.

However, the Pleiku and Xuan Loc xenoliths also display a strong correlation between $^{143}\text{Nd}/^{144}\text{Nd}$ and LREE enrichment, which strongly suggests a single mixing trend has affected both the refractory and fertile xenoliths. Figure 5 illustrates the correlations between $^{143}\text{Nd}/^{144}\text{Nd}$ and both Sm/Nd and Ce/Yb, across both (fertile and refractory) xenolith populations, suggesting modification by a single metasomatic agent for all of the xenoliths. Other geochemical patterns (e.g., Figures 3-4) show overlap and intermediate compositions between the xenolith types and locales, shedding doubt on the proposed bimodal character of the SCLM. As a second, alternative scenario, type-F xenoliths in Vietnam may thus record a related but less extensive history of melt-rock interactions to the type-R xenoliths, which would be consistent with the generally lower, and thus more easily modified trace element abundances in the refractory samples. That said, while type-F xenoliths from Pleiku and Xuan Loc share similar characteristics that suggest comparable processes have produced parts of the fertile SCLM in

both locations, samples from the two sites do exhibit a noteworthy difference in their trace element and radiogenic isotope compositions, specifically less radiogenic $^{208}\text{Pb}/^{204}\text{Pb}$ in Pleiku type-F xenoliths compared to those from Xuan Loc. This suggests a slightly different geochemical character to the metasomatic enrichment beneath Pleiku. Nonetheless, the fact that the Xuan Loc and Pleiku fertile xenoliths both fall along the same Ce/Yb - $^{143}\text{Nd}/^{144}\text{Nd}$ and Sm/Nd - $^{143}\text{Nd}/^{144}\text{Nd}$ trends, which also extend to the refractory xenolith compositions, strongly suggests that the metasomatism of the fertile and refractory xenoliths occurred as a single event.

In our proposed scenario, this single metasomatic event could be related to modification of the SCLM by subduction-related melts or fluids in the Mesozoic, or it could reflect more recent modification, possibly related to Cenozoic volcanism. A better understanding of the nature of the metasomatic agent(s) (e.g., whether or not the agent is subduction-related) may help to clarify the relationships between mantle lithospheric metasomatism and the tectonic evolution of the regional Indochinese SCLM. Nguyen and Kil (2019) previously suggested that the trace element characteristics of metasomatized, refractory Pleiku xenoliths suggest involvement of a subduction-related carbonate-rich melt and/or H_2O - CO_2 -rich fluids, and Anh et al. (2021) similarly advocated for a hydrous silicate metasomatizing agent beneath Dalat. Depletions of Ti relative to HREE and comparative enrichments in LREE in mantle rocks have previously been invoked to characterize the metasomatic agents responsible for melt-rock interactions (Coltorti et al., 1999). Given the notably low Ti/Eu (<1500) and high $(\text{La}/\text{Yb})_{\text{N}}$ in type-R cpx from Xuan Loc, such CO_2 -rich silicate melts (i.e., subduction-related melts) may be a plausible metasomatic agent here as well. Given the presence of Mesozoic subduction-related granitic batholiths throughout Indochina (Shellnutt et al., 2013; Gibbons et al., 2015), a carbonated silicate melt

would also likely support a Mesozoic origin for the metasomatism recorded in the refractory xenoliths.

Subduction-derived, CO₂-rich silicate melts, like those posited to have metasomatized the Pleiku xenoliths (Nguyen and Kil, 2019), are also expected to be enriched in large-ion lithophile elements (e.g., Rb, Sr) and depleted in high field-strength elements (HFSE) like Nb, Zr, and Hf (Ionov, 2002). At first glance, Vietnam xenolith cpx trace element patterns do appear consistent with involvement of a subduction-related, either carbonated or hydrous silicate melt (e.g., Anh et al., 2021), since many samples display depletions in Zr, Hf, and Ti (Figure 3). However, in most samples, these depletions in cpx are mirrored by positive anomalies in the same elements in coexisting opx (Figure 3e). Trace element patterns in cpx are controlled not only by the compositions of melts or fluids added to or removed from the host peridotite, but also by cpx/opx partitioning within the rock, which is a function of mineral composition, temperature, and pressure. Previous studies have demonstrated that HFSE anomalies in peridotitic cpx do not necessarily always reflect HFSE depletion of the bulk peridotite rock, and thus may not uniquely fingerprint a subduction origin for metasomatic enrichment (e.g., Byerly and Lassiter, 2015).

Although the trace element patterns described above provide somewhat ambiguous constraints on the origins of metasomatic overprinting, the isotopic compositions of the refractory and fertile xenoliths suggest the recorded metasomatism may be instead best explained by infiltration of melts similar to Cenozoic Vietnamese volcanic products. The refractory Xuan Loc xenoliths, which appear to record the greatest extent of metasomatic overprinting, have radiogenic isotope compositions that overlap with the depleted end of the field defined by Cenozoic basalts from south and central Vietnam. Many of the fertile samples from Xuan Loc, Dalat, and Pleiku, in contrast, overlap with the field defined by Indian MORB, and extend to

even more depleted compositions, particularly for Pb and Nd isotopes (Anh et al., 2021; Nguyen & Kil 2019, 2020). Although previous studies have proposed that the local SCLM is a significant melt source component in Cenozoic Vietnamese basalts, here we present an alternative interpretation from our xenolith measurements. We suggest that Cenozoic basalts, generated from enriched, possibly subduction-modified asthenospheric mantle components upwelling beneath the Vietnamese lithosphere, have themselves metasomatized and enriched the local SCLM, and that refractory lithospheric harzburgites were particularly susceptible to this overprinting.

In summary, although we cannot definitively preclude a scenario in which the refractory and fertile xenolith groups each experienced temporally separate metasomatic events with distinct origins, a more parsimonious interpretation of the correlations observed in Figure 5 is that both suites of xenoliths experienced a single metasomatic event, but were affected to differing degrees; and that this metasomatism is likely related to the same episode of mantle melting that generated the basalts which ultimately brought these xenoliths to the surface.

5.4. Working model of the evolution of SCLM beneath Vietnam

In the analysis above, we have made two potentially conflicting observations: 1) lithospheric mantle xenoliths from Vietnam mostly fall into two distinct populations (type-F and type-R) that have, on average, equilibrated at different depths and experienced different histories of melt extraction, and 2) the mantle xenoliths from Vietnam also exhibit trace element vs. radiogenic isotope compositions that suggest simultaneous metasomatism throughout the regional lithospheric mantle. Both of these observations appear robust: although they overlap, the type-F mantle xenoliths mostly have asthenosphere-like compositions, mineral compositions corresponding to equilibration depths between ~30-55 km (Figure 7), and a limited overall extent

of metasomatism compared to type-R xenoliths. Likewise, all measured Vietnamese mantle xenoliths exhibit $^{143}\text{Nd}/^{144}\text{Nd}$ vs. incompatible element correlations most consistent with a single, partially-overprinting metasomatic event.

Previous studies (e.g., Pan et al., 2013) have suggested that at some time in the past, some portions of the SCLM beneath Vietnam were replaced by asthenospheric mantle. In at least some locations, we also do observe that type-R xenoliths are restricted to shallower areas, suggesting they could derive from older refractory lithosphere that was not replaced. Our depth observations thus suggest that some portions of the older continental lithospheric mantle may have been partially removed by thinning or erosion during a prior event. Even neglecting our calculated mantle temperatures and xenolith trace element compositions, we do note that the xenolith equilibration pressures alone support a relatively thin continental lithospheric layer. The type-F xenoliths likewise exhibit $^{87}\text{Sr}/^{86}\text{Sr}$ (0.703724-0.703365) and ϵ_{Nd} isotopes (+5.51 to +30.28) that range from values resembling ultra-depleted mantle (Byerly and Lassiter, 2014; Cipriani et al., 2011; Mallick et al., 2014; Stracke et al., 2011) to those of MORB (Hofmann, 2007; Salters and Stracke, 2004; Workman and Hart, 2005). Type-F xenoliths also have relatively fertile major element compositions that likely experienced relatively small degrees of prior partial melting, suggesting the xenoliths were previously a part of a moderately depleted mantle region that underwent limited decompression melting.

Determining the pre-metasomatic origins of type-R xenoliths is more complex, given the high levels of overprinting of trace elements and isotopic compositions due to likely more thorough metasomatism (e.g., Figure 3). Mantle xenoliths with similar trace element and isotopic characteristics to the Vietnam type-R xenoliths have previously been attributed, but are not unique to pre-Phanerozoic lithosphere (e.g., Rio Grande Rift, Colorado Plateau, Zealandia, North

China Craton, Central Asian Orogenic Belt) (Byerly and Lassiter, 2012; Liu et al., 2012; McDonough, 1990; Pan et al., 2013; Scott et al., 2014; Warren, 2016). Type-R xenoliths from Vietnam (Nguyen and Kil, 2019, 2020) also exhibit some characteristics of peridotites that have experienced high degrees of fractional melting, and three Xuan Loc type-R xenoliths measured for radiogenic isotopes have both very consistent and relatively enriched isotopic signatures (e.g., moderately unradiogenic $^{143}\text{Nd}/^{144}\text{Nd} = 0.512755\text{--}0.512800$), in agreement with the previously published type-R xenolith from Dalat (0.512766; Nguyen & Kil, 2020). It is thus possible that the relatively shallow, type-R xenoliths are samples of an older lithospheric mantle that has now been partially removed (i.e., eroded or delaminated), consistent with past interpretations.

However, we note the presence of intermediate and even overlapping compositions and depths between the type-R and type-F groups, suggesting the distribution is not truly bimodal. This pattern, revealed by the compilation of more recent, larger data sets (Anh et al., 2021; Nguyen & Kil 2019, 2020) and this study, argues against the presence of such distinct regional layering and such a dramatic, wholesale lithospheric replacement scenario. While the ages and origins of Indochinese type-R and type-F xenoliths are uncertain, we posit that a single, major metasomatic episode has recently modified the entire lithospheric layer, but that metasomatic overprinting of prior geochemical signatures was incomplete. This scenario is most consistent with the observed geochemical correlations shown in Figure 5, which suggest one major metasomatic episode. The likely metasomatic agent is most consistent with infiltration and melt-rock reaction by Cenozoic magmas, indicating that metasomatism has been a recent event that may be particularly localized beneath the major Cenozoic volcanic centers.

We thus suggest an internally consistent, working model for the evolution and tectonic history of the lithospheric mantle beneath Vietnam and Indochina. In our model, the prior presence of both fertile and refractory mantle beneath Vietnam may not require a history of lithospheric removal and replacement, although the trend of generally higher fertility with depth is consistent with a fossil residue of ancient adiabatic upwelling. To place recent events in a tectonic context, during the India-Eurasia collision, Indochina was extruded along the Ailao Shan-Red River Fault Zone until the tectonic reorganization associated with a change in strike-slip sense along that fault zone and the coeval cessation of rifting in the South China Sea/East Vietnam Sea (Li et al., 2015; Zhu et al., 2009). Following that reorganization, the Sundaland block, which includes Indochina, lacked a “free surface” into which it could continue to laterally extrude due to the tectonic position of Borneo. It is unclear why such a scenario may have simultaneously induced such voluminous (70,000 km²) tholeiitic basalt production through mantle decompression, but one explanation is that the loss of a free surface during extrusion could have caused coupled lateral flow of the underlying asthenosphere to encounter an opposing mantle flow boundary beneath Borneo, triggering local upwelling and partial melting. Alternately or concurrently, the prior subduction and eventual sinking of the Pacific crustal slab under Indochina may have induced local upper mantle upwelling, as suggested by Yu et al. (2017b). Upwelling of the asthenosphere then resulted in small degrees of decompression melting, causing metasomatism of the overlying mantle lithosphere. This scenario would ultimately produce a relatively heterogeneous SCLM that partially preserves an older history of variable melt removal, but with metasomatic overprinting, perhaps especially under active Cenozoic volcanic centers. Although only one possible scenario, our working model can explain

the timing and geochemical compositions of Indochina basalts and xenoliths, in a way that is consistent with local tectonics.

6. Conclusions

In this study, we have characterized two suites of lithospheric mantle xenoliths from southern Vietnam. The measured major elements of peridotite mineral separates, trace elements of cpx and opx, and radiogenic isotope compositions of clinopyroxene in mantle peridotite xenoliths from two sites, Pleiku and Xuan Loc, provide the following insights into the lithospheric mantle of Vietnam:

1. The lithospheric mantle beneath Vietnam experienced a complex history of variable partial melting and subsequent metasomatism, which is preserved in lithospheric mantle-derived spinel peridotites.
2. Type-F or “fertile” spinel peridotites from Pleiku, Xuan Loc, and Dalat based on their spinel Cr#, exhibit fertile compositions with depleted, enriched, and spoon-shaped REE_{PM} patterns indicative of a lithosphere that has undergone low degrees of prior partial melting, and subsequently, variable degrees of silicate melt metasomatism. Type-F xenoliths from all three field locations have calculated equilibration temperatures and pressures that cover a wide range (Anh et al., 2021; Nguyen & Kil, 2019, 2020).
3. Similar to comparable xenoliths from previous work (Nguyen and Kil, 2019, 2020), type-R or “refractory” spinel peridotite xenoliths from Xuan Loc display refractory compositions with enriched and S-shaped REE_{PM} patterns, which indicate high degrees of partial melting followed by pervasive metasomatic enrichment, and share characteristics typical of ancient refractory SCLM. Calculated equilibration pressures for type-R xenoliths are notably restricted to relatively shallow equilibration depths.

4. The dramatic lithospheric erosion and replacement suggested by prior studies (e.g., Anh et al., 2021) may not in fact be necessary to explain Vietnam xenolith compositions. Regardless of the lithospheric mantle's more ancient origins, however, we posit that the underlying asthenosphere has recently upwelled and experienced decompression partial melting, both metasomatizing the overlying lithospheric mantle and producing extensive local basalt flows.

ACKNOWLEDGMENTS

We would like to thank Aaron Satkoski, Staci Loewy, Nate Miller, and Phil Orlandini for assistance in conducting major and trace element and isotopic analyses. Dinh Q. Sang, Le Duc Anh, Nguyen T. Mai, and Tran T. Huong assisted in fieldwork and sample collection in Vietnam, and Nicholas Richard assisted with some of the sample preparation and local map figures. This project was supported by a University of Nebraska-Lincoln College of Arts and Sciences international collaboration award and by NSF EAR-Tectonics award 1758972.

Data availability statement

Supporting data for this study are available in the data supplement provided, and in the EarthChem repository as three linked data sets (<https://doi.org/10.26022/IEDA/112849>, <https://doi.org/10.26022/IEDA/112848>, and <https://doi.org/10.26022/IEDA/112847>), which will be made fully public after peer review.

REFERENCES

830 Anh, H.T.H., Choi, S.H., Yu, Y., & Hieu, P.T. (2021). Geochemical constraints on the evolution
 831 of the lithospheric mantle beneath central and southern Vietnam. *Geosciences Journal*
 832 **25**(4), 433-451, doi:10.1007/s12303-020-0045-4.

833 Arai, S. (1994). Characterization of spinel peridotites by olivine-spinel compositional
 834 relationships: review and interpretation. *Chemical geology* **113**(3-4), 191-204.

835 Armstrong, J.T. (1988). Quantitative analysis of silicate and oxide materials: comparison of
 836 Monte Carlo, ZAF, and $\phi(\rho Z)$ procedures. *Microbeam Analysis*, 239–246.

837 Baker, J., Peate, D., Waight, T., & Meyzen, C. (2004). Pb isotopic analysis of standards and
 838 samples using a ^{207}Pb – ^{204}Pb double spike and thallium to correct for mass bias with a
 839 double-focusing MC-ICP-MS. *Chemical Geology* **211**(3-4), 275-303.

840 Ballhaus, C., Berry, R.F., & Green, D.H. (1991). High pressure experimental calibration of the
 841 olivine-orthopyroxene-spinel oxygen geobarometer: implications for the oxidation state
 842 of the upper mantle. *Contributions to Mineralogy and Petrology* **107**, 27–40,
 843 doi:10.1007/BF00311183.

844 Brey, G.P. & Kohler, T. (1990). Geothermobarometry in Four-phase Lherzolites II. New
 845 Thermobarometers, and Practical Assessment of Existing Thermobarometers. *Journal of*
 846 *Petrology* **31**, 1353–1378, doi:10.1093/petrology/31.6.1353.

847 Byerly, B.L. & Lassiter, J.C. (2012). Evidence from mantle xenoliths for lithosphere removal
 848 beneath the central Rio Grande Rift. *Earth and Planetary Science Letters* **355–356**, 82–
 849 93, doi:10.1016/j.epsl.2012.08.034.

850 Byerly, B.L. & Lassiter, J.C. (2015). Trace element partitioning and Lu–Hf isotope systematics
 851 in spinel peridotites from the Rio Grande Rift and Colorado Plateau: Towards improved

852 age assessment of clinopyroxene Lu/Hf–¹⁷⁶Hf/¹⁷⁷Hf in SCLM peridotite. *Chemical*
 853 *Geology* **413**, 146–158, doi:10.1016/j.chemgeo.2015.08.009.

854 Cipriani, A., Bonatti, E., & Carlson, R.W. (2011). Nonchondritic ¹⁴²Nd in suboceanic mantle
 855 peridotites. *Geochemistry, Geophysics, Geosystems* **12**, doi:10.1029/2010GC003415.

856 Coltorti, M., Bonadiman, C., Hinton, R.W., Siena, F., & Upton, B.G.J. (1999). Carbonatite
 857 Metasomatism of the Oceanic Upper Mantle: Evidence from Clinopyroxenes and Glasses
 858 in Ultramafic Xenoliths of Grande Comore, Indian Ocean. *Journal of Petrology* **40**, 133–
 859 165, doi:10.1093/petroj/40.1.133.

860 Cullen, A., Reemst, P., Henstra, G., Gozzard, S., & Ray, A. (2010). Rifting of the South China
 861 Sea: new perspectives. *Petroleum Geoscience* **16**, 273–282, doi:10.1144/1354-079309-
 862 908.

863 Dawson, J.B. (1984). Contrasting Types of Upper-Mantle Metasomatism?, in Kornprobst, J.,
 864 Ed., *Developments in Petrology*, 289–294, Elsevier. doi:10.1016/B978-0-444-42274-
 865 3.50030-5.

866 Flower, M., Tamaki, K., & Hoang, N. (1998). Mantle extrusion: A model for dispersed
 867 volcanism and DUPAL-like asthenosphere in East Asia and the western Pacific, in
 868 Flower, M.F.J., Chung, S., Lo, C., and Lee, T., Eds., *Geodynamics Series* **27**, 67–88,
 869 American Geophysical Union, Washington, DC. doi:10.1029/GD027p0067.

870 Frey, F.A. & Green, D.H. (1974). The mineralogy, geochemistry and origin of Iherzolite
 871 inclusions in Victorian basanites. *Geochimica et Cosmochimica Acta* **38**, 1023–1059,
 872 doi:10.1016/0016-7037(74)90003-9.

873 Frey, F.A. & Prinz, M. (1978). Ultramafic inclusions from San Carlos, Arizona: Petrologic and
874 geochemical data bearing on their petrogenesis. *Earth and Planetary Science Letters* **38**,
875 129–176, doi:10.1016/0012-821X(78)90130-9.

876 Gale, A., Dalton, C.A., Langmuir, C.H., Su, Y., & Schilling, J.-G. (2013). The mean composition
877 of ocean ridge basalts. *Geochemistry Geophysics Geosystems* **14**(3),
878 doi:10.1029/2012GC004334.

879 Gast, P.W. (1968). Trace element fractionation and the origin of tholeiitic and alkaline magma
880 types. *Geochimica et Cosmochimica Acta* **32**, 1057–1086, doi:10.1016/0016-
881 7037(68)90108-7.

882 Gibbons, A.D., Zahirovic, S., Müller, R.D., Whittaker, J.M., & Yatheesh, V. (2015). A tectonic
883 model reconciling evidence for the collisions between India, Eurasia and intra-oceanic
884 arcs of the central-eastern Tethys. *Gondwana Research* **28**, 451–492,
885 doi:10.1016/j.gr.2015.01.001.

886 Griffin, W.L., O'Reilly, S.Y., Afonso, J.C., & Begg, G.C. (2008). The Composition and
887 Evolution of Lithospheric Mantle: A Re-evaluation and its Tectonic Implications.
888 *Journal of Petrology* **50**, 1185–1204, doi:10.1093/petrology/egn033.

889 Hall, R. (2002). SE Asian heatflow: call for new data. *SEAPEX press* **5**, 54–56.

890 Hart, S.R. (1984). A large-scale isotope anomaly in the Southern Hemisphere mantle. *Nature*
891 **309**, 753–757, doi:10.1038/309753a0.

892 Hasterok, D. & Chapman, D.S. (2011). Heat production and geotherms for the continental
893 lithosphere. *Earth and Planetary Science Letters* **307**, 59–70,
894 doi:10.1016/j.epsl.2011.04.034.

895 Hellebrand, E., Snow, J.E., Dick, H.J.B., & Hofmann, A.W. (2001). Coupled major and trace
896 elements as indicators of the extent of melting in mid-ocean-ridge peridotites. *Nature*
897 **410**, 677–681, doi:10.1038/35070546.

898 Herzberg, C. (1999). Phase equilibrium constraints on the formation of cratonic mantle: Mantle
899 Petrology: Field Observations and High Pressure Experimentation, *Special Publications*
900 **6**, 241–257, Geochemical Society.

901 Hoang, N. & Flower, M. (1998). Petrogenesis of Cenozoic Basalts from Vietnam: Implication
902 for Origins of a ‘Diffuse Igneous Province’. *Journal of Petrology* **39**, 369–395,
903 doi:10.1093/petroj/39.3.369.

904 Hoang, N., Flower, M.F.J. & Carlson, R.W. (1996). Major, trace element, and isotopic
905 compositions of Vietnamese basalts: Interaction of hydrous EM1-rich asthenosphere with
906 thinned Eurasian lithosphere. *Geochimica et Cosmochimica Acta* **60**, 4329–4351,
907 doi:10.1016/S0016-7037(96)00247-5.

908 Hoang, N., Flower, M.F.J., Chi, C.T., Xuan, P.T., Quy, H.V. & Son, T.T. (2013). Collision-
909 induced basalt eruptions at Pleiku and Buôn Mê Thuột, south-central Viet Nam. *Journal*
910 *of Geodynamics*, **69**, 65–83, doi:10.1016/j.jog.2012.03.012.

911 Hofmann, A.W. (2007). Sampling Mantle Heterogeneity through Oceanic Basalts: Isotopes and
912 Trace Elements, in Carlson, R.W., Holland, H.D., and Turekian, K.K., Eds., *Treatise on*
913 *Geochemistry: The Mantle and Core*, 61–101. Elsevier, New York.

914 Ionov, D.A. (2002). Mechanisms and Sources of Mantle Metasomatism: Major and Trace
915 Element Compositions of Peridotite Xenoliths from Spitsbergen in the Context of
916 Numerical Modelling. *Journal of Petrology* **43**, 2219–2259,
917 doi:10.1093/petrology/43.12.2219.

918 Johnson, K.T.M., Dick, H.J.B., & Shimizu, N. (1990). Melting in the oceanic upper mantle: An
 919 ion microprobe study of diopsides in abyssal peridotites. *Journal of Geophysical*
 920 *Research* **95**, 2661, doi:10.1029/JB095iB03p02661.

921 Jolivet, L., Faccenna, C., Becker, T., Tesauro, M., Sternai, P., & Bouilhol, P. (2018). Mantle
 922 Flow and Deforming Continents: From India-Asia Convergence to Pacific Subduction.
 923 *Tectonics* **37**, 2887–2914, doi:10.1029/2018TC005036.

924 Lassiter, J.C., Blichert-Toft, J., Hauri, E.H., & Barszczus, H.G. (2003). Isotope and trace element
 925 variations in lavas from Raivavae and Rapa, Cook–Austral islands: constraints on the
 926 nature of HIMU- and EM-mantle and the origin of mid-plate volcanism in French
 927 Polynesia. *Chemical Geology* **202**, 115–138, doi:10.1016/j.chemgeo.2003.08.002.

928 Li, C.-F., Li, J., Ding, W., Franke, D., Yao, Y., Shi, H., Pang, X., Cao, Y., Lin, J., Kulhanek,
 929 D.K., & Williams, T. (2015). Seismic stratigraphy of the central South China Sea basin
 930 and implications for neotectonics. *Journal of Geophysical Research: Solid Earth* **120**,
 931 1377–1399, doi:10.1002/2014JB011686.

932 Liang, Y., Sun, C., & Yao, L. (2013). A REE-in-two-pyroxene thermometer for mafic and
 933 ultramafic rocks. *Geochimica et Cosmochimica Acta* **102**, 246–260,
 934 doi:10.1016/j.gca.2012.10.035.

935 Liermann, H.P. & Ganguly, J. (2003). Fe²⁺–Mg fractionation between orthopyroxene and spinel:
 936 experimental calibration in the system FeO–MgO–Al₂O₃–Cr₂O₃–SiO₂, and applications.
 937 *Contributions to Mineralogy and Petrology* **145**, 217–227, doi:10.1007/s00410-003-
 938 0444-3.

939 Liu, J., Carlson, R.W., Rudnick, R.L., Walker, R.J., Gao, S., & Wu, F. (2012). Comparative Sr–
 940 Nd–Hf–Os–Pb isotope systematics of xenolithic peridotites from Yangyuan, North China

941 Craton: Additional evidence for a Paleoproterozoic age. *Chemical Geology* **332–333**, 1–
942 14, doi:10.1016/j.chemgeo.2012.09.013.

943 Loubet, M., Shimizu, N., & Allègre, C.J. (1975). Rare earth elements in alpine peridotites.
944 *Contributions to Mineralogy and Petrology* **53**, 1–12, doi:10.1007/BF00402450.

945 Mallick, S., Dick, H.J.B., Sachi-Kocher, A., & Salters, V.J.M. (2014). Isotope and trace element
946 insights into heterogeneity of subridge mantle. *Geochemistry, Geophysics, Geosystems*
947 **15**, 2438–2453, doi:10.1002/2014GC005314.

948 McDonough, W.F. (1990). Constraints on the composition of the continental lithospheric mantle.
949 *Earth and Planetary Science Letters* **101**, 1–18, doi:10.1016/0012-821X(90)90119-I.

950 Metcalfe, I. (2013). Gondwana dispersion and Asian accretion: Tectonic and palaeogeographic
951 evolution of eastern Tethys. *Journal of Asian Earth Sciences* **66**, 1–33,
952 doi:10.1016/j.jseas.2012.12.020.

953 Michael, P.J. & Bonatti, E. (1985). Peridotite composition from the North Atlantic: regional and
954 tectonic variations and implications for partial melting. *Earth and Planetary Science*
955 *Letters* **73**, 91–104, doi:10.1016/0012-821X(85)90037-8.

956 Navon, O. & Stolper, E. (1987). Geochemical Consequences of Melt Percolation: The Upper
957 Mantle as a Chromatographic Column. *The Journal of Geology* **95**, 285–307,
958 doi:10.1086/629131.

959 Nguyen P.H., Bui, Q.C., & Nguyen, X.D. (2012). Investigation of earthquake tsunami sources,
960 capable of affecting Vietnamese coast. *Natural Hazards* **64**, 311–327.

961 Nguyen, T.C. & Kil, Y. (2019). The evolution of the lithospheric mantle beneath Ia Bang, Pleiku
962 plateau, Central Vietnam. *Journal of Asian Earth Sciences* **174**, 232–244,
963 doi:10.1016/j.jseas.2018.12.011.

964 Nguyen, C. & Kil, Y. (2020). Petrological characteristics of lithospheric mantle beneath Nui Nua
 965 and Ba Ria areas, southern Vietnam. *Geosciences Journal* **24**(5), 475-487,
 966 doi:10.1007/s12303-020-0017-8.

967 Pan, S., Zheng, J., Chu, L., & Griffin, W.L. (2013). Coexistence of the moderately refractory and
 968 fertile mantle beneath the eastern Central Asian Orogenic Belt. *Construction and*
 969 *Destruction of Cratons* **23**, 176–189, doi:10.1016/j.gr.2012.03.001.

970 Phach, P.V. & Anh, L.D. (2018). Tectonic evolution of the southern part of Central Viet Nam
 971 and the adjacent area. *Geodynamics & Tectonophysics* **9**, 801–825, doi:10.5800/GT-
 972 2018-9-3-0372.

973 Putirka, K.D. (2008). Thermometers and Barometers for Volcanic Systems. *Reviews in*
 974 *Mineralogy and Geochemistry* **69**, 61–120, doi:10.2138/rmg.2008.69.3.

975 Rohrmann, A., Kapp, P., Carrapa, B., Reiners, P.W., Guynn, J., Ding, L., & Heizler, M. (2012).
 976 Thermochronologic evidence for plateau formation in central Tibet by 45 Ma. *Geology*
 977 **40**, 187–190, doi:10.1130/G32530.1.

978 Royden, L.H., Burchfiel, B.C., & van der Hilst, R.D. (2008). The Geological Evolution of the
 979 Tibetan Plateau. *Science* **321**, 1054–1058, doi:10.1126/science.1155371.

980 Rudnick, R.L. & Fountain, D.M. (1995). Nature and composition of the continental crust: A
 981 lower crustal perspective. *Reviews of Geophysics* **33**, 267–309, doi:10.1029/95RG01302.

982 Salters, V.J.M. & Stracke, A. (2004). Composition of the depleted mantle. *Geochemistry,*
 983 *Geophysics, Geosystems* **5**, doi:10.1029/2003GC000597.

984 Scott, J.M., Hodgkinson, A., Palin, J.M., Waight, T.E., Van der Meer, Q.H.A., & Cooper, A.F.
 985 (2014). Ancient melt depletion overprinted by young carbonatitic metasomatism in the

986 New Zealand lithospheric mantle. *Contributions to Mineralogy and Petrology* **167**, 963,
987 doi:10.1007/s00410-014-0963-0.

988 Shaw, D.M. (1970). Trace element fractionation during anatexis. *Geochimica et Cosmochimica*
989 *Acta* **34**, 237–243, doi:10.1016/0016-7037(70)90009-8.

990 Shellnutt, J.G., Lan, C.-Y., Van Long, T., Usuki, T., Yang, H.-J., Mertzman, S.A., Iizuka, Y.,
991 Chung, S.-L., Wang, K.-L., & Hsu, W.-Y. (2013). Formation of Cretaceous Cordilleran
992 and post-orogenic granites and their microgranular enclaves from the Dalat zone,
993 southern Vietnam: Tectonic implications for the evolution of Southeast Asia. *Lithos* **182–**
994 **183**, 229–241, doi:10.1016/j.lithos.2013.09.016.

995 Sims, K. W., Hart, S. R., Reagan, M. K., Blusztajn, J., Staudigel, H., Sohn, R. A., Layne, G.D.,
996 Ball, L.A., & Andrews, J. (2008). ^{238}U - ^{230}Th - ^{226}Ra - ^{210}Pb - ^{210}Po , ^{232}Th - ^{228}Ra , and ^{235}U -
997 ^{231}Pa constraints on the ages and petrogenesis of Vailulu'u and Malumalu Lavas, Samoa.
998 *Geochemistry, Geophysics, Geosystems* **9**(4), <https://doi.org/10.1029/2007GC001651>.

999 Stracke, A., Snow, J.E., Hellebrand, E., von der Handt, A., Bourdon, B., Birbaum, K., &
1000 Günther, D. (2011). Abyssal peridotite Hf isotopes identify extreme mantle depletion.
1001 *Earth and Planetary Science Letters* **308**, 359–368, doi:10.1016/j.epsl.2011.06.012.

1002 Sun, S-S. & McDonough, W.F. (1989). Chemical and isotopic systematics of oceanic basalts:
1003 implications for mantle composition and processes. *Special Publications* **42**, 313–345,
1004 doi:10.1144/GSL.SP.1989.042.01.19. Geological Society, London.

1005 Tapponnier, P., Lacassin, R., Leloup, P.H., Shärer, U., Dalai, Z., Haiwei, W., Xiaohan, L.,
1006 Shaocheng, J., Lianshang, Z., and Jiayou, Z., 1990. The Ailao Shan/Red River
1007 metamorphic belt: Tertiary left-lateral shear between Indochina and South China. *Nature*
1008 **343** (6257), 431–437, doi: 10.1038/343431a0.

1009 Tapponnier P., Peltzer G., Armijo R., 1986. On the mechanics of the collision between India and
1010 Asia. In, Coward, M.P., Ries, A.C. (Eds.), *Collision Tectonics, Special Publications* **19**,
1011 115-157. Geological Society of London.

1012 Toramaru, A. & Fujii, N. (1986). Connectivity of melt phase in a partially molten peridotite.
1013 *Journal of Geophysical Research: Solid Earth* **91**, 9239–9252,
1014 doi:10.1029/JB091iB09p09239.

1015 Turcotte, D. & Schubert, G. (2002). *Geodynamics*, Cambridge University Press, Cambridge, UK,
1016 pp. 472. doi:10.1017/CBO9780511807442.

1017 Warren, J.M. (2016) Global variations in abyssal peridotite compositions. *Lithos* **248–251**, 193–
1018 219, doi:10.1016/j.lithos.2015.12.023.

1019 Workman, R.K. & Hart, S.R. (2005). Major and trace element composition of the depleted
1020 MORB mantle (DMM). *Earth and Planetary Science Letters* **231**, 53–72,
1021 doi:10.1016/j.epsl.2004.12.005.

1022 Workman, R. K., Hart, S. R., Jackson, M., Regelous, M., Farley, K. A., Blusztajn, J., Kurz, M.,
1023 & Staudigel, H. (2004). Recycled metasomatized lithosphere as the origin of the Enriched
1024 Mantle II (EM2) end-member: Evidence from the Samoan Volcanic Chain.
1025 *Geochemistry, Geophysics, Geosystems* **5**(4), <https://doi.org/10.1029/2003GC000623>.

1026 Yu, Y., Hung, T.D., Yang, T., Xue, M., Liu, K.H., & Gao, S.S. (2017). Lateral variations of
1027 crustal structure beneath the Indochina Peninsula. *Tectonophysics* **712–713**, 193–199,
1028 doi:10.1016/j.tecto.2017.05.023.

1029 Yu, Y., Gao, S.S., Liu, K.H., Yang, T., Xue, M., & Le, K.P. (2017), Mantle transition zone
1030 discontinuities beneath the Indochina Peninsula: Implications for slab subduction and

mantle upwelling. *Geophysical Research Letters* **44**(14), 7159-7167,
doi:10.1002/2017GL073528.

Zhu, M., Graham, S., & McHargue, T. (2009). The Red River Fault zone in the Yinggehai Basin,
South China Sea. *Tectonophysics* **476**, 397–417, doi:10.1016/j.tecto.2009.06.015.

FIGURE CAPTIONS

Figure 1. a. Map of southern and southeast Asia, illustrating extrusion of Indochina due to the
India-Eurasia collision (gray arrows) (after Taponnier et al., 1986, 1990). “Red river” = Ailao
Shan-Red River Fault Zone. **b.** Map of study area and sample collection localities. Major
mapped faults and volcanic centers are modified from published literature (Hoang et al., 1996,
2013; Nguyen et al., 2012; Phach & Anh, 2018).

Figure 2. a. Clinopyroxene Al_2O_3 rim/core ratio versus cpx Cr_2O_3 rim/core ratio. **b.** Cpx core
 Al_2O_3 wt% vs. cpx Mg# (molar $\text{Mg}/(\text{Mg} + \text{Fe})$). Small symbols show published xenolith data
from Pleiku (Nguyen & Kil, 2019) and Dalat (Anh et al., 2021; Nguyen & Kil, 2020).

Figure 3. Rare earth and trace element concentrations of samples from this study normalized to
primitive mantle (Sun and McDonough, 1989). Gray solid lines show type-F xenoliths and gray
dashed lines show type-R xenoliths from Pleiku and Dalat (Anh et al., 2021; Nguyen and Kil,
2019, 2020). Shaded grey fields show the range of cpx compositions in abyssal peridotites
(Warren, 2016), except where noted otherwise. **a.** Clinopyroxene rare earth element
concentrations for type-F Pleiku peridotites (solid blue lines), with Pleiku xenoliths from Nguyen
and Kil (2019) shown as solid (type-F) and dashed (type-R) gray lines for comparison. **b.**
Clinopyroxene rare earth element concentrations for Xuan Loc peridotites, where type-R
peridotites are shown by dashed and type-F peridotites by solid red lines; Dalat xenoliths from
Ang et al. (2021) and Nguyen and Kil (2020) are shown as solid (type-F) and dashed (type-R)
gray lines for comparison. **c.** Clinopyroxene trace element concentrations for Pleiku peridotite
xenoliths from this study; **d.** cpx trace element concentrations of Xuan Loc peridotites from this
study; and **e.** opx trace element concentrations of both Pleiku and Xuan Loc peridotites from this
study; the grey field represents the detection limits for each element.

Figure 4. **a.** $^{207}\text{Pb}/^{204}\text{Pb}$ and **c.** $^{208}\text{Pb}/^{204}\text{Pb}$ versus $^{206}\text{Pb}/^{204}\text{Pb}$; **b.** $^{206}\text{Pb}/^{204}\text{Pb}$ and **d.** ϵ_{Nd} versus $^{87}\text{Sr}/^{86}\text{Sr}$; and **e.** $^{206}\text{Pb}/^{204}\text{Pb}$ vs. ϵ_{Nd} for cpx from Pleiku (blue squares) and Xuan Loc (red circles) peridotites for this study (larger square and circle symbols). Uncertainty ranges (2σ) are smaller than symbol size. Small gray circles show Indian MORB compositions after Gale et al. (2013). Enriched mantle (EM) compositions are after Hofmann (2007), and the northern hemisphere reference line (NHRL) is after Hart (1984). Local basalt compositions from the Pleiku and Xuan Loc volcanic centers are from Hoang et al. (1996; 2013). Literature data for Pleiku and Dalat xenolith compositions (smaller symbols) are from Anh et al. (2021) and Nguyen and Kil (2019, 2020).

Figure 5. Spinel Cr# vs **a.** Al_2O_3 wt%, **b.** Yb_N , and **c.** [Ti] in cpx; **d.** ϵ_{Nd} in cpx vs. $(\text{Ce}/\text{Yb})_{\text{PM}}$ in cpx, and **e.** $^{143}\text{Nd}/^{144}\text{Nd}$ vs. Sm/Yb in cpx, for Vietnam xenoliths from this study and prior studies (Anh et al., 2021; Nguyen & Kil, 2019, 2020). Low Al_2O_3 , low Yb, and high sp Cr# characterize xenoliths that have undergone extensive melt extraction. The dashed line and tickmarks show the calculated degree of melting as a percentage, where the degrees of melting are a function of sp Cr# (Hellebrand et al., 2001) and Yb concentration (Johnson et al., 1990).

Figure 6. Comparison of equilibrium temperatures of Vietnam xenoliths with **a.** T_{36} and **b.** T_{REE} plotted versus T_{BKN} . The uncertainties (1σ) are $\pm 45^\circ\text{C}$ for T_{36} , $\pm 50^\circ\text{C}$ for T_{BKN} (Putirka, 2008), and as shown with error bars for T_{REE} . Black solid line is the 1:1 ratio, with the dotted lines representing $\pm 50^\circ\text{C}$.

Figure 7. Estimated equilibration temperatures (**a.** T_{36} and **b.** T_{BKN}) and **c.** measured $^{143}\text{Nd}/^{144}\text{Nd}$ isotopes, versus pressures (P_{38}) for xenolith samples from this study and prior studies (Anh et al., 2021; Nguyen & Kil, 2019, 2020). The calculated geothermal gradients beneath Pleiku and Xuan Loc, which were determined using measured lithospheric thickness and estimated heat flow, are also shown (using heat flow data after Hall (2002), and references therein).

Table 1. Sample locations for this study.

	Latitude (°N)	Longitude (°E)	Elevation (ft)	Xenolith type
Pleiku				
VN-2018-21-PL-1	13° 52.582'	108° 3.653'	2602	spinel lherzolite
VN-2018-21-PL-2	13° 52.582'	108° 3.653'	2602	spinel lherzolite
VN-2018-21-PL-3	13° 52.582'	108° 3.653'	2602	spinel lherzolite
VN-2018-21-PL-4	13° 52.582'	108° 3.653'	2602	spinel lherzolite
VN-2018-21-PL-5	13° 52.582'	108° 3.653'	2602	spinel lherzolite
VN-2018-21-PL-6	13° 52.582'	108° 3.653'	2602	spinel lherzolite
VN-2018-21-PL-7	13° 52.582'	108° 3.653'	2602	spinel lherzolite
VN-2018-21-PL-8	13° 52.582'	108° 3.653'	2602	spinel lherzolite
VN-2018-21-PL-9	13° 52.582'	108° 3.653'	2602	spinel lherzolite
Xuan Loc				
VN-2018-36-XL-1	10° 30.457'	107° 16.375'	229	dunite
VN-2018-36-XL-2	10° 30.457'	107° 16.375'	229	harzburgite
VN-2018-36-XL-3	10° 30.457'	107° 16.375'	229	spinel lherzolite
VN-2018-36-XL-4	10° 30.457'	107° 16.375'	229	harzburgite
VN-2018-36-XL-5	10° 30.457'	107° 16.375'	229	dunite
VN-2018-36-XL-7	10° 30.457'	107° 16.375'	229	spinel lherzolite
VN-2018-36-XL-8	10° 30.457'	107° 16.375'	229	spinel lherzolite
VN-2018-36-XL-9	10° 30.457'	107° 16.375'	229	spinel lherzolite
VN-2018-36-XL-11	10° 30.457'	107° 16.375'	229	dunite
VN-2018-36-XL-12	10° 30.457'	107° 16.375'	229	dunite
VN-2018-36-XL-13	10° 30.457'	107° 16.375'	229	harzburgite
VN-2018-36-XL-14	10° 30.457'	107° 16.375'	229	spinel peridotite
VN-2018-36-XL-15	10° 30.457'	107° 16.375'	229	spinel lherzolite
VN-2018-36-XL-16	10° 30.457'	107° 16.375'	229	spinel lherzolite
VN-2018-36-XL-18	10° 30.457'	107° 16.375'	229	spinel lherzolite
VN-2018-36-XL-19	10° 30.457'	107° 16.375'	229	spinel lherzolite
VN-2018-36-XL-20	10° 30.457'	107° 16.375'	229	spinel lherzolite

Table 2. Isotopic composition of Vietnam xenoliths.

	Type	$^{87}\text{Sr}/^{86}\text{Sr}$	$2\sigma^{**}$	$^{143}\text{Nd}/^{144}\text{Nd}$	$2\sigma^{**}$	$^{206}\text{Pb}/^{204}\text{Pb}$	$2\sigma^{**}$	$^{207}\text{Pb}/^{204}\text{Pb}$	$2\sigma^{**}$	$^{208}\text{Pb}/^{204}\text{Pb}$	$2\sigma^{**}$
Pleiku											
VN-2018-21-PL-1*	F	0.703267	±5	-	-	16.92	±1	15.45	±1	36.90	±3
VN-2018-21-PL-2	F	0.703365	±6	0.514190	±7	18.859	±2	15.579	±2	38.851	±4
VN-2018-21-PL-3*	F	0.702447	±6	0.513637	±7	16.69	±1	15.34	±1	36.48	±3
VN-2018-21-PL-4	F	0.704211	±6	0.512808	±4	18.33	±2	15.59	±2	38.50	±5
VN-2018-21-PL-5	F	0.702919	±5	0.513091	±4	18.591	±2	15.518	±2	38.579	±4
VN-2018-21-PL-6*	F	0.702425	±6	0.513776	±5	18.29	±1	15.49	±1	38.28	±3
VN-2018-21-PL-7*	F	0.702988	±7	-	-	17.39	±1	15.48	±1	37.30	±2
VN-2018-21-PL-8	F	0.702463	±6	-	-	18.1815	±2	15.560	±2	38.202	±5
VN-2018-21-PL-9*	F	0.702381	±5	0.513821	±5	18.27	±1	15.50	±1	38.27	±3
Xuan Loc											
VN-2018-36-XL-1	R	0.704050	±6	0.512800	±5	18.179	±2	15.548	±1	38.436	±4
VN-2018-36-XL-3	F	0.702645	±5	0.513196	±5	17.348	±2	15.461	±2	37.421	±4
VN-2018-36-XL-5	R	-	-	0.512755	±5	18.371	±2	15.572	±2	38.640	±5
VN-2018-36-XL-8	F	0.703724	±6	0.512921	±5	18.284	±2	15.541	±2	38.505	±5
VN-2018-36-XL-9	F	0.703162	±6	0.513036	±5	17.877	±2	15.494	±1	38.068	±3
VN-2018-36-XL-11	R	-	-	0.512755	±5	18.317	±2	15.570	±2	38.613	±5
VN-2018-36-XL-14	F	0.702952	±6	0.513176	±4	18.019	±2	15.512	±2	38.247	±5
VN-2018-36-XL-15	F	0.702565	±6	0.513358	±12	17.133	±1	15.435	±1	37.079	±3
VN-2018-36-XL-18	F	0.702995	±6	0.513087	±5	18.225	±1	15.529	±1	38.413	±3
VN-2018-36-XL-19	F	0.702967	±6	0.513371	±5	17.874	±2	15.509	±1	38.033	±3
VN-2018-36-XL-20	F	0.703070	±6	0.513110	±5	18.026	±2	15.520	±1	38.283	±3
Standards											
BCR-2	n = 2	0.705006	±8	0.512639	±5	18.756	±2	15.630	±2	38.740	±6
BHVO-2	n = 1	0.703472	±6	0.512987	±4	18.619	±2	15.542	±2	38.245	±5
BCR-2***	n = 5					18.76	±1	15.66	±1	38.77	±2
BHVO-2***	n = 5					18.61	±1	15.56	±1	38.20	±2

* Pb-isotopes measured via multi-Daly measurement with sample-standard bracketing

** Uncertainties are 2s internal standard error for the last digit expressed.

*** Pb-isotope measurement via multi-Daly measurement with sample-standard bracketing during same analytical session.

Table 3. Equilibrium temperatures (°C) and pressures (kbar) of Vietnam xenoliths. *

	Type	T _{BKN}	T ₃₆	T _{REE}	1 σ	P _{P38}
Pleiku						
VN-2018-21-PL-1	F	912	959	927	±47	11.7
VN-2018-21-PL-2	F	963	956	907	±10	15.8
VN-2018-21-PL-3	F	891	965	926	±25	13.4
VN-2018-21-PL-5	F	1067	1034	993	±30	13.6
VN-2018-21-PL-6	F	958	966	892	±13	15.1
VN-2018-21-PL-7	F	945	945	907	±40	13.1
VN-2018-21-PL-8	F	1000	1008	1032	±18	12.3
VN-2018-21-PL-9	F	967	972	887	±13	15.6
Xuan Loc						
VN-2018-36-XL-2	R	921	1006	-		9.6
VN-2018-36-XL-3	F	856	951	845	±59	12.2
VN-2018-36-XL-4	R	765	942	-		8.1
VN-2018-36-XL-5	R	795	923	1004	±94	7.1
VN-2018-36-XL-7	F	805	940	1076	±63	12.7
VN-2018-36-XL-8	F	859	935	1070	±37	12.1
VN-2018-36-XL-9	F	851	932	861	±49	13.2
VN-2018-36-XL-11	R	927	942	997	±70	10.0
VN-2018-36-XL-14	F	924	936	821	±25	15.3
VN-2018-36-XL-15	F	1026	992	843	±42	16.1
VN-2018-36-XL-18	F	907	927	908	±32	12.6
VN-2018-36-XL-19	F	946	959	916	±31	12.8
VN-2018-36-XL-20	F	894	929	885	±23	13.2

* Major element equilibrium temperatures calculated from the two-pyroxene thermometer (T_{BKN}) of Brey and Kohler (1990). T_{BKN} has an uncertainty (1 σ) of ± 50 °C. Temperatures are also calculated using the two-pyroxene thermometer of Putirka (2008) (T₃₆), with an uncertainty (1 σ) of ± 45 °C. Trace element equilibrium temperatures are calculated from the two-pyroxene REE thermometer (T_{REE}) of Liang et al. (2013); uncertainties reported are from linear fit of the inversion diagrams from that method (see Supplementary Information). Equilibration pressures (P_{P38}) are calculated using the two-pyroxene barometer of Putirka (2008), with an uncertainty of ± 3.7 kbar.




Figure 1. a. Map of southern and southeast Asia, illustrating extrusion of Indochina due to the India-Eurasia collision (gray arrows) (after Taponnier et al., 1986, 1990). "Red river" = Ailao Shan - Red River Fault Zone. **b.** Map of study area and sample collection localities. Major mapped faults and volcanic centers are modified from published literature (Hoang et al., 1996, 2013; Nguyen et al., 2012; Phach & Anh, 2018).




Figure 2. a. Clinopyroxene Al_2O_3 rim/core ratio versus cpx Cr_2O_3 rim/core ratio. **b.** Cpx core Al_2O_3 wt% vs. cpx Mg# (molar $\text{Mg}/(\text{Mg} + \text{Fe})$). Small symbols show published xenolith data from Pleiku (Nguyen & Kil, 2019) and Dalat (Anh et al., 2021; Nguyen & Kil, 2020).




Figure 3. Rare earth and trace element concentrations of samples from this study normalized to primitive mantle (Sun and McDonough, 1989). Gray solid lines show type-F xenoliths and gray dashed lines show type-R xenoliths from Pleiku and Dalat (Anh et al., 2021; Nguyen and Kil, 2019, 2020). Shaded grey fields show the range of cpx compositions in abyssal peridotites (Warren, 2016), except where noted otherwise. **a.** Clinopyroxene rare earth element concentrations for type-F Pleiku peridotites (solid blue lines), with Pleiku xenoliths from Nguyen and Kil (2019) shown as solid (type-F) and dashed (type-R) gray lines for comparison. **b.** Clinopyroxene rare earth element concentrations for Xuan Loc peridotites, where type-R peridotites are shown by dashed and type-F peridotites by solid red lines; Dalat xenoliths from Ang et al. (2021) and Nguyen and Kil (2020) are shown as solid (type-F) and dashed (type-R) gray lines for comparison. **c.** Clinopyroxene trace element concentrations for Pleiku peridotite xenoliths from this study; **d.** cpx trace element concentrations of Xuan Loc peridotites from this study; and **e.** opx trace element concentrations of both Pleiku and Xuan Loc peridotites from this study; the grey field represents the detection limits for each element.




Figure 4. **a.** $^{207}\text{Pb}/^{204}\text{Pb}$ and **c.** $^{208}\text{Pb}/^{204}\text{Pb}$ versus $^{206}\text{Pb}/^{204}\text{Pb}$; **b.** $^{206}\text{Pb}/^{204}\text{Pb}$ and **d.** ϵ_{Nd} versus $^{87}\text{Sr}/^{86}\text{Sr}$; and **e.** $^{206}\text{Pb}/^{204}\text{Pb}$ vs. ϵ_{Nd} for cpx from Pleiku (blue squares) and Xuan Loc (red circles) peridotites for this study (larger square and circle symbols). Uncertainty ranges (2σ) are smaller than symbol size. Small gray circles show Indian MORB compositions after Gale et al. (2013). Enriched mantle (EM) compositions are after Hofmann (2007), and the northern hemisphere reference line (NHRL) is after Hart (1984). Local basalt compositions from the Pleiku and Xuan Loc volcanic centers are from Hoang et al. (1996; 2013). Literature data for Pleiku and Dalat xenolith compositions (smaller symbols) are from Anh et al. (2021) and Nguyen and Kil (2019, 2020).




Figure 5. Spinel Cr# vs **a.** Al_2O_3 wt%, **b.** Yb_{N} , and **c.** [Ti] in cpx; **d.** ϵ_{Nd} in cpx vs. $(\text{Ce}/\text{Yb})_{\text{PM}}$ in cpx, and **e.** $^{143}\text{Nd}/^{144}\text{Nd}$ vs. Sm/Yb in cpx, for Vietnam xenoliths from this study and prior studies (Anh et al., 2021; Nguyen & Kil, 2019, 2020). Low Al_2O_3 , low Yb, and high sp Cr# characterize xenoliths that have undergone extensive melt extraction. The dashed line and tickmarks show the calculated degree of melting as a percentage, where the degrees of melting are a function of sp Cr# (Hellebrand et al., 2001) and Yb concentration (Johnson et al., 1990).




Figure 6. Comparison of equilibrium temperatures of Vietnam xenoliths with **a.** T_{36} and **b.** T_{REE} plotted versus T_{BKN} . The uncertainties (1σ) are ± 45 °C for T_{36} , ± 50 °C for T_{BKN} (Putirka, 2008), and as shown with error bars for T_{REE} . Black solid line is the 1:1 ratio, with the dotted lines representing ± 50 °C.




Figure 7. Estimated equilibration temperatures (**a.** T_{36} and **b.** T_{BKN}) and **c.** measured $^{143}\text{Nd}/^{144}\text{Nd}$ isotopes, versus pressures (P_{38}) for xenolith samples from this study and prior studies (Anh et al., 2021; Nguyen & Kil, 2019, 2020). The calculated geothermal gradients beneath Pleiku and Xuan Loc, which were determined using measured lithospheric thickness and estimated heat flow, are also shown (using heat flow data after Hall (2002), and references therein).