



Discovery based research experiences: gauche effects in group 16 elements.

HENRY RZEPA

READ REVIEWS

WRITE A REVIEW

CORRESPONDENCE:

h.rzepa@imperial.ac.uk

DATE RECEIVED:

March 02, 2016

DOI:

10.15200/winn.145691.11862

ARCHIVED:

March 02, 2016


CITATION:

Henry Rzepa, Discovery based research experiences: gauche effects in group 16 elements., *The Winnower* 3:e145691.11862, 2016, DOI: 10.15200/winn.145691.11862


© Rzepa This article is distributed under the terms of the [Creative Commons Attribution 4.0 International License](#), which permits unrestricted use, distribution, and redistribution in any medium, provided that the original author and source are credited.



The upcoming ACS national meeting in San Diego has a CHED (chemical education division) session entitled [Implementing Discovery-Based Research Experiences in Undergraduate Chemistry Courses](#). I had previously explored what I called *extreme gauche effects* in the molecule F-S-S-F. Here I take this a bit further to see what else can be discovered about molecules containing bonds between group 16 elements (QA= O, S, Se, Te).




The search definition is shown above, with DIST1 being the QA-QA bond length, the QA-QA bond being acyclic, each QA bearing only two bonded atoms and NM being any non-metal. The first result shown is for QA=S.




1. The first discovery is that the most common torsion (red-hot spot) is about 90° , but there appears to be a statistically significant distortion towards **longer** S-S distances as the torsion deviates from this angle. For those who are so inclined it would perhaps be worth improving my term "*appears to be*" with a more formal numerical analysis of the distribution shown above and its significance. Any offers?
2. The other discovery worth exploring is the number of occurrences with an angle of 180° . With F-S-S-F itself (not a solid), I had previously *noted* that this angle actually represented a transition state in the torsion! So what might be inferred from these examples?

The next search includes a further constraint that the temperature the data was recorded at be $<140\text{K}$. This reduces vibrational "noise" and so should increase the significance.




1. Here we discover the same "V"-shaped distribution as before, possibly more significant statistically than the previous search. Again, a proper statistical analysis of the significance of this result is desirable.

The next search is for QA = Se or Te.



1. The Se and Te distributions can clearly be distinguished, with a weak "V-shape" visible for Se, but absent for Te. Again, those hits at 180!
2. There are a few instances "in-between" the two distributions, which appear to be Se-Te systems. Finally, QA=QB = O.



1. The discovery here is the apparent **absence** of any "V-shaped" distribution.
2. The hot spot now occurs at 180° , but with a tail down to 60° or less. Clearly, the definition of "NM" as any non-metal probably needs to be explored further for specific instances to see what influence the nature of NM has. NM for example could be another O, which might be a severe perturbation. So here I have tried to tease out seven directions for further discovery. I am attending/presenting at the session I noted at the top and will report back on any interesting observations.