

Exploring potential atmospheric methane removal approaches: an example research roadmap for chlorine radical enhancement

Katrine A. Gorham¹, Sam Abernethy¹, Tyler R. Jones², Peter Hess³, Natalie M. Mahowald⁴, Daphne Meidan⁵, Matthew S. Johnson⁶, Maarten M. J. W. van Herpen⁷, Yangyang Xu⁸, Alfonso Saiz-Lopez⁵, Thomas Röckmann⁹, Chloe Brashear⁹, Erika Reinhardt¹, David Mann¹

¹ Spark Climate Solutions, Covina, CA, USA

² Institute of Arctic and Alpine Research, University of Colorado, Boulder, CO, USA

³ Department of Biological and Environmental Engineering, Cornell University, Ithaca, NY, USA

⁴ Department of Earth and Atmospheric Sciences, Cornell University, Ithaca, NY, USA

⁵ Department of Atmospheric Chemistry and Climate, Institute of Physical Chemistry Blas Cabrera, CSIC, Madrid, Spain

⁶ Department of Chemistry, University of Copenhagen, Copenhagen, Denmark

⁷ Acacia Impact Innovation BV, Bernheze, The Netherlands

⁸ Department of Atmospheric Sciences, Texas A&M University, College Station, TX, USA

⁹ Institute for Marine and Atmospheric Research Utrecht, Department of Physics, Utrecht University, Utrecht, The Netherlands

Keywords: climate intervention, methane removal, iron salt aerosols, ISA, chlorine radical, roadmap

Abstract:

The escalating climate crisis requires rapid action to reduce the concentrations of atmospheric greenhouse gases and lower global surface temperatures. Methane will play a critical role in near-term warming due to its high radiative forcing and short atmospheric lifetime. Methane emissions have accelerated in recent years and there is significant risk and uncertainty associated with the future growth in natural emissions. The largest natural sink of methane occurs through oxidation reactions with atmospheric hydroxyl and chlorine radicals. Enhanced atmospheric oxidation could be a potential approach to remove atmospheric methane. One method proposes the addition of iron salt aerosols (ISA) to the atmosphere, mimicking a natural process that is proposed to occur when mineral dust mixes with chloride from sea spray to form iron chlorides, which are photolyzed by sunlight to produce chlorine radicals. Under the right conditions, lofting ISA into the atmosphere could potentially reduce atmospheric methane concentrations and lower global surface temperatures. Recognizing that potential atmospheric methane removal must only be considered as an additive measure – in addition to, not replacing, crucial anthropogenic greenhouse gas emission reductions and carbon dioxide removal – roadmaps can be a valuable tool to organize and streamline interdisciplinary and multifaceted research to efficiently move towards an understanding of whether an approach may be viable and socially acceptable, or if it is nonviable and further research should be deprioritized. Here we present an example of a five-year research roadmap to explore whether ISA enhancement of the chlorine radical sink could be a viable and socially acceptable atmospheric methane removal approach.

1. Introduction

Driven by anthropogenic greenhouse gas emissions, Earth's average surface temperature has increased by 1.1 °C since 1850 (Forster et al., 2021). As global temperature increases, irreversible changes in the Earth system will likely occur, such as ice sheet collapses in Greenland and Antarctica, coral reef die-off, abrupt permafrost thaw, mountain glacier loss, and Amazon rainforest dieback (Lenton et al., 2008; McKay et al., 2022). To mitigate both near-term and long-term warming we must undertake rapid, sustained action to establish a diverse portfolio of approaches to slow and then reverse the increase of atmospheric greenhouse gas concentrations, ideally reducing them to preindustrial levels. Emissions reductions must be prioritized. In addition, as near-term warming threatens to trigger climate tipping points, negative emissions approaches may be used to remove greenhouse gases already in the atmosphere and counter rising natural and uncontrollable emissions

Methane (CH_4) has contributed 0.5 °C to the current 1.1 °C temperature increase relative to preindustrial times, second only to carbon dioxide (CO_2) (IPCC, 2023). Methane lasts roughly a decade in the atmosphere (UNEP, 2021), with a Global Warming Potential 83 times that of carbon dioxide over 20 years and 30 times that of carbon dioxide over 100 years (Forster et al., 2021). The concentration of methane in the atmosphere is 2.5 times pre-industrial levels, and the growth rate has accelerated since 2006, with record increases in 2020 and 2021 (NOAA, 2023). Methane emissions come from anthropogenic sources (fossil fuel use, agriculture, waste and wastewater, biomass burning, etc.) and natural sources (wetlands, oceans, freshwaters, termites, permafrost, etc.), both of which are increasing (Jackson et al., 2020; Nisbet et al., 2023; Saunio et al., 2020). As the planet continues to warm and precipitation patterns change, natural methane emissions are expected to increase from wetlands, as well as from permafrost due to abrupt thaw, thermokarst lake formation and expansion, and bacterial processes (Dean et al., 2018; Neumann et al., 2019; Paudel et al., 2016; Peng et al., 2022; Zhang et al., 2023). One estimate is that natural emissions could increase by 20% (~50 Tg CH_4 /yr) to 150% (~330 Tg CH_4 /yr) by 2100, becoming larger than anthropogenic emissions in most modeled scenarios (Kleinen et al. 2021). There is evidence that wetland emissions increases are already underway, with roughly half of the 2020 atmospheric methane increase attributed to wetlands (Peng et al., 2022; Nisbet et al., 2023; Zhang et al., 2023).

The natural sinks for atmospheric methane are oxidation by gas phase radicals (~95 %) and uptake into soil by methanotrophic bacteria and archaea (~5 %) (Saunio et al., 2020). The atmospheric lifetime of methane is thus mainly determined by the oxidative capacity of the atmosphere. Approximately 550 Tg methane per year is removed by reactions with hydroxyl radicals ($\text{OH}\cdot$); chlorine radicals ($\text{Cl}\cdot$) destroy ~11 Tg methane per year (Saunio et al., 2020). However, the chlorine radical sink estimate is poorly constrained, ranging from 1 to 35 Tg methane per year (Saunio et al., 2020). Recent research suggests the atmospheric chlorine radical sink could be underestimated (van Herpen et al., 2023).

Complex and non-linear atmospheric chemistry dictates the oxidation of methane, which results in the production of formaldehyde and carbon monoxide which is further oxidized to carbon dioxide. Hydroxyl radical oxidation of methane can result in the formation or loss of ozone, depending on the nitrogen oxide (NO_x) concentrations. Chlorine radicals react to remove both methane and ozone, the principal

precursor of hydroxyl radical production (Lelieveld et al., 2002; Seinfeld and Pandis, 2016). Therefore, chlorine radicals may either increase or decrease the atmospheric lifetime of methane depending on the concentration of chlorine radicals and atmospheric conditions (Li et al., 2022, van Herpen et al., 2023, Li et al., 2023). Volatile organic compounds and NO_x will change the methane response since they directly and indirectly affect the concentrations of hydroxyl and chlorine radicals. Furthermore, since reactions with carbon monoxide and methane are the primary sinks for hydroxyl radicals, the abundance of methane impacts the oxidative character of the atmosphere in a self feedback process (Lelieveld et al., 2002; Staniaszek et al., 2022).

Enhanced atmospheric oxidative sinks could increase the rate of atmospheric methane removal and therefore reduce near-term warming. One untested proposal involves iron salt aerosols (ISA). This potential approach involves lofting iron-based particles into the troposphere (e.g., from ships or towers) to catalytically produce chlorine radicals (Oeste, 2009; Oeste et al., 2017), mimicking a natural phenomenon proposed to occur when mineral dust combines with sea spray aerosols (van Herpen et al., 2023). Discussing natural analogues of this process and the current state of research, this paper presents a roadmap for research and development that is needed to understand whether ISA enhancement of the chlorine radical sink may be a feasible, scalable, and safe approach for atmospheric methane removal.

2. State of the Research

Currently, research into ISA falls into three categories: laboratory experiments to quantify the details of the mechanism, observational analysis of the natural analogue of ISA, known as mineral dust sea spray aerosol (MDSA), and numerical modeling evaluating the potential impacts of ISA.

a. Laboratory experiments

In a series of papers, Wittmer et al. demonstrated the production of chlorine atoms from iron-doped salts and aerosols (Wittmer et al., 2015a; Wittmer et al., 2015b; Wittmer et al., 2017). Reproducing and expanding upon these laboratory studies is of the highest importance, since the mechanism of chlorine generation is poorly understood (van Herpen et al., 2023; Wittmer et al., 2015; Wittmer et al., 2017; Zhu et al., 1997). The ISA mechanism is catalytic in iron and may be catalytic in chlorine in natural environments (Wittmer et al., 2017). Understanding of catalytic efficiency requires study of dependencies on conditions such as aerosol size distribution and number density; humidity; rate of supply of acidity to the system; rate of coagulation of aerosols; effects of organic chelating agents on iron activity; changes in chlorine escape probability due to aqueous chemical conditions; changes in iron activity due to aerosol chemistry; and, understanding how the atmosphere behaves under high ClO_x conditions (Pennacchio et al., submitted). Furthermore, studies of real atmospheric conditions are needed to better understand suppression of the ISA mechanism in the presence of sulfate and oxalate (Wittmer et al., 2015; Wittmer et al., 2017).

b. The natural analogue of ISA: mineral dust sea spray aerosol

The natural mineral dust sea spray aerosol (MDSA) mechanism is proposed to occur when iron from mineral dust mixes with the chlorine in sea spray, forming iron chlorides which are photolyzed to produce chlorine radicals (van Herpen et al., 2023). ISA could mimic this natural MDSA phenomenon by only aerosolizing what is currently believed to be the key component of the mineral dust: the photoactive iron.

van Herpen et al. (2023) provided the first evidence that methane may be removed over the North Atlantic by chlorine radicals produced by the MDSA mechanism. The semi-arid regions of North Africa are the dominant source of iron-containing mineral dust, with frequent transport over the North Atlantic (Prospero et al., 2021). Using air samples collected during North African dust events from 1996-2000 in Barbados (Mak et al., 2003), a model parameterized with the MDSA mechanism of chlorine radical production produced results consistent with a previously unexplained ^{13}C -depletion in the reaction product carbon monoxide (CO) (van Herpen et al., 2023). Carbon monoxide produced from chlorine radical oxidation of methane is extremely depleted in ^{13}C which makes $\delta^{13}\text{C}(\text{CO})$ a very sensitive indirect detection method of chlorine radicals (Röckmann et al., 1999). The variability of $\delta^{13}\text{C}(\text{CO})$ in atmospheric air is the main evidence that the MDSA process is occurring, as there is no other mechanism proposed that can explain the carbon monoxide isotope signature (Mak et al., 2003).

As a proxy for methane oxidation by chlorine radicals, studies of the isotopic composition of carbon monoxide in the mid-Atlantic boundary layer are underway to further confirm the MDSA mechanism. This includes a regular air sampling program at four ground-based stations in the North Atlantic, as well as Atlantic transect sampling onboard commercial vessels. Time series at the ground-based stations provide high-resolution observations at the longitudinal margins of trans-Atlantic dust transport, including the site where seasonal depletions in $\delta^{13}\text{C}(\text{CO})$ were previously observed (Mak et al., 2003). Air samples collected in north-south ship transects may allow spatial correlation of the $\delta^{13}\text{C}(\text{CO})$ excursions with North African dust plumes. Together, these samples provide an opportunity to investigate the formation process of MDSA, as well as the seasonal and spatial influence of North African dust on tropospheric chlorine radical oxidation.

c. Modeling

Early modeling studies indicate that atmospheric chlorine additions can increase or decrease methane concentrations, depending on the concentration of chlorine that is present in the atmosphere (Horowitz et al., 2020; Li et al., 2022; Saiz-Lopez et al., 2023). Chlorine radicals readily oxidize ozone, the main precursor of hydroxyl radicals, which results in less methane oxidation via hydroxyls; at low atmospheric concentrations, chlorine radicals reduce ozone concentrations without having a commensurate impact on methane. Even though chlorine radicals react 16x faster with methane than the reaction of hydroxyl radicals with methane (Atkinson, 2006), hydroxyl radicals dominate the methane oxidation sink because they are much more abundant than chlorine radicals. As more chlorine is emitted, ozone concentrations will be reduced so that proportionally more chlorine radicals react with methane. The increased methane destruction by chlorine radicals will eventually outcompete decreased destruction by hydroxyl. In an initial, highly simplified model scenario, Li et al. (2023) found that a reduction in methane concentration could be achieved if more than 90 Tg Cl/yr (three times the estimated present-day emission rate) was added evenly to the atmosphere over all ocean surfaces. However, assuming a uniform increase of chlorine over all ocean surfaces may underestimate the potential effectiveness of local chlorine radical generation where efficiency may be condition-dependent (e.g., NO_x , CO, and chlorine concentrations, humidity, temperature, altitude, etc.) (Meidan et al., submitted). For example, considering the MDSA natural analogue of ISA, van Herpen et al (2023) found that high dust concentrations in the North Atlantic corresponded with net methane removal, while globally lower dust concentrations led to a net increase of methane.

Current models may not accurately capture the speed and efficiency of producing chlorine radicals via the ISA mechanism due to assumptions of the percentage of photoactive iron in the emitted iron, aerosol pH, aerosol mixing rates, and more. Another challenge with Earth system models is that they instantaneously dilute emissions to model grid dimensions which could lead to underestimates or overestimates of the effectiveness of the ISA mechanism, especially when considering iron emission additions from point sources like ships (e.g. Meidan et al., submitted). For example, the mixing of the iron and sea salt within the aerosol is modeled to occur instantaneously (Meidan et al., submitted; van Herpen et al., 2023); however, in reality it would likely take hours to days, leading the global model to overestimate the rate of chlorine radical production. Furthermore, the ISA mechanism is likely to occur faster in high NO_x environments (Oeste et al., 2017) but could be less efficient in high sulfate environments (Bondy et al., 2017; Chen et al., 2020; Legrand et al., 2017; Pio et al., 1998), and both NO_x and sulfate may be co-emitted with iron (e.g. from a ship plume). Thus, models that instantaneously dilute emissions across the grid dimensions may misrepresent the ISA mechanism. Overall, it is unclear whether current Earth system models overestimate or underestimate the efficiency of the oxidation mechanism. Additional detailed box modeling focusing on deployment sites and constrained by field observations are necessary to assess the effectiveness of the mechanism. However, local box models are less reliable over timescales where mixing between different air masses is relevant. This motivates the need for high-resolution regional and seasonal modeling over ocean basins and variable resolution configurations embedded in global models.

Considering the difficulty of simulating variable atmospheric chemical conditions (e.g., atmospheric chemical composition, solar radiation, wind mixing, etc.) across different geographic locations, it is important to develop an ensemble of models that enable uncertainty assessment. Such a model ensemble will allow a comprehensive exploration of different iron salt aerosol deployment scenarios (magnitude each year), aerosol particle sizes, and deployment location and timing.

3. Roadmap

a. Roadmap framework

Roadmaps are used in climate research to define knowledge gaps, needs, and associated outputs as they relate to interdependencies and timelines (e.g. Wanser et al., 2022; Maesano et al., 2022; Aschwanden et al., 2021; Ocean Visions, 2023), particularly in instances that benefit from integrated, interdisciplinary research. A coordinated, thorough, and science-based approach is needed to ensure that resources are used efficiently, stakeholders and interdisciplinary teams are engaged on appropriate timelines, and efforts are focused towards sequenced research questions and milestones.

b. Viability assessment

The viability of an atmospheric methane removal approach can be assessed by considering its potential for feasibility, scalability, and social license to operate. A feasible approach must be climate beneficial, safe, acceptable for its side effects, and cost-plausible. Determination of scalability will be approach-specific, acknowledging that the scale of increased natural emissions is anticipated to be tens of millions of tons of methane per year (Kleinen et al., 2021).

The key milestone questions below can help determine the viability of ISA and whether it should continue to be prioritized. The research that informs the key milestones questions should be pursued in parallel (Table 1).

1. Is enhancement of the chlorine radical oxidative sink of methane via the ISA mechanism effective and climate beneficial? At what scale?
2. What impacts could the ISA approach have on Earth systems and human systems, both positive and negative? Is there a cost-plausible ISA deployment method?
3. What is needed to advance a structure of ethical governance and social license for utilizing atmospheric intervention to reduce atmospheric methane concentrations?

Milestone question #1: Is the ISA mechanism effective and climate beneficial and scalable?

The complexity and nonlinearity of atmospheric chemistry and meteorology requires laboratory, field, and plume and global modeling studies of the efficiency of chlorine radical production, its dependence on atmospheric conditions and other gases, and the impact on methane removal.

An important assumption in previous studies (van Herpen et al., 2023; Meidan et al., submitted) is that only 2% of iron is photoactive (van Herpen et al., 2023; Meidan et al., submitted). However, the amount of ISA that is photoactive may vary by emission source (e.g. ship emissions may have more photoactive iron relative to mineral dust; Ito, 2013; Rodriguez et al., 2021), geographical location, aerosol pH, the presence of other chemical constituents (e.g. sulfate and NO_x), and altitude (Mahowald et al., 2018). The efficiency, cost, safety (e.g. air quality), and net radiative forcing of ISA will depend on the percentage of iron that is photoactive.

Current studies assume that the chlorine radicals released from the photochemical reaction with iron will react (e.g., with methane) to form hydrochloric acid, which will then be reabsorbed back into the aerosol and thus recycled (van Herpen et al., 2023; Oeste et al., 2017). It is unclear under which atmospheric conditions this cycle occurs, but if some chlorine radicals are lost (e.g. to reactions with ozone instead), then the cycling would be less efficient. After reacting with ozone there is a much longer chain of reactions before hydrochloric acid is created, thus reducing the cycling efficiency (Wang et al., 2019). Therefore, further laboratory measurements and detailed box models are needed to further study hydrochloric acid recycling efficiency by ISA.

Better understanding of how sulfur dioxide and NO_x concentrations impact the ISA mechanism is also needed (Oeste et al., 2017). Sulfur dioxide and NO_x concentrations vary regionally and locally and their emissions may sometimes coincide with those of iron.

Smaller ISA particles have greater surface area to mass ratios and stay longer in the atmosphere, likely increasing the efficiency of the ISA mechanism. Furthermore, smaller particles tend to be more acidic (Pye et al., 2020), which is required for the mechanism to be active (Wittmer et al., 2015; 2017). However, smaller aerosols may be transported further to coastal or inland locations where these particles could contribute directly to negative human health effects or deposit on terrestrial or ice-covered surfaces with unintended consequences. The aerosol size may also affect marine cloud cover, thereby influencing local radiative forcing. In addition, there tends to be more sulfate in smaller aerosols, which may reduce the effectiveness of the ISA mechanism (Bondy et al., 2017; Chen et al., 2020; Legrand et al., 2017; Pio et al., 1998).

Studying MDSA (the natural analogue of ISA) through field studies is essential to understand the effectiveness of this mechanism under different atmospheric conditions and its geographical extent, thereby better constraining atmospheric and Earth system models. Early MDSA field studies are underway to explore the seasonality and spatial extent of methane oxidation by chlorine radicals that may occur in natural dust plumes through proxy measurements of $\delta^{13}\text{C}(\text{CO})$. Further studies – both natural analogue and in-situ ISA enhancement studies – would benefit from alternative ISA detection and quantification approaches, including direct chlorine measurements or additional proxy measurements to reinforce existing observations.

Ideally, models will be developed to include the entire MDSA mechanism, including implementation of the isotope effect in the chlorine radical reaction with methane, thus enabling direct comparison of model results to observations of $\delta^{13}\text{C}(\text{CO})$. At present, some models (EMAC; Gromov et al., 2014) include complete carbon monoxide isotope representation, but not the MDSA mechanism, whereas other models (CAM-Chem; van Herpen et al., 2023) include an initial representation of the MDSA mechanism but do not incorporate the isotopic effects.

Isotopic signatures and dust from ice core paleo records may elucidate evidence of historical MDSA activity. Methane isotope measurements of air trapped in polar ice cores have been used to constrain the methane budget in the past (Bock et al. 2017; Sapart et al., 2012; Fischer et al., 2008; Ferretti et al., 2005), but possible variations in the chlorine-based methane sink have not been taken into account in these studies. Dust levels have undergone strong changes in the past (e.g. Fischer et al., 2007; Han et al., 2018; Yuan et al., 2020; Yue et al., 2023), and associated changes in MDSA may have affected the paleo records of $\delta^{13}\text{C}(\text{CH}_4)$.

If modeling, laboratory studies, and natural analogue field studies prove that a promising, safe mechanism exists to produce chlorine radicals at sufficient scale and under diverse atmospheric conditions, it may be appropriate to consider field studies with intentional enhancement of ISA, using a broad suite of atmospheric measurements to understand how the mechanism performs in the real atmosphere. Prior to performing any ISA enhancement field studies – even at a small and controlled scale – it is essential to engage and work collaboratively with potentially impacted communities, policy and science leaders, governmental bodies, NGOs, media, and other stakeholders to ensure that actions are conducted with social license and an appropriate governance framework with free, prior, and informed consent (UNFAO, 2016). For example, a study could involve controlled enhancement of dust or emitted aerosolized iron, or could investigate existing anthropogenic emissions of iron (e.g. from a ship plume, power plant, iron foundry, etc). The scale of the study should be suitable to accommodate a likely non-linear atmospheric response, where substantial increases in chlorine – thus iron emissions – may be needed before there is a decrease in methane.

Milestone question #2: What are the potential Earth system and human systems impacts of ISA, and is it cost-plausible?

Lifecycle analyses are necessary to assess the potential benefits, tradeoffs, risks, uncertainties, and costs of ISA. As part of this analysis, understanding the potential impacts of ISA enhancement on the Earth system and human system is imperative before considering large scale deployment. Human system impacts may include human health outcomes, as well as indirect human impacts from Earth system

changes. For example, if ISA resulted in ocean acidification there could be marine life implications which could have far reaching ecosystem impacts, as well as economic (e.g. fisheries) and cultural (e.g. Indigenous practices) impacts for coastal communities.

Earth system modeling must be conducted to understand the impact of atmospheric conditions, aerosol size, and release locations, magnitudes, and timing, on ocean systems, terrestrial systems, the cryosphere, the stratosphere, and clouds and precipitation. Results from atmospheric modeling (plume and global), laboratory studies, and initial field studies will inform priorities and research directions for Earth system studies and could potentially inform the design of field studies to verify model results.

The potential broader impacts of ISA deployment beyond removing methane are not well understood. These may include co-benefits such as reductions in atmospheric black carbon (Li et al., 2021, Oeste et al., 2017) and ozone (Li et al., 2023; van Herpen et al., 2023), both of which are climate warming agents. However, there are also potential negative impacts that must be further explored including ocean acidification, stratospheric ozone loss, adverse chemical side reactions (such as COCl_2 formation), and albedo changes from deposition on ice surfaces (Li et al., 2023). Iron aerosols absorb light and thus tend to warm the planet, offsetting some of the lowered radiative forcing from oxidized methane (Matsui et al., 2018; Li et al., 2021; Meidan et al., submitted). There are also potential side effects such as indirect radiative forcing due to marine cloud brightening and carbon dioxide absorption by ocean iron fertilization (Emerson, 2019) that could be either favorable or unfavorable.

The potential effects of ISA on air quality and human health are also poorly understood. Enhanced chlorine could lead to beneficial reductions in tropospheric ozone. However, the reduced hydroxyl radical production may increase lifetimes of volatile organic compounds that are detrimental to human health. Moreover, iron aerosols themselves present a human health risk, especially when small (O'Day et al., 2022). Therefore, more research is required to determine under which conditions (including deployment location/timing, particle composition, and ISA size) the air quality impact is beneficial or detrimental.

Engineering deployment modalities and implementation scenarios is one of the later steps in this roadmap, only to be pursued if earlier dependencies are addressed and ISA proves effective and climate beneficial with acceptable side effects. Nevertheless, to avoid delaying potential future deployment-readiness and to iteratively refine design in advance of any in-situ field studies, development and engineering of a nozzle sprayer delivery system could begin in parallel with early research activities.

As part of a lifecycle analysis, the cost of materials (e.g., iron), infrastructure, and other implementation resources must be assessed. To be cost-plausible, these costs at scale must be lower than the social cost of methane, a monetary valuation that estimates the socioeconomic impact caused by an additional metric ton of methane (Azar et al., 2023).

Milestone question #3: What is needed to advance ethical governance and social license?

Policy and social science aspects of this roadmap must occur in parallel with the research and technical activities and should be iterative, where outcomes from policy and social science efforts inform development of research and technical direction, and vice versa. A misaligned or siloed approach may

result in outcomes where physical science results are not answering the social science questions to understand if there could be a safe, cost-plausible deployment method and pathway to ethical governance, which may stall or halt progress. Conversely, detrimental outcomes could also be realized if research and technical aspects were delayed.

It is imperative that governance and social impacts be considered for any atmospheric methane removal approach. Addressing the climate crisis requires engagement beyond technical solutions; collaboration and transparency between scientists, behavioral scholars, media, the public, policy-makers, NGOs, Indigenous peoples, and other stakeholders is essential to ensure that an ethical governance framework is established (Dowell et al., 2020; Diamond et al., 2022; Scott-Buechler et al., 2023; Carbon180, 2022). There must be effective engagement and education to co-create research questions and iteratively communicate research findings and results, as well as risks and co-benefits. Failure to do so jeopardizes the trust and sound decision making of communities and governments (The Arctic Institute, 2021), threatening our ability to critically and openly assess potential climate solution approaches through a scientific process. Ideally, an external governance framework is developed which is enforceable and legally binding; however, there is also value in internal governance frameworks which may, for example, be based around a code of conduct, advisory or review boards, or other non-binding structures.

If ISA remains promising, potential implementation scenarios could be considered. Some methane removal solutions may be deployed immediately, whereas others may be higher risk and may only be suitable for emergency deployment.

Table 1: Key research and development activities for a five-year timeline beginning in 2023. Funding is needed for all research and development activities, including those that are already underway.

Category	Research and development activity	Year in which work becomes a funding priority	Anticipated duration (variable depending on outcomes)	Academia	Non-profits	Government	Engineering consultant
Key milestone question #1: Is enhancement of the chlorine oxidative sink of methane via the iron salt aerosol (ISA) mechanism effective and climate beneficial? At what scale?							
Modeling studies	Atmospheric plume and high-resolution regional modeling to understand if ISA-driven chlorine radical production results in net methane loss using various ISA sizes and concentrations of iron, NO _x , and other chemical species.	underway	3 years	✓	✓	✓	
	Global atmospheric modeling to understand if ISA-driven chlorine radical production results in net cooling, considering radiative forcing from methane, ozone, and iron aerosols, the photoactivity of the iron, meteorological and atmospheric chemistry conditions, and ISA size.	underway	4 years	✓	✓	✓	
	Paleo modeling to understand variability in the chlorine-based methane sink that may have been associated with past dust changes.	2024	2 years	✓	✓	✓	
Laboratory studies	Chamber studies to understand the photoactive iron fraction, chlorine radical production efficiencies, methane oxidation efficiencies, and reactions with other species	underway	3 years	✓	✓	✓	

Field studies	Natural analogue studies to (a) better constrain the atmospheric radical sinks; (b) understand the spatial extent and magnitude of the natural analogue, MDSA; and (c) understand the sources and distribution of photoactive iron (from MDSA as well as combustion iron sources)	underway	3 years	✓	✓	✓	
	In-situ ISA enhancement field studies (e.g. ship plume and/or ground based) at a controlled experiment site to measure changes in chlorine, methane, and other species	2025	2+ years	✓	✓	✓	
Engineering	Initial design study for ISA sprayer system.	2024	1 year	✓			✓
Key milestone question #2: What impacts could this approach have on Earth systems and human systems, both positive and negative? Is there a cost-plausible deployment method?							
Modeling studies	Modeling of ocean system impacts considering different release locations/timing, ISA sizes, and atmospheric conditions.	2024	3+ years	✓	✓	✓	
	Modeling of ISA deposition impacts on different land and ice surfaces considering different release locations/timing, ISA sizes, and atmospheric conditions.	2024	3+ years	✓	✓	✓	
	Modeling of atmospheric impacts including clouds, precipitation, and stratospheric ozone considering different release locations/timing, ISA sizes, and atmospheric conditions.	2024	3+ years	✓	✓	✓	
	Lifecycle analysis to quantify ISA climate benefits, tradeoffs, and cost.	2024	Ongoing	✓	✓	✓	
Governance and social impacts	Study of potential human health impacts considering different release locations/timing, ISA sizes, and atmospheric conditions.	2025	3 years	✓	✓	✓	

Engineering	Advanced design study for ISA sprayer system.	2026	1 year	✓			✓
	Design study of deployment modalities and implementation scenarios.	2027	2 years	✓			✓
Key milestone #3: What is needed to advance a structure of ethical governance and social license for utilizing atmospheric intervention to reduce atmospheric methane concentrations?							
Governance and social impacts	Engage with stakeholders regarding the research findings and results, risks, and potential.	underway	Ongoing	✓	✓	✓	
	Develop a collaborative governance framework to monitor and report on environmental and social impacts.	2024	Ongoing		✓	✓	
	Stakeholder engagement for identification of potential deployment locations and strategies.	2026	3 years			✓	✓

c. Priorities and timeline

It may be advantageous to pursue multiple research questions in parallel because the output from one research question may inform the inputs for other research questions. As such, activities can be sequenced using a prioritized timeline (Figure 1), where later research activities and action areas often have multiple dependencies on earlier activities. For example, the engineering design study is suggested to begin in year 4 (2027) because it can start prior to having complete Earth system modeling results, human health study outcomes, or conclusions from ISA enhancement field studies. However, the engineering design study cannot advance to its later stages until earlier activities have been thoroughly addressed. This expedited schedule advances possible timelines to avoid delaying potential deployment-readiness, but is not meant to accelerate the timeline beyond appropriate caution and due diligence. Setbacks in addressing early research questions and action areas will likely result in timeline delays.




Figure 1: Roadmap of ISA research and development needs. Duration of research and development timelines may be longer or shorter than depicted and/or exceed the five-year horizon.

4. Conclusion

The activities outlined in this roadmap require coordinated efforts across multiple government agencies to financially support research and development, to ensure robust assessment and governance processes, and to foster international engagement. This work is valuable for multiple reasons:

1. We need to understand if ISA is a feasible, scalable, and safe methane removal method, or if it is nonviable and further research should be deprioritized.
2. Though this roadmap is ISA-specific, the research and development needs identified here contribute to fundamental understanding of processes and mechanisms that are broadly applicable to exploration of other methane removal approaches.
3. The research outlined in this roadmap will contribute to constraining the global methane budget and oxidative character of the atmosphere, which will improve our understanding of atmospheric chemistry, Earth system dynamics, and air quality.

Addressing the climate crisis requires a diverse portfolio of climate solutions. It is essential that atmospheric methane removal approaches are considered in addition to, not replacing, crucial anthropogenic greenhouse gas emission reductions and carbon dioxide removal. All atmospheric methane removal approaches are at a very early stage (Jackson et al., 2021; Ming et al., 2022; Spark, 2023); all require further research and none are ready for deployment. We hope that this ISA roadmap, and other atmospheric methane removal roadmaps that follow, will help accelerate, prioritize, and parallelize research that is essential to understanding which climate solutions to pursue.

References

- Aschwanden, A., Bartholomäus, T., Brinkerhoff, D.J., Truffer, M. (2021) Brief communication: A roadmap towards credible projects of ice sheet contribution to sea level. *The Cryosphere*, 15, 5705-5715.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., ... & Subcommittee, I. U. P. A. C. (2006). Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II—gas phase reactions of organic species. *Atmospheric chemistry and physics*, 6(11), 3625-4055.
- Azar, C., Martin, G.J., Johansson, D.J.A., Sterner, T. (2023) The social cost of methane. *Climatic Change*, 176, 71.
- Bock, M., Schmitt, J., Beck, J., Seth, B., Chappellaz, J., Fischer, H. (2017) Glacial/interglacial wetland, biomass burning, and geologic methane emissions constrained by dual stable isotopic CH₄ ice core records. *Proceedings of the National Academies of Sciences*, 114, E5778-E5786.
- Bondy, A.L., Wang, B., Laskin, A., Craig, R.L., Nhliziyo, M.V., Bertman, S.B., Pratt, K.A., Shepson, P.B., Ault, A.P. (2017) Inland Sea Spray Aerosol Transport and Incomplete Chloride Depletion: Varying Degrees of Reactive Processing Observed during SOAS. *Environmental Science and Technology*, 51, 17, 9533-9542.
- Carbon180. Setting DAC on Track. (2022) Available online at: <https://static1.squarespace.com/static/5b9362d89d5abb8c51d474f8/t/6261d1890b76863f1047a2dd/1650577901659/Carbon180-SettingDAConTrack.pdf> (Accessed August 15, 2023).
- Chen, Y., Cheng, Y., Ma, N., Wei, C., Ran, L., Wolke, R., Grob, J., Wang, Q., Pozzer, A., Denier van der Gon, H.A.C., Spindler, G., Lelieveld, J., Tegen, I., Su, H., Wiedensohler, A. Natural sea-salt emissions moderate the climate forcing of anthropogenic nitrate. (2020) *Atmospheric Chemistry and Physics*, 20, 2, 771-786.
- Dean, J.F., Middelburg, J.J., Rockmann, T., Aerts, R., Blauw, L.G., Egger, M., Jetten, M.S.M., de Jong, A.E.E., Meisel, O.H., Rasigraf, O., Slomp, C.P., in't Zandt, M.H., Dolman, A.J. (2018) Methane Feedback to the Global Climate System in a Warmer World. *Reviews of Geophysics*, 56, 207-250.
- Diamond, M.S., Gettelman, A., Lebsock, M.D., McComiskey, A., Russell, L.M., Wood, R., Feingold, G. (2022) To assess marine cloud brightening's technical feasibility, we need to know what to study - and when to stop. *PNAS*, 119, 4.
- Dowell, G., Niederdeppe, J., Vanucchi, J., Dogan, T., Donaghy, K., Jacobson, R., Mahowald, N., Milstein, M., Zelikova, T.J. (2020) Rooting carbon dioxide removal research in the social sciences. *Interface Focus*, 10:20190138.
- Emerson, D. (2019) Biogenic Iron Dust: A Novel Approach to Ocean Iron Fertilization as a Means of Large Scale Removal of Carbon Dioxide From the Atmosphere. *Front. Mar. Sci.* 6:22.

Fischer, H., Behrens, M., Bock, M., Richter, U., Schmitt, J., Loulergue, L., Chappellaz, J., Spahni, R., Blunier, T., Leuenberger, M. and Stocker, T.F. (2008). Changing boreal methane sources and constant biomass burning during the last termination. *Nature*, 452, 7189, 864-867.

Fischer, H., Siggaard-Andersen, M.L., Ruth, U., Rothlisberger, R., Wolff, E. (2007) Glacial/interglacial changes in mineral dust and sea-salt records in polar ice cores: sources, transport, and deposition. *Reviews of Geophysics*, 45, RG1002.

Ferretti, D.F., Miller, J.B., White, J.W., Etheridge, D.M., Lassey, K.R., Lowe, D.C., Meure, C.M., Dreier, M.F., Trudinger, C.M., Van Ommen, T.D., Langenfelds, R.L. (2005) Unexpected changes to the global methane budget over the past 2000 years. *Science*, 309, 1714-7.

Forster, P. et al. The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity. in *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* vol. In Press (Cambridge University Press, 2021).

Gromov, S.S. (2014) Stable isotope composition of atmospheric carbon monoxide: a modeling study. Johannes Gutenberg- Universität, Mainz. Available online at: <https://openscience.ub.uni-mainz.de/handle/20.500.12030/4250> (accessed August 18, 2023).

Han, C., Do Hur, S., Han, Y., Lee, K., Hong, S., Erhardt, T., Fischer, H., Svensson, A.M., Steffensen, J.P., Vallerønga, P. (2018) High-resolution isotopic evidence for a potential Saharan provenance of Greenland glacial dust. *Scientific Reports*, 8, 15582.

Horowitz, H.M., Holmes, C., Wright, A., Sherwen, T., Wang, X., Evans, M., et al. (2020). Effects of sea salt aerosol emissions for marine cloud brightening on atmospheric chemistry: Implications for radiative forcing. *Geophysical Research Letters*, 47.

IPCC, 2023. *Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland, 35-115.

Ito, A. Global modeling study of potentially bioavailable iron input from shipboard aerosol sources to the ocean. (2013) *Global Biogeochemical Cycles*, 27, 1-10.

Jackson, R.B., Abernethy, S., Canadell, J.G., Cargnello, M., Davis, S.J., Feron, S., Fuss, S., Heyer, A.J., Hong, C., Jones, C.D., Matthews, H.D., O'Conner, F.M., Pisciotta, M., Rhoda, H.M., de Richter, R., Solomon, E.I., Wilcox, J.L., Zickfeld, K. (2021) Atmospheric methane removal: a research agenda. *Philosophical Transactions Royal Society A*, 379, 20200454.

Jackson, R.B., Saunio, M., Bousquet, P., Canadell, J.G., Poulter, B., Stavert, A.R., Bergamaschi, P., Niwa, Y., Segers, A., Tsuruta, A. (2020) Increasing anthropogenic methane emissions arise equally from agricultural and fossil fuel sources. *Environmental Research Letters*, 15, 071002.

Kleinen, T., Gromov, S., Steil, B., Brovkin, V. (2021) Atmospheric methane underestimated in future climate projections. *Environmental Research Letters*, 16, 9, 094006.

Legrand, M., Preunkert, S., Wolff, E., Weller, R., Jourdain, B., Wagenbach, D. (2017) Year-round records of bulk and size-segregated aerosol composition in central Antarctica (Concordia site) - Part 1: Fractionation of sea-salt particles. *Atmospheric Chemistry and Physics*, 17, 22, 14,039-14,054.

Lelieveld, J., Peters, W., Dentener, F.J., Krol, M.C. (2002) Stability of tropospheric hydroxyl chemistry. *Journal of Geophysical Research*, 107, D23, 4715.

Lenton, T.M., Held, H., Hall, J.W., Lucht, W., Rahmstorf, S., Schellnhuber, H.J. (2008) Tipping elements in the Earth's climate system. *PNAS*, 105:6, 1786-1793.

Li, Q., Fernandez, R.P., Hossaini, R., Iglesias-Suarez, F., Cuevas, C.A., Apel, E.C., Kinnison, D.E., Lamarque, J.F., Saiz-Lopez, A. (2022) Reactive halogens increase the global methane lifetime and radiative forcing in the 21st century. *Nature Communications*, 13, 2768.

Li, Q., Meidan, D., Hess, P., Anel, J.A., Cuevas, C.A., Doney, S., Fernandez, R.P., van Herpen, M., Hoglund-Isaksson, L., Johnson, M.S., Kinnison, D.E., Lamarque, J.F., Röckmann, T., Mahowald, N.M., Saiz-Lopez, A. (2023) Global environmental implications of atmospheric methane removal through chlorine-mediated chemistry-climate interactions. *Nature Communications*, 14, 4045.

Li, L., Mahowald, N.M., Miller, R.L., Pérez García-Pando, C., Klose, M., Hamilton, D.S., Gonçalves Ageitos, M., Ginoux, P., Balkanski, Y., Green, R.O., Kalashnikova, O., Kok, J.F., Obiso, V., Paynter, D., Thompson, D.R. (2021) Quantifying the range of the dust direct radiative effect due to source mineralogy uncertainty. *Atmospheric Chemistry and Physics*, 21, 3973–4005.

Li, Y., Song, Y., Li, X., Kaskaoutis, D.G., Gholami, H., Li, Y. Disentangling variations of dust concentration in Greenland ice cores over the last glaciatiion: An overview of current knowledge and new initiative (2023) *Earth-Science Reviews*, 242, 104451.

Maesano, C.N., Campbell, J.S., Foteinis, S., Furey, V., Hawrot, O., Pike, D., Aeschlimann, S., Reginato, P.L., Goodwin, D.R., Looger, L.L., Boyden, E.S., Renforth, P. (2022) Geochemical Negative Emissions Technologies: Part II. Roadmap. *Frontiers in Climate*, 4.

Mahowald, N.M., Hamilton, D.S., Mackey, K.R.M., Moore, J.K., Baker, A.R., Scanza, R.A., Zhang, Y. (2018) Aerosol trace metal leaching and impacts on marine microorganisms. *Nature Communications*, 9, 2614.

Mak, J., Kra, G., Sandomenico, T. (2003) The seasonality varying isotopic composition of the sources of carbon monoxide at Barbados, West Indies. *Journal of Geophysical Research*, 108.

Matsui, H., Mahowald, N.M., Moteki, N., Hamilton, D.S., Ohata, S., Yoshida, A., Koike, M., Scanza, R.A., Flanner, M.G. (2018) Anthropogenic combustion iron as a complex climate forcer. *Nature Communications*, 1593.

McKay, D.I.A., Stall, A., Abrams, J.F., Winkelmann, R., Sakschewski, B., Loriani, S., Fetzer, I., Cornell, S.E., Rockstrom, J., Lenton, T.M. (2022) Exceeding 1.5°C global warming could trigger multiple climate tipping points, *Science*, 377, 6611, 7950.

Meidan et al., Evaluating the potential of iron-based interventions in methane reduction and climate mitigation, in preparation for submission to *Environmental Research Letters*.

Ming, T., Li, W., Yuan, Q., Davies, P., de Richter, R., Peng, C., Deng, Q., Yuan, Y., Caillol, S., Zhou, N. (2022) Perspectives on removal of atmospheric methane. *Advances in Applied Energy* 5, 100085.

Neumann, R.B., Moorberg, C.J., Lundquist, J.D., Turner, J.C., Waldrop, M.P., McFarland, J.W., Euskirchen, E.S., Edgar, C.W., Turetsky, M.R. (2019) Warming Effects of Spring Rainfall Increase Methane Emissions From Thawing Permafrost. *Geophysical Research Letters*, 46, 1393-1401.

Nisbet et al. Manuscript responding to referees' comments: submitted to *Global Biogeochemical Cycles*. Authorea. June 11, 2023.

NOAA. Global Monitoring Laboratory, Earth System Research Laboratories. Trends in Atmospheric Methane. (2023) Available online at: https://gml.noaa.gov/ccgg/trends_ch4/ (accessed July 21, 2023).

Ocean Visions (2023). Ocean-Based Carbon Dioxide Removal: Road Maps. Available online at: <https://www2.oceanvisions.org/roadmaps/> (accessed July 21, 2023).

O'Day, P.A., Pattammattel, A., Aronstein, P., Leppert, V.J., Forman, H.J. (2022) Iron Speciation in Respirable Particulate Matter and Implications for Human Health. *Environmental Science and Technology*, 56, 7006-7016.

Oeste, F. D. (2009) Ferrous aerosol emission method for self-releasing cooling of atmosphere, involves adding compound of iron and/or bromine and/or chlorine to solid fuel and/or gas fuel and mixing flue gases of solid fuel and/or gas fuel, DE102009004281A, Bundesrepublik Deutschland Deutsches Patent- und Markenamt. Available online at: <https://patentimages.storage.googleapis.com/ac/64/72/49f3c79762e44f/AU2010203265B2.pdf> (accessed August 23, 2023).

Oeste, F. D., de Richter, R., Ming, T., Caillol, S. (2017) Climate engineering by mimicking natural dust climate control: the iron salt aerosol method. *Earth System Dynamics*, 8, 1–54.

Paudel, R., Mahowald, N.M., Hess, P.G.M., Meng, L., Riley, W.J. (2016) Attribution of changes in global wetland methane emissions from pre-industrial to present using CLM4.5-BGC. *Environmental Research Letters*, 11, 034020.

Pennacchio, L., van Herpen, M., Liisberg, J.B., Meidan, D., Saiz-Lopez, A., Johnson, M.S. Catalytic efficiencies for methane removal via iron chloride photochemistry in plumes: Impact of HO_x, NO_x and the high-chlorine regime, submitted to *Environmental Research Letters*.

Peng, S., Lin, X., Thompson, R.L., Xi, Y., Liu, G., Hauglustaine, D., Lan, X., Poulter, B., Ramonet, M., Saunois, M., Yin, Y., Zhang, Z., Zheng, B., Ciais, P. (2022) Wetland emission and atmospheric sink changes explain methane growth in 2020. *Nature*, 612, 7940, 477–482.

Pio, C.A., Lopes, D.A. Chlorine loss from marine aerosol in a coastal atmosphere. (1998) *Journal of Geophysical Research*, 103, D19, 25,263-25,272.

Prospero, J.M., Delany, A.C., Delany A.C., Carlson, T.N. (2021) The Discovery of African Dust Transport to the Western Hemisphere and the Saharan Air Layer. *Bulletin of the American Meteorological Society*, 102,6, E1239-E1260.

Pye, H.O.T, Nenes, A., Alexander, B., Ault, A.P., Barth, M.C., Clegg, S.L., Collett, J.L., Fahey, K.M., Hennigan, C.J., Herrmann, H., Kanakidou, M., Kelly, J.T., Ku, I.T., Faye McNeill, .V, Riemer, N., Schaefer, T., Shi, G., Tilgner, A., Walker, J.T., Wang, T., Weber, R., Xing, J., Zaveri, R.A., and Zuend, A. (2020) The acidity of atmospheric particles and clouds. *Atmospheric Chemistry and Physics*, 20, 4809–88.

Röckmann, T., Brenninkmeijer, C.A.M., Crutzen, P.J., and Platt, U. (1999) Short term variations in the $^{13}\text{C}/^{12}\text{C}$ ratio of CO as a measure of Cl activation during tropospheric ozone depletion events in the Arctic, *Journal of Geophysical Research*, 104, 1691–1697.

Rodriguez, S., Prospero, J.M., Lopez-Darias, J., Garcia-Alvarez, M.I., Zuidema, P., Nava, S., Lucarelli, F., Gaston, C.J., Galindo, L., Sosa, E. (2021) Tracking the changes of iron solubility and air pollutants traces as African dust transits the Atlantic in the Saharan dust outbreaks. *Atmospheric Environment*, 246, 118092.

Saiz-Lopez, A., Fernandez, R.P., Li, Q., Cuevas, C.A., Fu, X., Kinnison, D.E., Times, S., Mahajan, A.S., Martin, J.C.G., Iglesias-Suarez, F., Hossaini, R., Plane, J.M.C., Myhre, G., Lamarque, J.F. (2023) Natural short-lived halogens exert an indirect cooling effect on climate. *Nature*, 618, 967-973.

Sapart, C.J., Monteil, G., Prokopiou, M., van de Wal, R.S., Kaplan, J.O., Sperlich, P., Krumhardt, K.M., Van der Veen, C., Houweling, S., Krol, M.C. and Blunier, T. (2012) Natural and anthropogenic variations in methane sources during the past two millennia. *Nature*, 490, 85-88.

Saunois et al., The Global Methane Budget 2000–2017 (2020) *Earth Systems Science Data*, 12, 3, 1561–1623.

Scott-Buechler, C., Jeanty, J., Fraser, C., Adcox, G., Scott, C. (2023) Advancing Equitable Deployment of Regional DAC Hubs. *Data For Progress*. Available online at: <https://www.filesforprogress.org/memos/advancing-equitable-deployment-of-regional-dac-hubs.pdf> (accessed August 15, 2023).

Seinfeld, J. H., & Pandis, S. N. (2016). *Atmospheric chemistry and physics: from air pollution to climate change*. John Wiley & Sons.

Spark (2023). *Approaches to Atmospheric Methane Removal*. Available online at: <https://www.sparkclimate.org/methane-removal/101/approaches> (accessed July 21, 2023).

Staniaszek, A., Griffiths, P.T., Folberth, G.A., O'Connor, F.M., Abraham, N.L., Archibald, A.T. (2022) The role of future anthropogenic methane emission in air quality and climate. *NPJ Climate and Atmosphere Science*, 5, 21.

The Arctic Institute (2021). Sami Council resistance to SCoPEX highlights the complex questions surrounding geoengineering and consent. Available online at: <https://www.thearcticinstitute.org/sami-council-resistance-scopex-highlights-complex-questions-geo-engineering-consent/> (accessed July 21, 2023).

UNFAO (2016); United Nations, Food and Agriculture Organization. Free Prior and Informed Consent: An indigenous peoples' right and a good practice for local communities. Available online at: <https://www.fao.org/3/i6190e/i6190e.pdf> (accessed July 21, 2023).

UNEP (2021); United Nations Environment Programme. Global Methane Assessment: Benefits and Costs of Mitigating Methane Emissions. Available online at: <https://www.unep.org/resources/report/global-methane-assessment-benefits-and-costs-mitigating-methane-emissions> (accessed July 21, 2023).

van Herpen, M.M.J.W., Li, Q., Saiz-Lopez, A., Liisberg, J.B., Röckmann, T., Cuevas, C.A., Fernandez, R.P., Mak, J.E., Mahowald, N.M., Hess, P., Meidan, D., Stuut, J.-B. W., Johnson, M.S. (2023) Photocatalytic Chlorine Atom Production on Mineral Dust-Sea Spray Aerosols over North Atlantic, *Proceedings of the National Academy of Sciences of the United States of America*, 120, 31.

Wang, X., Jacob, D.J., Eastham, S.D., Sulprizio, M.P., Zhu, L., Chen, Q., Alexander, B., Sherwen, T., Evans, M.J., Lee, B.H., Haskins, J.D., Lopez-Hilfiker, F.D., Thornton, J.A., Huey, G.L., Liao, H. (2019) The role of chlorine in global tropospheric chemistry. *Atmospheric Chemistry and Physics*, 19, 6, 3981-4003.

Wanser, K., Doherty, S.J., Hurrell, J.W., Wong, A. (2022) Near-term climate risks and sunlight reflection modification: a roadmap approach for physical science research. *Climate Change*, 174:23.

Wittmer, J., Bleicher, S., Ofner, J., Zetzsch, C. (2015a) Iron (III)-induced activation of chloride from artificial sea-salt aerosol. *Environmental Chemistry*, 12, 461-475.

Wittmer, J., Bleicher, S., Zetzsch, C. (2015b) Iron (III)-induced activation of chloride and bromide from modeled salt pans. *The Journal of Physical Chemistry A*, 119, 19, 4373-4385.

Wittmer, J., Zetzsch, C. (2017) Photochemical activation of chlorine by iron oxide aerosol. *Journal of Atmospheric Chemistry*, 74, 187-204.

Yuan, T., Yu, H., Chin, M., Remer, L.A., McGee, D., Evan, A. (2020) Anthropogenic decline of African dust: Insights from the Holocene records and beyond. *Geophysical Research Letters*, 47, 22.

Zhang, Z., Poulter, B., Feldman, A.F., Ying, Q., Ciais, P., Peng, S., Li, X. (2023) Recent intensification of wetland methane feedback. *Nature Climate Change*, 13, 430-433.

Zhu, X. R., Prospero, J. M., Millero, F. J. (1997) Diel variability of soluble Fe(II) and soluble total Fe in North African dust in the trade winds at Barbados, *Journal of Geophysical Research*, 102(D17), 21297– 21305.